



Planck's constant



Francesco Gonnella
Matteo Mascolo




Aristotelian mechanics




The speed of an object's motion is proportional to the force being applied.

$$F = mv$$


Aristotle's trajectories



Issues of Aristotelian mechanics




- It hardly explains how an arrow keeps on flying after being shot;



- Can be easily disproved through a simple experiment.


The scientific method

- Observation of phenomenon;
- Hypothesis and prediction;
- Experiment;
 - Reproducible;
- Write a law using the “language of mathematics”.



Falsifiability of the law

**1874: The task of physics is
“nearly completed”**



Quantum mechanics and relativity




- Is classical mechanics as wrong as Aristotelian mechanics?
- How can classical mechanics, relativity, and quantum mechanics to reconcile?

**Every theory has its own scope of validity,
one has to know its limits!**


For example, how is the atom made?

The plum pudding model of the atom by J.J. Thomson




The atoms [...] consist of a number of negatively electrified corpuscles enclosed in a sphere of uniform positive electrification.

Rutherford's model of the atom




Negatively-charged electrons orbit around a positively charged nucleus.

Geiger–Marsden experiment 1911




©NCSSM 2002

Scattering Helium nuclei (aka α -particles) off thin gold foil.



The more the particle travels close to the centre, the smaller is the deflection angle



The more the particle travels close to the centre, the larger is the deflection angle. Even backscattering occurs!

Some calculations



$$E = T + V = \frac{1}{2}mv^2 + \frac{1}{4\pi\epsilon_0} \frac{q_1q_2}{r}$$

The total energy is the kinetic energy of the α -particle T plus the Coulomb-energy V due to the repulsion between charges of the same sign. When the particle turns back $T=V$.

Putting $T=V$ we get:
$$T = \frac{1}{4\pi\epsilon_0} \frac{q_1q_2}{r}$$

If we solve for r we get:
$$r = \frac{1}{4\pi\epsilon_0} \frac{q_1q_2}{T}$$

Some more calculations



The elementary charge e , the dielectric constant of vacuum ϵ_0 and the kinetic energy T appear in the previous formula:

$$e = 1.6 \cdot 10^{-19} \text{ C}$$

$$\epsilon_0 = 8.85 \cdot 10^{-12} \text{ m}^{-3} \text{ kg}^{-1} \text{ s}^4 \text{ A}^2$$

$$T \simeq 5 \text{ MeV} = 5 \cdot 10^6 \cdot 1.6 \cdot 10^{-19} \text{ J}$$

It is useful to know that $1/(4\pi\epsilon_0)$ is about $9 \cdot 10^9$ in the International System of Units.

Last calculations

Protons in
 α -particle

Protons in
gold

Elementary charge

$$r = \frac{1}{a \pi \epsilon_0} \frac{Z_\alpha Z_{Au} e^2}{T} = 9 \cdot 10^9 \frac{2 \cdot 79 \cdot 1.6 \cdot 10^{19} \cdot 1.6 \cdot 10^{19}}{5 \cdot 10^6 \cdot 1.6 \cdot 10^{19}} =$$


Dielectric
constant
of vacuum

Kinetic energy of α -particle

$$r = 4.55 \cdot 10^{-14} m \simeq 45 fm$$

At that time, the radius of the atom of gold was well known: 150'000 fm. **It is 3000 times larger!**

“It was quite the most incredible event that has ever happened to me in my life. It was almost as if you fired a **15-inch shell into a piece of tissue paper and it came back and hit you.**”




On consideration, I realized that this scattering backward must be the result of a single collision, and when I made calculations I saw that it was impossible to get anything of that order of magnitude unless you took a system in which the greater part of the mass of the atom was concentrated in a minute nucleus. It was then that I had the idea of an **atom with a minute massive center, carrying a charge.**

Boltzmann proposes a statistical interpretation of thermodynamics




At the beginning of 1800, Boltzmann proposed a weird theory to explain laws of thermodynamics.




He said that gases were made of many tiny particles moving randomly and hitting one another. The well-known laws of thermodynamics could be interpreted as statistical laws.

People did not like this theory...

Kirchhoff: the black body radiation



A **black body** is a cavity (e.g. an oven) with a tiny hole, kept at a constant temperature.




The radiation entering the black body is reflected by the inner walls a huge number of times before getting out.

Sunday, October 7, 1900
teatime.



Max Planck solves the puzzle of the spectrum of black body radiation. In its calculations, Planck simply replaced an **integral** with a **sum**, in order to get things right. The implications of this “mathematical trick” revealed a bizarre behaviour of nature.

Sum vs. integral:



Planck's law explains the black-body spectrum even at short wavelengths (small λ). The Rayleigh-Jeans law goes to infinity when λ goes to zero.

What was Planck's idea?



Planck assumed that the inner walls of the black body behaved like many tiny oscillators.

They absorbed and emitted electromagnetic waves at **all wavelengths and all energies**.

From this hypothesis, he calculated the **mean energy** $\langle E \rangle$ emitted by each oscillator, “multiplied by” (actually integrated over) the number of oscillators.

Max Planck's calculation (1/2)



Planck disagreed with atomic theories, and did not like statistical thermodynamics...

Nevertheless, Planck took his start from Boltzmann's statistical approach:

$$P(E)dE = \frac{e^{-\frac{E}{kT}}}{\int_0^\infty e^{-\frac{E}{kT}} dE} dE$$

Diagram illustrating the components of the Boltzmann distribution formula:

- Probability of finding a particle with energy between E and $E+dE$** (points to $P(E)dE$)
- Distribution** (points to $e^{-\frac{E}{kT}}$)
- Normalization** (points to the denominator $\int_0^\infty e^{-\frac{E}{kT}} dE$)

In the same fashion as in thermodynamics, this equation provides the probability to find a particle of energy E within a gas at temperature T .

How much do you understand this formula?

Max Planck's calculation (2/2)



The mean value of a distribution is given by its **integral** “weighted” with the value of the energy. This mean value is $\langle E \rangle = kT$.

Using this results, one obtains the Rayleigh-Jeans' black-body spectrum.

$$\langle E \rangle = \frac{\int_0^\infty E e^{-\frac{E}{kT}} dE}{\int_0^\infty e^{-\frac{E}{kT}} dE} = \dots = kT \quad \Rightarrow \quad \rho(\lambda) = \frac{8\pi}{\lambda^4} KT$$

Planck used a sum instead. To do that he had to replace with . So the energy is no longer a continuous variable , but rather times the value .

$$\langle E \rangle_{Planck} = \frac{\sum_0^\infty nh\nu e^{-\frac{nh\nu}{kT}}}{\sum_0^\infty e^{-\frac{nh\nu}{kT}}} = \dots = \frac{hc}{\lambda} \frac{1}{e^{\frac{hc}{\lambda kT}} - 1} \Rightarrow \rho(\lambda) = \frac{8\pi hc}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda kT}} - 1}$$

“Quantum” of energy



It would seem that not any amount of energy can be emitted and absorbed. Only certain discrete values are permitted by nature: the multiples of the so-called “**Quanta**” (one quantum, two quanta, ...)

$$E = h\nu$$

$$h = 6.62618 \cdot 10^{-34} \text{ Js}$$

Bohr's atom (1913)




A charge acceleratin within an electric field emits radiation and loses some enrgy.

So why electrons do not fall over the nucleus?

$$F_c = F_E \rightarrow \frac{mv^2}{r} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q^2}{r^2}$$

$$\frac{m^2 v^2 r^2}{r m r^2} = \frac{L^2}{m r^3} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q^2}{r^2} \rightarrow r = L^2 \cdot \frac{4\pi\epsilon_0}{m q^2}$$




What would we obtain if L were quantized ($L = n \frac{h}{2\pi} = n\hbar$)?

Clearly the radius would be quantized as well:

$$r = n^2 \hbar^2 \cdot 4 \frac{\pi\epsilon_0}{m q^2}$$

Discrete atom spectra



The quantization of angular momentum explains also the well known and yet inexplicable lines in the emission spectra of atoms.


The Compton effect

The Compton effect is the result of a high-energy radiation hitting an electron. The scattered radiation experiences a **wavelength shift** that cannot be explained in terms of classical wave theory, thus lending support to

Einstein's photon theory.

The effect was first demonstrated in 1923 by

Arthur Holly Compton
(Nobel Prize in 1927).



De Broglie's hypothesis

- For a photon, the following formulas hold:


$$E = h\nu \rightarrow p = \frac{h\nu}{c} = \frac{h}{\lambda}$$

- Louis De Broglie (Nobel in 1929) stated that a material particle should behave in the same way. It's only because of the small value of h and the small dimension of particles that **we don't experience quantum mechanics** in everyday life
- In the end if an **electron is a wave**, what is its wavelength?

$$\lambda = \frac{h}{p} = \frac{h}{mv}$$

- This means that an electron can only fit in an orbit if it has a circumference being a multiple of the “**electron wavelength**” $n\lambda = 2\pi r$:

$$L = rmv = r\mathbf{p} = \frac{n\lambda}{2\pi} \cdot \frac{h}{\lambda} = \frac{nh}{2\pi} = n\hbar$$



Conclusion

Quantum mechanics describes a lot of mysterious behaviours of nature.



There is no classical way to describe phenomena like: black-body spectrum, atom discrete spectra, Compton effect, photoelectric effect, double-slit experiment, and much more...

But how can it be like that?

Nobody knows how it can be like that.



Thanks for your attention



s t u d e n t s
a g e
e s **infn-lnf**

