

Evidence for a kaon-bound state K^-pp produced in K^- absorption reactions at rest

M. Agnello,^{1,2} G. Beer,³ L. Benussi,⁴ M. Bertani,⁴ S. Bianco,⁴ E. Botta,^{5,2} T. Bressani,^{5,2} L. Busso,^{6,2} D. Calvo,² P. Camerini,^{7,8} P. Cerello,² B. Dalena,^{9,10} F. De Mori,^{5,2} G. D'Erasmus,^{9,10} D. Di Santo,^{9,10} F. L. Fabbri,⁴ D. Faso,^{6,2} A. Feliciello,² A. Filippi,² V. Filippini,^{11,*} E. M. Fiore,^{9,10} H. Fujioka,¹² P. Gianotti,⁴ N. Grion,⁸ V. Lucherini,⁴ S. Marcello,^{5,2} T. Maruta,¹² N. Mirfakhrai,¹³ O. Morra,¹⁴ T. Nagae,¹⁵ A. Olin,¹⁶ H. Ota,¹⁷ E. Pace,⁴ M. Palomba,^{9,10} A. Pantaleo,¹⁰ A. Panzarasa,¹¹ V. Paticchio,¹⁰ S. Piano,^{7,8} F. Pompili,⁴ R. Rui,^{7,8} G. Simonetti,^{9,10} H. So,¹⁸ S. Tomassini,⁴ A. Toyoda,¹⁵ R. Wheadon,² and A. Zenoni^{19,11}

(FINUDA collaboration)

¹*Dipartimento di Fisica del Politecnico di Torino, I-10129 Torino, Italy*

²*INFN Sezione di Torino, I-10125 Torino, Italy*

³*University of Victoria, Victoria, Canada*

⁴*INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*

⁵*Dipartimento di Fisica Sperimentale, Università di Torino, I-10125 Torino, Italy*

⁶*Dipartimento di Fisica Generale, Università di Torino, I-10125 Torino, Italy*

⁷*Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy*

⁸*INFN Sezione di Trieste, I-34127 Trieste, Italy*

⁹*Dipartimento di Fisica, Università di Bari, I-70126 Bari, Italy*

¹⁰*INFN Sezione di Bari, I-70126 Bari, Italy*

¹¹*INFN Sezione di Pavia, I-27100 Pavia, Italy*

¹²*Department of Physics, University of Tokyo, Bunkyo, Tokyo, 113-0033, Japan*

¹³*Department of Physics, Shahid Beheshti University, 19834 Teheran, Iran*

¹⁴*CNR-IFSI Sezione di Torino and INFN Sezione di Torino, I-10125 Torino, Italy*

¹⁵*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki, 305-0801, Japan*

¹⁶*TRIUMF, Vancouver, BC, V6T 2A3, Canada*

¹⁷*RIKEN, Wako, Saitama, 351-0198, Japan*

¹⁸*Department of Physics, Seoul National University, Seoul, 151-742, South Korea*

¹⁹*Dipartimento di Meccanica, Università di Brescia, 25123 Brescia, Italy*

(Dated: March 4, 2005)

We have searched for a deeply-bound kaonic state by using the FINUDA spectrometer installed at the e^+e^- collider DAΦNE. Almost monochromatic K^- 's produced through the decay of $\phi(1020)$ mesons are used to observe K^- absorption reactions stopped on very thin nuclear targets. Taking this unique advantage, we have succeeded to detect a kaon-bound state K^-pp through its two-body decay into a Λ hyperon and a proton. The binding energy and the decay width are determined from the invariant mass distribution as 115_{-5}^{+6} (stat.) $_{-4}^{+3}$ (sys.) MeV and 67_{-11}^{+14} (stat.) $_{-3}^{+2}$ (sys.) MeV, respectively.

PACS numbers: 21.30.Fe, 21.45.+v, 21.80.+a, 25.80.Nv, 36.10.Gv, 26.60.+c

At the last conference on hypernuclei and strangeness production HYP2003, two groups reported possible signatures of deeply-bound kaonic states in two different reactions: $^{16}\text{O}(K^-, n)$ [1] and $^4\text{He}(K^-_{stop}, nX)$ [2, 3]. The observed signals were interpreted as the productions of $^{15}_{K^-}\text{O}$ and K^-ppn , respectively. Later, the latter group has also reported the observation of a much clearer signal of the production of a tri-baryonic state $S^0(3115)$ in the $^4\text{He}(K^-_{stop}, pX)$ reaction [3, 4]. If these interpretations are true, the K^- nucleus potential seems to be deeper than 100 MeV.

Thus, it is an urgent experimental subject to verify or confirm if these signals really correspond to the production of kaon bound states or not. Unfortunately, all of these measurements used missing-mass method, so that there exist some ambiguities whether a kaon was really bound in the system. In the FINUDA (acronym for "Fisica NUcleare a DAΦNE") experiment, we can stop

K^- 's in very thin nuclear targets, and detect multi-tracks in the FINUDA spectrometer. It enables us to detect a Λ hyperon through its decay to proton and π^- . We can clearly identify the formation of bound states with strangeness from their decays to $\Lambda + X$.

As for the K^-p interaction in free space, the existing experimental information on low-energy K^-p scattering [5], K^-p threshold branching ratios [6], a kaonic-hydrogen atomic X-ray measurement [7, 8], etc. leaves rooms for different theoretical analyses.

The information on K^- nucleus interaction has been obtained with the kaonic atomic data [9]. Unfortunately, we have qualitatively two different predictions for the depth of the K^- nucleus potential at normal nuclear matter density: very deep attractive potentials ($-\text{Re } V_{opt}(\rho_0) \approx 150\text{--}200$ MeV) [10, 11] and much shallower potentials ($-\text{Re } V_{opt}(\rho_0) \approx 50\text{--}75$ MeV) [12–16]. Both types of potentials reasonably describe the shifts

and widths of the X-ray data.

Recently, an interesting idea has been proposed suggesting the possible existence of deeply bound nuclear \bar{K} states in light nuclei [17, 18], for which the potential is very deep. Since the potential is strongly attractive in the $I=0$ $\bar{K}N$ interaction, proton-rich \bar{K} -nuclei such as K^-pp and K^-ppn are predicted to have large binding energies of 50–100 MeV. A further interesting suggestion is that the nucleus might be shrunk because of the large binding energy and form a high-density state, which is also confirmed with the method of anti-symmetrized molecular dynamics for the cases of K^-ppn and $K^-^8\text{Be}$ [19].

If such a \bar{K} -nucleus is observed, it would give us direct information of the K^- nucleus potential at nuclear matter density. Another proposed idea to get the information of the K^- nucleus potential is to study the (K^-_{stop}, π^-) reaction into specific Λ -hypernuclear states [12].

The experiment is performed at an intersection region of the e^+e^- collider DAΦNE in Laboratori Nazionali di Frascati (LNF), which produces copious $\phi(1020)$ mesons. A key characteristic of the FINUDA spectrometer is the achievement of high resolution through the use of very thin nuclear targets of ≈ 200 mg/cm² and low mass chambers in a He atmosphere [20]. The nearly monochromatic K^- from the ϕ decays are fully stopped in these targets. The spectrometer covers a solid angle larger than 2π sr. This good performance enables us to carry out the spectroscopy of Λ -hypernuclei with a good energy resolution better than 1 MeV FWHM and their weak decay studies as the main objective of the FINUDA experiment.

Figure 1 (TOP) shows an overview of the FINUDA spectrometer. It consists of three regions: the interaction/target region, the tracking system and the external time-of-flight system. The K^+K^- pair from the ϕ decay is detected by a barrel of 12 thin scintillator slabs (TOFINO) and an octagonal array of silicon microstrip detectors (ISIM). The kaon loses its energy through the beam pipe, TOFINO and ISIM, and finally stops in one of thin targets. After K^- absorption in the target, the emitted particles are detected by the external tracking system. Charged particles are detected with a second layer of silicon microstrip detectors (OSIM), two layers of low-mass drift chambers (LMDC), and six layers of aluminized mylar straw tubes, arranged in three groups, the first one beam-oriented, the others tilted by $\pm 12^\circ$. The dE/dx information obtained in the OSIM detector is used for particle identification. 72 external scintillator slabs (TOFONE) are installed in the outermost part, which are used to measure the time-of-flight of a charged or neutral particle and for triggering purpose.

In the first data taking in 2003–2004, five kinds of targets (^6Li , ^7Li , ^{12}C , ^{27}Al and ^{51}V) were installed as shown in Fig. 1 (BOTTOM). The integrated luminosity during the run reached ~ 250 pb⁻¹, of which ~ 190 pb⁻¹ was effectively used in the present analyses.




FIG. 1: Global views (top: complete apparatus, bottom: interaction/target region) of the FINUDA apparatus.

The momentum resolution of the FINUDA spectrometer at this stage of analysis was determined from the two monochromatic peaks corresponding to $K_{\mu 2}(K^+ \rightarrow \mu^+\nu_\mu)$ and $K_{\pi 2}(K^+ \rightarrow \pi^+\pi^0)$ decays of K^+ stopping in the targets. It was found to be $\Delta p/p \sim 0.6\%$ FWHM. It is remarkable that the spectrometer delivered the right momenta of the monochromatic particles to a precision better than 200 keV/c, without the need of further correction.

Λ hyperons are produced with a sizable fraction in kaon absorption through various processes. The quasifree process ($K^-N \rightarrow \Lambda\pi$) emits a slow Λ with a momentum ~ 300 MeV/c. Above ~ 400 MeV/c, a main contribution comes from the Λ hyperons emitted from two-nucleon absorptions ($K^- + \text{“}NN\text{”} \rightarrow \Lambda N, \Sigma^0 N$). The acceptance of the FINUDA spectrometer cuts the Λ hyperons with a momentum lower than 300 MeV/c which is restricted by the low momentum threshold for π^- from the $\Lambda \rightarrow p + \pi^-$ decay. Therefore, the Λ from the quasifree process is hardly observed in the FINUDA spectrometer.

Λ particles can be identified by reconstructing the invariant mass of a proton and a negative pion as shown in Fig. 2(a). The peak position agrees well with the known Λ mass, and the width of the peak is as narrow as 6 MeV/c² FWHM.




FIG. 2: (a) Invariant mass distribution of a proton and a π^- for all the events in which these two particles are observed, fitted by a single Gaussian together with a linear background in the invariant-mass range of 1100–1130 MeV/c^2 . (b) Opening angle distribution between a Λ and a proton. (solid line : ${}^6\text{Li}$, ${}^7\text{Li}$ and ${}^{12}\text{C}$, dashed line : ${}^{27}\text{Al}$ and ${}^{51}\text{V}$). The shaded area ($\cos\theta^{\text{Lab.}} < -0.8$) is selected as the back-to-back events.

When a K^- interacts with two protons, one expects that a hyperon-nucleon pair ($\Lambda + p$, $\Sigma^0 + p$ or $\Sigma^+ + n$) is emitted in the opposite direction, ignoring a final state interaction inside the nucleus. The angular correlation between a Λ and a proton from the same point in the target [Fig. 2(b)] clearly indicates the existence of this kind of reaction. Even for heavy nuclei such as ${}^{27}\text{Al}$ and ${}^{51}\text{V}$ the similar correlations were observed although less pronounced as compared to those for light nuclei.

In the following analysis, we use the Λ - p pairs emitted in the opposite direction ($\cos\theta^{\text{Lab.}} < -0.8$) only from the light nuclear targets (${}^6\text{Li}$, ${}^7\text{Li}$ and ${}^{12}\text{C}$). In this way, the possible contaminations from other processes can be minimized, and the signal-to-noise ratio as well as the available statistics are kept to a good level.

Since the back-to-back angular correlation between a Λ and a proton is so clear, it is naturally expected that the two particles are emitted from a “ K^-pp ” intermediate system. If the reaction process were simply two-nucleon absorption process, the mass of the system should be close to the sum of a kaon and two proton mass, namely $2.370 \text{ GeV}/c^2$.




FIG. 3: Invariant mass of a Λ and a proton in back-to-back correlation ($\cos\theta^{\text{Lab.}} < -0.8$) from light targets before the acceptance correction. The inset shows the result after the acceptance correction for the events which have two protons with well-defined good tracks. Only the bins between 2.22–2.33 GeV/c^2 are used for the fitting.

The invariant mass distribution of the Λ - p pairs is shown in Fig. 3. A significant mass decrease of the K^-pp system with respect to its expected mass is observed. It can be interpreted as a bound state composed of a kaon and two protons; hereafter abbreviated as K^-pp .

In the inset in Fig. 3, the acceptance corrected invariant mass distribution for events with two well-defined long-track protons is shown. Since the trigger and detection acceptance are monotonically increasing functions of the invariant mass in this mass region, the peak further shifts to a lower mass side. The binding energy $B_{K^-pp} = 115_{-5}^{+6}(\text{stat.})_{-4}^{+3}(\text{sys.}) \text{ MeV}$ and the width $\Gamma = 67_{-11}^{+14}(\text{stat.})_{-3}^{+2}(\text{sys.}) \text{ MeV}$ are obtained from the fitting with a Lorentzian function (folded with a Gaussian with $\sigma = 4 \text{ MeV}/c^2$, corresponding to the detector resolution, estimated with a Monte Carlo simulation) in the region of 2.22–2.33 GeV/c^2 . Although we still have ambiguities on absolute normalization, a rough estimate on the yield of $K^-pp \rightarrow \Lambda + p$ is of the order of 0.1% per stopped K^- .

The K^-pp decay into $\Sigma^0 + p$ should still have a back-to-back correlation between a Λ and a proton, but a γ from the Σ^0 decay is missing when reconstructing the invariant mass (Σ^0 at rest emits a 74 MeV γ). An excess in a low-mass region below $\sim 2.2 \text{ GeV}/c^2$ may be attributed to this decay mode. An analysis of this decay mode is under way for future publication.

From old bubble chamber and emulsion data [21] it is known that kaon two-nucleon absorption processes take place in 15–20% per stopped K^- in the broad range of the periodic table. However, the experimental information was statistically too limited to further investigate the reaction mechanism, and nobody has identified

the kaon two-nucleon absorption process experimentally. Several theoretical analyses were performed in the Sixties to understand the reaction mechanism. A more recent theoretical analysis [22] roughly explained it with two-nucleon absorption process including meson-rescattering diagrams. Here, it should be noted that the main mode for the kaon two-nucleon absorption is on a pn pair with $S=1$ and $T=0$ and not on a pp pair, which only contributes by $\sim 10\%$ according to the calculation. Thus, there is no a priori reason that the two-nucleon absorption are to be observed in the Λ - p coincidence events.

On the other hand, the detector system is very sensitive to the existence of the two-nucleon absorption mode $K^- + "pp" \rightarrow \Lambda p$ since its invariant mass resolution is $10 \text{ MeV}/c^2$ FWHM. The effect of the nuclear binding of two protons is only to move the peak position to the lower mass side of the order of separation energies of two protons ($\sim 30 \text{ MeV}$), and not to broaden the peak. A sharp spike around $2.34 \text{ GeV}/c^2$ may be attributed to this process. There could be, in addition, the two-nucleon absorption mode $K^- + "pp" \rightarrow \Sigma^0 p$. In this case, the Λ - p invariant mass distribution is shifted to the lower mass region by about 74 MeV and broadened because a γ from the $\Sigma^0 \rightarrow \Lambda + \gamma$ decay is missing. However, the observed invariant mass distribution is too broad to be attributed to this process only. Also, according to the old data in helium bubble chamber [23], the branching ratios of $K^- NN \rightarrow \Lambda N$ and $\Sigma^0 N$ are estimated to be $9.3 \pm 2.6\%$ and $2.3 \pm 1.0\%$ per stopped K^- in ^4He , respectively. The theoretical calculation [22] also suggests the ΛN mode has a larger branching ratio than the $\Sigma^0 N$ mode. Therefore, we can assume that the kaon two-nucleon absorption mode is not dominant in this channel, nor in the $\Sigma^0 + p$ one which has even a lower branching ratio.

Thus, we attribute the observed events in the bound region to the formation of the deeply-bound $K^- pp$ state. Since the $K^- pp$ mass that we have obtained is so far above the $\Lambda + p$ threshold, the $\Lambda + p + \pi$ decay would be possible. However, it would contribute to the non back-to-back events in the opening angular distribution of Λ - p pairs. From the data, it seems to be a small contribution. With the present invariant mass precision, it is not clear that the state is below the $\Sigma\pi$ thresholds.

In summary, we have successfully observed evidence of the kaon bound state $K^- pp$ through its decay into a Λ and a proton. The invariant mass distribution of the Λ - p pair shows a significant mass decrease with respect to the mass of the system expected in the case of a simple kaon two-nucleon absorption. The state $K^- pp$ has a binding energy of 115^{+6}_{-5} (stat.) $^{+3}_{-4}$ (sys.) MeV and a decay width of 67^{+14}_{-11} (stat.) $^{+2}_{-3}$ (sys.) MeV.

We are greatly indebted with Dr. S. Bertolucci, former

Director of LNF, for his continuous encouragement and help. Dr. P. Raimondi and the DAΦNE machine staff are warmly acknowledged for their very skillful handling of the DAΦNE collider.

This work is supported in part by the Grant-In-Aid for Scientific Research (B) 15340087 of Ministry of Education, Culture, Sports, Science and Technology (MEXT).

* deceased

- [1] T. Kishimoto *et al.*, to be published in Proceedings of HYP2003, Newport News, USA, November 2003.
- [2] M. Iwasaki *et al.*, nucl-ex/0310018.
- [3] T. Suzuki *et al.*, to be published in Proceedings of HYP2003, Newport News, USA, November 2003.
- [4] T. Suzuki *et al.*, Phys. Lett. B **597**, 263 (2004).
- [5] A. D. Martin, Nucl. Phys. B **179**, 33 (1981); J. Ciborowski *et al.*, J. Phys. G **8**, 13 (1982); D. Evans *et al.*, *ibid.* **9**, 885 (1983).
- [6] D. N. Tovee *et al.*, Nucl. Phys. B **33**, 493 (1971); R. J. Nowak *et al.*, *ibid.* **139**, 61 (1978); W. E. Humphrey and R. R. Ross, Phys. Rev. **127**, 1305 (1962).
- [7] M. Iwasaki *et al.*, Phys. Rev. Lett. **78**, 3067 (1997); T. M. Ito *et al.*, Phys. Rev. C **58**, 2366 (1998).
- [8] J. Marton, Acta Physica Slovaca **55**, 69 (2005).
- [9] C. J. Batty, E. Friedman, and A. Gal, Phys. Rep. **287**, 385 (1997).
- [10] E. Friedman, A. Gal, J. Mareš, and A. Cieplý, Phys. Rev. C **60**, 024314 (1999).
- [11] E. Friedman, A. Gal, and C.J. Batty, Nucl. Phys. A **579**, 518 (1994).
- [12] A. Cieplý, E. Friedman, A. Gal, and J. Mareš, Nucl. Phys. A **696**, 173 (2001).
- [13] N. Kaizer, P.B. Siegel, and W. Weise, Nucl. Phys. A **594**, 325 (1995).
- [14] T. Waas, N. Kaiser, and W. Weise, Phys. Lett. B **365**, 12 (1996); *ibid* **379**, 34 (1996).
- [15] A. Ramos and E. Oset, Nucl. Phys. A **671**, 481 (2000).
- [16] A. Baca, C. García-Recio, and J. Nieves, Nucl. Phys. A **673**, 335 (2000).
- [17] Y. Akaishi and T. Yamazaki, Phys. Rev. C **65**, 044005 (2002).
- [18] T. Yamazaki and Y. Akaishi, Phys. Lett. B **535**, 70 (2002).
- [19] A. Doté, H. Horiuchi, Y. Akaishi, and T. Yamazaki, Phys. Lett. B **590**, 51 (2004).
- [20] T. Bressani in *Proceedings of Workshop on Physics and Detectors for DAΦNE, Frascati, April 9-12, 1991*, edited by G. Pancheri (Laboratori Nazionali di Frascati), p. 475.
- [21] V. Vander Velde-Wilquet *et al.*, Nuovo Cimento **39A**, 537 (1977).
- [22] T. Onaga, H. Narumi, and T. Kohmura, Prog. Theor. Phys. **82**, 222 (1989).
- [23] P. A. Katz *et al.*, Phys. Rev. D **1**, 1267 (1970).




Figure 1a