

KAONIS

M. Bazzi, M. Cargnelli (Ric. Str.), A. Clozza, C. Curceanu (Resp. Naz.),
R. Del Grande (Ass.), L. De Paolis (Dott.), S. Dabagov, C. Guaraldo,
D. Hampai, P. Levi Sandri, M. Iliescu (Ric. Str.), M. Merafina (Assoc.),
M. Miliucci (Dott.), S. Niedźwiecki (Ric. Str.), E. Pace,
A. Scordo, D. Sirghi (Ass.), F. Sirghi (Ass.), M. Skurzok (Post. Doc.), A. Spallone (Assoc.),
O.Vazquez Doce (Ric. Str.), J. Zmeskal (Ric. Str.)

1 KAONNIS: the scientific program

KAONNIS represents an integrated initiative in the field of experimental low-energy kaon-nucleon/nuclei interaction studies. Under KAONNIS the following activities are performed:

- the study of kaonic atoms by the SIDDHARTA-2 experiment
- the study of kaon-nuclei interaction at low-energies in the framework of the AMADEUS Collaboration
- participation at experiments at J-PARC (Japan) dedicated to strangeness studies.

We present in what follows these scientific lines, together with the 2019 activities and plans for 2020.

The KAONNIS activities, in particular the collaboration with Japan, are partially financed within the “Strange Matter” project by MAECI and by the STRONG-2020 European project (grant agreement No. 824093).

2 The SIDDHARTA-2 experiment

The objective of the SIDDHARTA-2 (Silicon Drift Detector for Hadronic Atom Research by Timing Application) experiment is to perform high precision measurements of X-ray transitions in exotic (kaonic) atoms at the DAΦNE collider.

The precise measurement of the shift and width of the $1s$ level, with respect to the purely electromagnetic calculated values, in kaonic hydrogen and kaonic deuterium, induced by the strong interaction, through the measurement of the X-ray transitions to this level, will allow the first precise experimental determination of the isospin dependent antikaon-nucleon scattering lengths, fundamental quantities for the understanding of the low-energy QCD in strangeness sector.

The accurate determination of the scattering lengths will place strong constraints on the low-energy K^-N dynamics, which, in turn, constraints the $SU(3)$ description of chiral symmetry breaking in systems containing the strange quark. The implications go from particle and nuclear physics to astrophysics (the equation of state of neutron stars).

In 2009 SIDDHARTA performed the most precise measurement of kaonic hydrogen and the first exploratory study of kaonic deuterium. Moreover, the kaonic helium 4 and 3 transitions to the $2p$ level were measured, for the first time in gas in $He4$ and for the first time ever in $He3$. Presently, a major upgrade of SIDDHARTA, namely SIDDHARTA-2, is under way, with the aim to measure kaonic deuterium and other types of kaonic atoms starting in 2020.

2.1 Previous to SIDDHARTA-2: the SIDDHARTA setup

In the first decade of this century, SIDDHARTA represented a new phase in the study of kaonic atoms at DAΦNE. The previous DEAR experiment's precision was limited by a signal/background ratio of about 1/70 for the kaonic hydrogen measurement, due to the high machine background. To significantly improve this ratio, an experimental breakthrough was necessary. An accurate study of the background sources at DAΦNE was done. The background includes two main sources:

- *synchronous background*: coming from the K^- interactions in the setup materials and ϕ -decay processes; it can be defined as *hadronic background*;
- *asynchronous background*: final products of electromagnetic showers in the machine pipe and in the setup materials, originating from particles lost from primary circulating beams either due to the interaction of particles in the same bunch (Touschek effect) or due to the interaction with the residual gas.

Accurate studies showed that the main background source in DAΦNE is of the second type, which points to the way to reduce it. A fast trigger correlated to a kaon entering into the target cut the main part of the asynchronous background. X rays were detected by DEAR using CCDs (Charge-Coupled Devices), which are excellent X-ray detectors, with very good energy resolution (about 140 eV FWHM at 6 keV), but having the drawback of being non-triggerable devices (since the read-out time per device is at the level of 10 s). A new device, which preserves all good features of CCDs (energy resolution, stability and linearity), but additionally is triggerable - i.e. fast (at the level of $1\mu\text{s}$), was implemented. The new detector was a large area Silicon Drift Detector (SDD), specially designed for SIDDHARTA. The development of the new 1 cm^2 SDD device, together with its readout electronics and very stable power supplies, was partially performed under the Joint Research Activity JRA10 of the I3 project "Study of strongly interacting matter (HadronPhysics)" within FP6 of the EU.

The trigger in SIDDHARTA was given by a system of scintillators which recognized a kaon entering the target making use of the back-to-back production mechanism of the charged kaons at DAΦNE from ϕ decay:

$$\phi \rightarrow K^+ K^- . \quad (1)$$

The SIDDHARTA setup contained 144 SDD chips, 1 cm^2 each, placed around a cylindrical target, filled with high density cryogenic gaseous hydrogen (deuterium or helium). The target was made of kapton, $75\mu\text{m}$ thick, reinforced with aluminium grid.

The SIDDHARTA setup was installed on DAΦNE in late summer 2008, and the period till the end of 2008 was used to debug and optimize the setup performances (degrader optimization included). The kaonic atoms measurements were done in 2009 and data analysis followed in the coming years, which produced the most precise measurement of kaonic hydrogen ¹⁾ and measurements of kaonic helium ^{3 2)} and ^{4 3), 4)}. Kaonic deuterium could not be measured by SIDDHARTA.

2.2 The SIDDHARTA-2 setup

The upgrade of SIDDHARTA to SIDDHARTA-2 is based on the following essential modifications:

- *Trigger geometry and target density*: By placing the upper kaon-trigger detector in front of the target entrance window the probability that a triggered kaon really enters the gas and is stopped there is improved. Making the detector smaller than the entry area gives away some signal, but suppresses efficiently the kaonic lines from "wall-stops" (kaons entering the

gas volume, but passing from the inside of the target to the cylindrical walls). The number “signal per trigger” goes up, which also reduces the accidental background coming along with every trigger. We will also plan to double the gas density which enhances the gas stops and further reduces the wall-stops.

- *Active shielding:* The scintillators surrounding the target will also be used in prompt anti-coincidence if the spatial correlation of SDD and scintillator hits indicates that it originated from a pion (“charged particle veto”). An anticoincidence covering the SDD time window of about 600 ns (with the exception of the 4 ns of the gas stopping time) will reduce the accidental background. Although the scintillators have low efficiency for gammas, the abundance of secondaries from the electromagnetic showers allows a relevant reduction of accidental (“beam”) background. The upper trigger scintillator has 2 functions, it is also used as an anticoincidence counter: after the kaon and eventual prompt kaon-absorption secondaries pass, it vetos beam background.
- *Use of new SDD detectors,* produced by FBK, having a much better active/total surface ratio (about 85%, with respect to 40% in SIDDHARTA SDDs) (see Figure 1).
- *Operating SDDs at a lower temperature:* tests indicate that an improvement of the timing resolution by a factor of 1.5 is feasible by more cooling. The signal enhancement by a factor 2 to 3 is due to moving the target cell closer to the IP, by changing its shape, by a better solid angle of the SDDs and by the higher gas density. In such conditions, with an integrated luminosity of 800 pb^{-1} a precision similar to that obtained for kaonic hydrogen is reachable.

A scheme of the SIDDHARTA-2 internal region of the setup is shown in Figure 2.

In 2019 the first setup, SIDDHARTINO, see Fig. 3, containing 8 SDDs units, aiming to measure kaonic helium to quantify the background in the new DAΦNE configuration, previous to the kaonic deuterium measurement, was installed and tested on DAΦNE (Figure 4). The luminometer detector and SDD calibration system were optimized, while DAΦNE was in commissioning. In early 2020 first collisions were achieved in DAΦNE and SIDDHARTINO is in optimization run.

More details can be found in the presentations to the 2019 LNF International Scientific Committee meetings on the LNF-INFN dedicated web-site.

2.3 Plan for the SIDDHARTA-2 activities in 2020

The LNF group main activities in SIDDHARTA-2 for 2020 will be the following ones:

- run with of the SIDDHARTINO setup on DAΦNE with kaonic helium measurement
- Monte Carlo simulations for the SIDDHARTINO and SIDDHARTA-2 setups and physics;
- data analysis
- preparation and installation of the SIDDHARTA-2 setup on DAΦNE (when background conditions measured by SIDDHARTINO are as in SIDDHARTA as run for kaonic deuterium measurement)
- run with a High Purity Germanium detector for testing the feasibility of other kaonic atoms measurements

In Figure 5 we show the kaonic deuterium simulated spectrum and expected results for an integrated luminosity of 800 pb^{-1} .




Figure 1: The new 2 x 4 SIDDHARTA-2 SDD array together with the readout electronics.




Figure 2: Schematic view of the SIDDHARTA-2 setup




Figure 3: The SIDDHARTINO setup



Figure 4: Photo of SIDDHARTINO setup installed in DAΦNE




Figure 5: Monte Carlo simulated kaonic deuterium spectrum for 800 pb^{-1} .

3 AMADEUS: 2019

The low-energy kaon-nuclei interaction studies represent the main aim of AMADEUS. The negatively charged kaons from DAΦNE can stop inside the target or interact at low energies, giving birth of a series of processes we plan to study. Among these, a key-role is played by the production of $\Lambda(1405)$ which can decay into $\Sigma^0\pi^0$, $\Sigma^+\pi^-$ or $\Sigma^-\pi^+$. We study all these three channels in the same data taking. Another important item is represented by the debated case of the “kaonic nuclear clusters”, especially the K^-pp , and K^-ppn ones. We study these channels by measuring their decays to Λp and to Λd . In the same time, many other kaon-nuclei processes are investigated, either for the first time, or in order to obtain more accurate results than those actually reported in literature. Cross sections, branching ratios, rare hyperon decay processes are investigated, taking advantage of the unique kaon-beam quality delivered by DAΦNE.

In the summer of 2012 a first dedicated target, half cylinder done in pure carbon was realized and installed inside the Drift Chamber of KLOE as a first setup towards the realization of AMADEUS. The target thickness was optimized to have a maximum of stopped kaons (about 24% of the generated ones) without degrading too much the energy of resulting charged particles inside the target material. In the period of data taking a total integrated luminosity of about 90 pb^{-1} was achieved. The ongoing analysis of these data will provide new insights in the low-energy interactions of charged kaons in the nuclear matter.

Activities done in 2019:

- analysis of 2002-2005 KLOE data searching for processes generated by negatively charged kaons interacting at rest or in-flight in the setup materials (wall of the Drift Chamber and gas inside the Drift Chamber); the analyses of the K^- absorption delivering Λp and $\Sigma^0 p$ final states was finalized and results published in ⁵⁾;

- analysis of Λt and $\Sigma^0\pi^0$;
- analysis of the 2012 Carbon target data;
- Monte Carlo dedicated simulations.

3.1 AMADEUS activities in 2020

The main activities of AMADEUS in 2020 will be:

- analyses of data taken with the dedicated carbon target
- Monte Carlo dedicated simulations
- definition of the future strategy for dedicated experiment on DAΦNE and J-PARC.

3.2 Workshops organization

In 2019 the following workshops,

- Symposium on Strange Matter: Kaonic Atoms research in Italy and Japan, LNF-INFN, 15 May 2019.
- Strange Matter Workshop - Strangeness studies in Italy and Japan, LNF-INFN, 15-17 October 2019.
- STRANEX: Recent progress and perspectives in STRange EXotic atoms studies and related topics, ECT*, Trento, 21-25 October 2019.

where the KAONNIS physics was discussed, were organized.

Acknowledgements

The support from LNF Director, Pierluigi Campana and from the DAΦNE, KLOE2 and BTF-LNF teams are gratefully acknowledged.

4 Publications in 2019

1. J. Zmeskal *et al*, Probing Strong Interaction with SIDDHARTA-2, JPS Conf. Proc. **26** (2019) 023012.
2. J-PARC E57 and E62 Collaborations, Kaonic Atom Experiments at J-PARC, JPS Conf. Proc. **26** (2019) 023013.
3. M. Skurzok *et al*, Low-energy K^- Hadronic Interactions with Light Nuclei by AMADEUS, JPS Conf. Proc. **26** (2019) 023011.
4. J-PARC E15 Collaboration, Results of K^- NN Search via the (K^-,n) Reaction at J-PARC, JPS Conf. Proc. **26** (2019) 023008.

5. S. Kawasaki *et al*, $\Lambda(1405)$ Spectroscopy via the In-flight $d(K^-, n)$ Reaction at the J-PARC K1.8BR, JPS Conf. Proc. **26** (2019) 022009.
6. L. Paolis *et al*, Kaonic Atoms Measurement at DAΦNE: SIDDHARTA and SIDDHARTA-2, Springer Proc. Phys. **225** (2019) 191.
7. D. Sirghi *et al*, Kaonic atoms measurements at the DAΦNE Collider, PoS Confinement2018 (2019) 215.
8. H. Asano *et al*, Spectroscopic study of the $\Lambda(1405)$ resonance via the $d(K^-, n)$ reaction at J-PARC, AIP Conf. Proc. **2130** (2019) no.1, 040018.
9. K. Piscicchia *et al*, Low energy antikaon-nucleon/nuclei interaction studies by AMADEUS, AIP Conf.Proc. **2130** (2019) no.1, 020021.
10. C. Curceanu *et al*, The modern era of light kaonic atom experiments, Rev. Mod. Phys. **91** (2019) no.2, 025006.
11. C. Curceanu *et al*, X-ray Detectors for Kaonic Atoms Research at DAΦNE, Condens. Mat. **4** (2019) no.2, 42.
12. M. Miliucci *et al*, Energy Response of Silicon Drift Detectors for Kaonic Atom Precision Measurements, Condens. .Mat. **4** (2019) no.1, 31.
13. A. Scordo, C. Curceanu, M. Miliucci, F. Sirghi, J. Zmeskal, Development of a compact HAPG crystal Von Hamos X-ray spectrometer forextended and diffused sources, arXiv:1903.02826, 9 March 2019.
14. C. Broggini *et al*, Experimental nuclear astrophysics in Italy. Riv. Nuovo Cim. **42** (2019) no.3, 103.
15. R. Del Grande *et al*, Λp correlated production from low energy $K^- {}^{12}C$ interactions by AMADEUS, EPJ Web Conf. **199** (2019) 03010.
16. J. Marton *et al*, Spectroscopy of kaonic atoms at DAΦNE and J-PARC, EPJ Web Conf. **199** (2019) 03004.
17. K. Piscicchia *et al*, Low Energy Antikaon-nucleon/nuclei interaction studies by AMADEUS, EPJ Web Conf. **199** (2019) 01014
18. J. Adamczewski-Musch (HADES Collaboration) *et al*, Strong absorption of hadrons with hidden and open strangeness in nuclear matter, Phys. Rev. Lett. **123** (2019) no.2, 022002.
19. R. Del Grande *et al*, K^- multi-nucleon absorption cross sections and branching ratios in Λp and Σ^0 final states, Eur. Phys. J. C **79** (2019) no.3, 190.
20. S. Ajimura (E15 Collaboration) *et al*, “ $K^- pp$ ” a \bar{K} -Meson Nuclear Bound State, Observed in ${}^3\text{He}(K^-, \Lambda p)n$ Reactions, Phys. Lett. B **789** (2019) 620.
21. B. Adeva *et al*, First measurement of a long-lived $\pi^+\pi^-$ atom lifetime , Phys. Rev. Lett **122** (2019), no 8, 082003.
22. M. Miliucci *et al*, Kaonic Deuterium Precision Measurement at DAΦNE: The SIDDHARTA-2 Experiment, Springer Proc. Phys. **238** (2020) 965.

23. M. Skurzok *et al*, Low-Energy K^- Nucleon/Multi-nucleon Interaction Studies by AMADEUS, Springer Proc. Phys. **238** (2020) 937.
24. C. Curceanu *et al*, Kaonic Deuterium Measurement with SIDDHARTA-2 on DAΦNE, Acta Phys. Polon. B **51** (2020) 251.
25. R. Del Grande *et al*, Recent Experimental Results on the Low-energy K^- Interaction with Nucleons by AMADEUS, Acta Phys. Polon. B **51** (2020) 121.
26. D. Bosnar *et al*, Revisiting the Charged Kaon Mass, Acta Phys. Polon. B **51** (2020) 115.

References

1. M. Bazzi *et al*, Phys. Lett. B **704**, 113 (2011).
2. M. Bazzi *et al*., Phys. Lett. B **697** (3), 199 (2011).
3. M. Bazzi *et al*., Phys. Lett. B **681** , 310 (2009).
4. M. Bazzi *et al*., Phys. Lett. B **714** , 40 (2012).
5. R. Del Grande *et al*., Eur. Phys. J. C **79** 190 (2019).