

Frascati, 7 April '04

Neutrino Masses as a Probe of Grand Unification

G. Altarelli
CERN

Some recent work by our group

G.A., F. Feruglio, I. Masina, hep-ph/0210342
(Addendum: v2 in Nov. '03), hep-ph/0402121.

Reviews:

G.A., F. Feruglio, hep-ph/0206077/0306265

ν Oscillations Imply Different ν Masses

flavour mass

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$


U: mixing matrix

$$\begin{aligned} \nu_e &= \cos\theta \nu_1 + \sin\theta \nu_2 \\ \nu_\mu &= -\sin\theta \nu_1 + \cos\theta \nu_2 \end{aligned} \quad \leftarrow \text{e.g 2 flav.}$$

$\nu_{1,2}$: different mass, different x-dep:

$$\nu_a(x) = e^{i p_a x} \nu_a$$

$$p_a^2 = E^2 - m_a^2$$




$$P(\nu_e \leftrightarrow \nu_\mu) = |\langle \nu_\mu(L) | \nu_e \rangle|^2 = \sin^2(2\theta) \cdot \sin^2(\Delta m^2 L / 4E)$$

At a distance L , ν_μ from μ^- decay can produce e^- via charged weak interact's

Solid evidence for ν oscillations
(+LSND unclear)

$\Delta m^2_{\text{atm}} \sim 2.5 \cdot 10^{-3} \text{ eV}^2,$
 $\Delta m^2_{\text{sol}} \sim 7 \cdot 10^{-5} \text{ eV}^2$
 $(\Delta m^2_{\text{LSND}} \sim 1 \text{ eV}^2)$




Solar ν 's



For the LA solution the oscill's occur inside the sun thru the MSW effect

Mikhaev and Smirnov; Wolfenstein

Atmospheric ν 's



$$\nu_{\mu} \rightarrow \nu_{\tau}$$

atmospheric ν 's traverse different L depending on azimuth θ (up-down asymm.)

Evolution in vacuum and in matter

$$\nu_e = \cos\theta \nu_1 + \sin\theta \nu_2$$

$$\Delta m^2 = m_2^2 - m_1^2 > 0 \quad \nu_\mu = -\sin\theta \nu_1 + \cos\theta \nu_2$$

$$i \frac{d}{dt} \begin{bmatrix} \nu_e \\ \nu_\mu \end{bmatrix} = H_{eff} \begin{bmatrix} \nu_e \\ \nu_\mu \end{bmatrix} \quad H_{eff} = \frac{\Delta m^2}{4E} \begin{bmatrix} -\cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$$

In vacuum, 2 flavours, apart from multiples of the identity

In matter CC int's on electrons introduce a flavour dep.
(coherent forward scattering on electrons)

$$H_{eff} = \frac{\Delta m^2}{4E} \begin{bmatrix} -\cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix} + \begin{bmatrix} \sqrt{2} G_F N_e & 0 \\ 0 & 0 \end{bmatrix} \quad N_e: \text{ n. of e per unit V}$$

The mixing angle is changed
A resonance can appear (MSW)

$$\tan 2\theta_m = \frac{\tan 2\theta}{1 - \frac{2\sqrt{2}EG_F N_e}{\Delta m^2 \cos 2\theta}}$$


G. Altarelli

Mikhaev and Smirnov; Wolfenstein

Atmospheric Neutrinos

Smirnov,
Aachen'03

SuperKamiokande:



Best fit point:

$$\sin^2 2\theta_{23} = 1.0$$

$$\Delta m_{32}^2 = 2.0 \cdot 10^{-3} \text{ eV}^2$$

$$\Delta m_{32}^2 = (1.3 - 3.0) \cdot 10^{-3} \text{ eV}^2$$
$$\sin^2 2\theta_{23} > 0.9 \quad (90\% \text{ C.L.})$$

Confirmed by
MACRO,
SOUDAN
K2K

Combined analysis of CHOOZ,
atmospheric (SK) and solar data:

$$\sin^2 2\theta_{13} < 0.067 \quad (3\sigma)$$

G.L. Fogli et al, hep-ph/p0308055

ν Reactions in SNO

CC




- Good measurement of ν_e energy spectrum
- Weak directional sensitivity $\propto 1 - 1/3 \cos(\theta)$
- ν_e only.

NC




- Equal cross section for all ν types
- Measure total ^8B ν flux from the sun.

ES



- Low Statistics
- Mainly sensitive to ν_e , some sensitivity to ν_μ and ν_τ
- Strong directional sensitivity

April '02



$$\frac{\Phi_{cc}}{\Phi_{es}} = \frac{\nu_e}{\nu_e + 0.154(\nu_\mu + \nu_\tau)} = 1?$$

$$\frac{\Phi_{cc}}{\Phi_{nc}} = \frac{\nu_e}{\nu_e + \nu_\mu + \nu_\tau} = 1?$$

G. Alta

Results of April '02

Signal Extraction in Φ_{CC} , Φ_{NC} , Φ_{ES} · $E_{\text{Threshold}} > 5 \text{ MeV}$

$$\Phi_{CC}(\nu_e) = 1.76^{+0.06}_{-0.05} \text{ (stat.) } ^{+0.09}_{-0.09} \text{ (syst.) } \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$

$$\Phi_{ES}(\nu_\chi) = 2.39^{+0.24}_{-0.23} \text{ (stat.) } ^{+0.12}_{-0.12} \text{ (syst.) } \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$

$$\Phi_{NC}(\nu_\chi) = 5.09^{+0.44}_{-0.43} \text{ (stat.) } ^{+0.46}_{-0.43} \text{ (syst.) } \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$

Signal Extraction in Φ_e , $\Phi_{\mu\tau}$

$$\Phi_e = 1.76^{+0.05}_{-0.05} \text{ (stat.) } ^{+0.09}_{-0.09} \text{ (syst.) } \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$

$$\Phi_{\mu\tau} = 3.41^{+0.45}_{-0.45} \text{ (stat.) } ^{+0.48}_{-0.45} \text{ (syst.) } \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$$

Note: $\Phi_{\mu,\tau} \sim 2 \Phi_e$


(We receive an equal amount of ν_e , ν_μ , ν_τ)

G. Altarelli

The measured total ν flux is in perfect agreement with the Solar Standard Model!!

But: $\Phi_e \sim 1/3 (\Phi_e + \Phi_\mu + \Phi_\tau)$

$$\Phi_{\text{ssm}} = 5.05^{+1.01}_{-0.81} \quad \Phi_{\text{sno}} = 5.09^{+0.44+0.46}_{-0.43-0.43}$$




G. Altarelli

Direct evidence for $\nu_e \rightarrow \nu_{\mu,\tau}$ oscill's as solution of the solar ν_e deficit!

Recent important results from KamLAND

Dec'02

Kamioka
Liquid
scintillator
AntiNeutrino
Detector






1 kton

Reactor $\bar{\nu}_e$ ($E > 2.6$ MeV) detected 180 Km
away at Kamiokande site

G. Altarelli

First results from KamLAND


- Solar oscill.'s confirmed on earth
 - Large angle sol. established
- Best fit: $\Delta m^2 \sim 7.10^{-5} \text{ eV}^2$, $\sin^2 2\theta = 1$
- $\bar{\nu}_e$ from reactors behave as ν_e from sun:
Constraint on ~~CPT~~ models



In summary for solar ν 's:

Before Kamland

After Kamland



G. Altarelli

Note the change of scale





Sept.'03: SNO new results

Salt added to D₂O:
Better NC sensitivity

- Previous results confirmed
- More precision
- The upper Δm^2 part of the LA sol. now disfavoured
- θ_{12} is now 5.4σ from maximal

All data now



ν Oscillations: Summary of Exp. Facts

Homestake, Gallex, Sage, (Super)Kamiokande, Macro...

GNO, K2K, ...

Atmospheric:

$$\Delta m_{\text{atm}}^2 \sim 2.5 \cdot 10^{-3} \text{ eV}^2$$

$$\sin^2 \theta_{23} \sim 1/2$$

$\nu_{\mu} \rightarrow \nu_{\tau}$ dominant

$\nu_{\mu} \rightarrow \nu_e$ small 

(Chooz $|U_{13}| < \sim 0.2$)

$\nu_{\mu} \rightarrow \nu_{\text{sterile}}$ small

Solar:

The MSW-LA solution selected

$$\Delta m^2 \sim 7 \cdot 10^{-5} \text{ eV}^2, \sin^2 \theta_{12} \sim 0.3$$

$\nu_e \rightarrow \nu_{\mu}, \nu_{\tau}$ dominant

$\nu_e \rightarrow \nu_{\text{sterile}}$ small

after KAMLAND,
SNO-salt

LSND:

true or false?

MINIBOONE (in progress)


$$\Delta m^2 \sim 1 \text{ eV}^2, \sin^2 \theta \sim \text{small}$$

$\nu_{\mu} \rightarrow \nu_e, \nu_{\text{sterile}}$

CPT violation?

G. Altarelli

parameter	best fit	2σ	3σ	5σ
Δm_{21}^2 [10^{-5}eV^2]	6.9	6.0–8.4	5.4–9.5	2.1–28
Δm_{31}^2 [10^{-3}eV^2]	2.6	1.8–3.3	1.4–3.7	0.77–4.8
$\sin^2 \theta_{12}$	0.30	0.25–0.36	0.23–0.39	0.17–0.48
$\sin^2 \theta_{23}$	0.52	0.36–0.67	0.31–0.72	0.22–0.81
$\sin^2 \theta_{13}$	0.006	≤ 0.035	≤ 0.054	≤ 0.11



ν oscillations measure Δm^2 . What is \bar{m} ?

$$\Delta m^2_{\text{atm}} \sim 2.5 \cdot 10^{-3} \text{ eV}^2; \quad \Delta m^2_{\text{sun}} \sim 7 \cdot 10^{-5} \text{ eV}^2$$

- Direct limits (PDG '02)

$$m_{\nu_e} < 2.8 \text{ eV}$$

$$m_{\nu_\mu} < 170 \text{ KeV}$$

$$m_{\nu_\tau} < 18.2 \text{ MeV}$$


End-point tritium β decay (Mainz)

- $0\nu\beta\beta$


- Cosmology $\Omega_\nu h^2 \sim \sum_i m_i / 94 \text{ eV}$ ($h^2 \sim 1/2$)

$$\sum_i m_i \sim 0.69 \text{ eV (95\%)} \quad [\Omega_\nu \sim 0.014]$$

WMAP



Any ν mass 0.23-1 eV



Why ν 's so much lighter than quarks and leptons?


New powerful cosmological limit


All info on the absolute scale of ν mass is very important!

Finding $0\nu\beta\beta$ would also prove Majorana ν 's

G. Altarelli

Assumes some priors!
Could be somewhat relaxed





Neutrino masses
are really special!

$m_t / (\Delta m^2_{\text{atm}})^{1/2} \sim 10^{12}$

Massless ν 's?

- no ν_R
- L conserved

Small ν masses?

- ν_R very heavy
- L not conserved

How to guarantee a massless neutrino?

1) ν_R does not exist



No Dirac mass

$$\bar{\nu}_L \nu_R + \nu_R \bar{\nu}_L$$

and

2) Lepton Number is conserved



No Majorana mass

$$\bar{\nu}^c \nu \rightarrow \nu_R^T C \nu_R \text{ or } \nu_L^T C \nu_L$$

G. Altarelli


$$C = i\gamma^0 \gamma^2$$

Neutrinos:

Dirac mass: $\bar{\nu}_L \nu_R + \bar{\nu}_R \nu_L$
(needs ν_R)

ν 's have no electric charge. Their only charge is lepton number L

IF L is not conserved (not a good quantum number)
 ν and $\bar{\nu}$ are not really different



Majorana mass: $\nu_R^T \nu_R$ or $\nu_L^T \nu_L$
(we omit the charge conj. matrix C)

Violates L, B-L by $|\Delta L| = 2$

Weak isospin I

$$\nu_L \Rightarrow I = 1/2, I_3 = 1/2$$

$$\nu_R \Rightarrow I = 0, I_3 = 0$$

Dirac Mass:

$$\bar{\nu}_L \nu_R + \bar{\nu}_R \nu_L \quad |\Delta I| = 1/2$$

Can be obtained from Higgs doublets: $\nu_L \bar{\nu}_R H$

Majorana Mass ($\Delta L=2$):

- $\nu_L^T \nu_L \quad |\Delta I| = 1$

Non ren., dim. 5 operator: $\nu_L^T \nu_L H H$

- $\nu_R^T \nu_R \quad |\Delta I| = 0$

G. Altarelli

Directly
compatible
with $SU(2) \times U(1)$!

See-Saw Mechanism

Yanagida; Glashow;
Gell-Mann, Ramond, Slansky;
Mohapatra, Senjanovic

$M \nu_R^T \nu_R$ allowed by $SU(2) \times U(1)$
 Large Majorana mass M (as large as the cut-off)

$$m_D \bar{\nu}_L \nu_R$$

Dirac mass m from
Higgs doublet(s)

$$\begin{matrix} & \nu_L & \nu_R \\ \nu_L & \left[\begin{array}{cc} 0 & m_D \\ m_D & M \end{array} \right] \end{matrix}$$

$$M \gg m_D$$

Eigenvalues

$$\nu_{\text{light}} = \frac{-m_D^2}{M}, \quad \nu_{\text{heavy}} = M$$

sign conventional
for fermions

In general ν mass terms are:

$$\mathcal{L}_\nu = \bar{L}h\nu_R H + \text{h.c.} + \nu_R^T M_R \nu_R + \nu_L^T \frac{\lambda}{M_L} \nu_L H H$$

Dirac $m_D = h v$
 $v = \langle 0 | H | 0 \rangle$

Majorana $m = \frac{\lambda v^2}{M_L}$

More general see-saw mechanism:

$$\begin{pmatrix} \nu_L \\ \nu_R \end{pmatrix} \begin{bmatrix} \lambda v^2 / M_L & m_D \\ m_D & M_R \end{bmatrix}$$


$m_{\text{light}} \sim \frac{m_D^2}{M_R}$ and/or $\frac{\lambda v^2}{M_L}$
 $m_{\text{heavy}} \sim M_R$

$$m_{\text{eff}} = \nu_L^T m_{\text{light}} \nu_L$$

Neutrinos are (probably) Majorana particles:

$$\nu_L^T m_\nu \nu_L$$

See-saw




$$m_\nu = m_D^T M^{-1} m_D$$

connection with m_D

More in general: non ren. O_5 operator

$$\lambda/M \nu_L^T H H^T \nu_L$$

e.g from



N: new particle $I_w=0,1$

A very natural and appealing explanation:

ν 's are nearly massless because they are Majorana particles and get masses through L non conserving interactions suppressed by a large scale $M \sim M_{\text{GUT}}$


$$m_\nu \sim \frac{m^2}{M}$$

$m \sim m_t \sim v \sim 200 \text{ GeV}$
 M : scale of L non cons.

Note:

$$m_\nu \sim (\Delta m_{\text{atm}}^2)^{1/2} \sim 0.05 \text{ eV}$$

$$m \sim v \sim 200 \text{ GeV}$$



$$M \sim 10^{15} \text{ GeV}$$

Neutrino masses are a probe of physics at M_{GUT} !


B and L conservation in SM:

"Accidental" symmetries: in SM there is no dim.4 gauge invariant operator that violates B and/or L (if no ν_R , otherwise $M \nu_R^T \nu_R$ is dim-3 $|\Delta L|=2$)
 The same is true in SUSY with R-parity cons.

e. g. for the $\Delta B = \Delta L = -1$ transition $u + u \rightarrow e^+ + \bar{d}$

all good quantum numbers are conserved:
 e.g. colour $u \sim 3$, $\bar{d} \sim \bar{3}$ and $3 \times 3 = 6 + \bar{3}$ but

$$\frac{\lambda}{M^2} \bar{d}^c \Gamma u \bar{e}^c \Gamma u \quad \longrightarrow \quad \text{dim-6}$$



Once ν_R is introduced (Dirac mass) large Majorana mass is naturally induced \longrightarrow see-saw

Dark Matter

WMAP

Most of the Universe is not made up of atoms: $\Omega_{\text{tot}} \sim 1$, $\Omega_b \sim 0.044$, $\Omega_m \sim 0.27$
Most is Dark Matter and Dark Energy

Most Dark Matter is Cold (non relativistic at freeze out)
Significant Hot Dark matter is disfavoured
Neutrinos are not much cosmo-relevant: $\Omega_\nu < 0.015$ (WMAP)

SUSY has excellent DM candidates: Neutralinos
Also Axions are still viable

For 3 neutrinos: $\Omega_\nu < 0.015 \rightarrow m_\nu < 0.23 \text{ eV} \sim 5(\Delta m^2_{\text{atm}})^{1/2}$


the exact value depends on the cosmological model: can be somewhat relaxed

G. Altarelli

$m_\nu < \sim 1 \text{ eV}$

Baryogenesis

A most attractive possibility:

BG via Leptogenesis near the GUT scale

$T \sim 10^{12 \pm 3}$ GeV (after inflation)

Buchmuller, Yanagida,
Plumacher, Ellis, Lola,
Giudice et al, Fujii et al

Only survives if $\Delta(B-L)$ is not 0
(otherwise is washed out at T_{ew} by instantons)

Main candidate: decay of lightest ν_R ($M \sim 10^{12}$ GeV)

L non conserv. in ν_R out-of-equilibrium decay:

B-L excess survives at T_{ew} and gives the obs. B asymm.

Quantitative studies confirm that the range of m_i from ν oscill's is compatible with BG via (thermal) LG

In particular the bound
was derived

$$m_i < 10^{-1} \text{ eV}$$

Close to WMAP

G. Altarelli

Buchmuller, Di Bari, Plumacher
Giudice et al

The scale of the cosmological constant is a big mystery.

$\Omega_\Lambda \sim 0.65 \quad \longrightarrow \quad \rho_\Lambda \sim (2 \cdot 10^{-3} \text{ eV})^4 \sim (0.1 \text{ mm})^{-4}$

In Quantum Field Theory: $\rho_\Lambda \sim (\Lambda_{\text{cutoff}})^4$ Similar to m_ν !?

If $\Lambda_{\text{cutoff}} \sim M_{\text{Pl}}$ \longrightarrow $\rho_\Lambda \sim 10^{123} \rho_{\text{obs}}$

Exact SUSY would solve the problem: $\rho_\Lambda = 0$

But SUSY is broken: $\rho_\Lambda \sim (\Lambda_{\text{SUSY}})^4 \sim 10^{59} \rho_{\text{obs}}$

It is interesting that the correct order is $(\rho_\Lambda)^{1/4} \sim (\Lambda_{\text{EW}})^2 / M_{\text{Pl}}$


So far no solution:

- A modification of gravity at 0.1 mm? (large extra dim.)
- Leak of vac. energy to other universes (wormholes)?

...

G. Altarelli

Other problem:
Why now?



Quintessence?

The current experimental situation is still unclear

- LSND: true or false?
- what is the absolute scale of ν masses?
-


Different classes of models are possible:

If LSND true

sterile ν (s)??
CPT violat'n??

• "3-1"

ν_{sterile}



$m^2 \sim 1-2 \text{eV}^2$

If LSND false



3 light ν 's are OK

We assume this case here

• Degenerate ($m^2 \gg \Delta m^2$)



$m^2 < o(1) \text{eV}^2$

• Inverse hierarchy

sol



$m^2 \sim 10^{-3} \text{eV}^2$

• Normal hierarchy




$m^2 \sim 10^{-3} \text{eV}^2$

3-ν Models

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

flavour

mass



$$U = U_{\text{P-MNS}}$$

Pontecorvo

Maki, Nakagawa, Sakata

In basis where e^- , μ^- , τ^- are diagonal:

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \sim$$

$s = \text{solar: large}$

$$\sim \begin{pmatrix} c_{13} & c_{12} & c_{13}s_{12} \\ \dots & \dots & \dots \\ \dots & \dots & \dots \end{pmatrix}$$

CHOOZ: $|s_{13}| < \sim 0.2$

atm.: $\sim \text{max}$



$$U = \begin{pmatrix} c & -s & 0 \\ \frac{s}{\sqrt{2}} & \frac{c}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{s}{\sqrt{2}} & \frac{c}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$

(some signs are conventional)

G. Altarelli

$m_\nu \sim U \begin{bmatrix} e^{i\phi_1} m_1 & 0 & 0 \\ 0 & e^{i\phi_2} m_2 & 0 \\ 0 & 0 & m_3 \end{bmatrix} U^T$

In general 9 parameters:
 3 masses, 3 angles,
 3 phases

$L^T m_\nu L$

For $s_{13} \sim 0$:

$m_\nu \sim \begin{bmatrix} m_1 c^2 + m_2 s^2 & (m_1 - m_2) cs / \sqrt{2} & (m_1 - m_2) cs / \sqrt{2} \\ \dots & (m_1 s^2 + m_2 c^2 + m_3) / 2 & (m_1 s^2 + m_2 c^2 - m_3) / 2 \\ \dots & \dots & (m_1 s^2 + m_2 c^2 + m_3) / 2 \end{bmatrix}$

$0\nu\beta\beta \longrightarrow$

Note:

- m_ν is symmetric
- phases included in m_i

Relation between masses and frequencies:

$$P(\nu_e \leftrightarrow \nu_\mu) = P(\nu_e \leftrightarrow \nu_\tau) = 1/2 \sin^2 2\theta_{12} \cdot \sin^2 \Delta_{\text{sun}}$$

$$P(\nu_\mu \leftrightarrow \nu_\tau) = \sin^2 \Delta_{\text{atm}} - 1/4 \sin^2 2\theta_{12} \cdot \sin^2 \Delta_{\text{sun}}$$

$$\Delta_{\text{sun}} = \frac{m_2^2 - m_1^2}{4E} L \quad ; \quad \Delta_{\text{atm}} = \frac{m_3^2 - m_{1,2}^2}{4E} L$$

G. Altarelli

In our def.: $\Delta_{\text{sun}} > 0$, $\Delta_{\text{atm}} >$ or < 0

$0\nu\beta\beta$ can tell degenerate, inverted or normal hierarchy

$$|m_{ee}| = c_{13}^2 [m_1 c_{12}^2 + e^{i\alpha} m_2 s_{12}^2] + m_3 e^{i\beta} s_{13}^2$$

LA: $\sim 0.3-1$ 

Degenerate: $\sim |m| |c_{12}^2 + e^{i\alpha} s_{12}^2|$


$$|m_{ee}| \sim |m| (0.3 - 1) < 0.23-1 \text{ eV}$$

IH: $\sim (\Delta m_{\text{atm}}^2)^{1/2} |c_{12}^2 + e^{i\alpha} s_{12}^2|$

$$|m_{ee}| \sim (1.6-5) 10^{-2} \text{ eV}$$

NH: $\sim (\Delta m_{\text{sol}}^2)^{1/2} s_{12}^2 + (\Delta m_{\text{atm}}^2)^{1/2} e^{i\beta} s_{13}^2$

$$|m_{ee}| \sim (\text{few}) 10^{-3} \text{ eV}$$



Present exp. limit: $m_{ee} < 0.3-0.5 \text{ eV}$
(and a hint of signal?????)

Evidence for $0\nu\beta\beta$?

Heidelberg-Moscow
Klapdor-Kleingrothaus et al

Not at all compelling!!!!

New recent ('04) paper


Iff true: (WMAP ??)

$$m_{ee}/z = 0.39 \pm 0.11 \text{ eV} \gg (\Delta m^2_{\text{atm}})^{1/2}$$


($z \sim 0.6-2.8$
uncert. matrix element)

would clearly point
to degenerate models

G. Altarelli



Feruglio, Strumia, Vissani



Degenerate ν 's

$$m^2 \gg \Delta m^2$$

- Apriori compatible with hot dark matter ($m \sim 1-2$ eV)
 - was considered by many
- Limits on m_{ee} from $0\nu\beta\beta$ then imply large mixing also for solar oscillations: (Vissani; Georgi, Glashow)

→ $m_{ee} < 0.3-0.5$ eV (Exp)

$$m_{ee} = c_{13}^2 (m_1 c_{12}^2 + m_2 s_{12}^2) + s_{13}^2 m_3 \sim m_1 c_{12}^2 + m_2 s_{12}^2$$

If $|m_1| \sim |m_2| \sim |m_3| \sim 1-2$ eV → $m_1 = -m_2$ and $c_{12}^2 \sim s_{12}^2$

LA solution: $\sin^2\theta \sim 0.3$ → $\cos^2\theta - \sin^2\theta \sim 0.4$ ↷

a moderate suppression factor!

Trusting WMAP: $|m| < 0.23$ eV, only a moderate degeneracy is allowed: for LA, $m/(\Delta m_{atm}^2)^{1/2} < 5$, $m/(\Delta m_{sol}^2)^{1/2} < 30$.

Less constraints from $0\nu\beta\beta$ (both $m_1 = \pm m_2$ allowed)

G. Altarelli

Recall: leptogenesis prefers $|m| < 0.1$ eV

After KamLAND, SNO and WMAP not too much hierarchy is needed for ν masses:



$$r \sim \Delta m_{\text{sol}}^2 / \Delta m_{\text{atm}}^2 \sim 1/40$$

Precisely at 3σ : $0.018 < r < 0.053$

or

$$m_{\text{heaviest}} < 1 - 0.23 \text{ eV}$$

$$m_{\text{next}} > \sim 7 \cdot 10^{-3} \text{ eV}$$




Anarchical or semi-anarchical models

Anarchy (or accidental hierarchy):
No structure in the leptonic sector

Hall, Murayama, Weiner

See-Saw:
 $m_\nu \sim m^2/M$
produces hierarchy
from random m, M


could fit LA



But: all mixing angles
should be large

marginal for LA \rightarrow
predicts θ_{13} near
bound

G. Altarelli



Semianarchy: no structure in 23

Consider a matrix like $m_\nu \sim \begin{pmatrix} \lambda^2 & \lambda & \lambda \\ \lambda & 1 & 1 \\ \lambda & 1 & 1 \end{pmatrix}$

Note: $\theta_{13} \sim \lambda$
 $\theta_{23} \sim 1$

with coeff.s of $o(1)$ and $\det 23 \sim o(1)$
[$\lambda \sim 1$ corresponds to anarchy]


After 23 and 13 rotations $m_\nu \sim \begin{pmatrix} \lambda^2 & \lambda & 0 \\ \lambda & \eta & 0 \\ 0 & 0 & 1 \end{pmatrix}$

Normally two masses are of $o(1)$ and $\theta_{12} \sim \lambda$
But if, accidentally, $\eta \sim \lambda$, then the solar angle is also large.

The advantage over anarchy is that θ_{13} is small, but
the hierarchy $m^2_3 \gg m^2_2$ is accidental

Inverted Hierarchy

Zee, Joshipura et al;
 Mohapatra et al; Jarlskog et al;
 Frampton, Glashow; Barbieri et al
 Xing; Giunti, Tanimoto



An interesting
 model for double
 maximal mixing (bimixing):

$$U \sim \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 1/2 & 1/2 & -1/\sqrt{2} \\ 1/2 & 1/2 & 1/\sqrt{2} \end{bmatrix}$$

1st approximation

$$m_{\nu \text{diag}} = \begin{bmatrix} m & 0 & 0 \\ 0 & -m & 0 \\ 0 & 0 & 0 \end{bmatrix} ; \quad U m_{\nu \text{diag}} U^T = 1/\sqrt{2} \begin{bmatrix} 0 & m & m \\ m & 0 & 0 \\ m & 0 & 0 \end{bmatrix}$$

Can arise from see-saw or dim-5 $L^T H H^T L$
 e.g. by approximate $L_e - L_\mu - L_\tau$ symmetry

- 1-2 degeneracy stable under rad. corr.'s

G. Altarelli

1st approximation

$$m_{\nu\text{diag}} = \begin{bmatrix} m & 0 & 0 \\ 0 & -m & 0 \\ 0 & 0 & 0 \end{bmatrix} ; \quad Um_{\nu\text{diag}}U^T = 1/\sqrt{2} \begin{bmatrix} 0 & m & m \\ m & 0 & 0 \\ m & 0 & 0 \end{bmatrix}$$

- LA? This texture prefers θ_{sol} closer to maximal than θ_{atm}
i.e $\theta_{\text{sol}} - \pi/4$ small for $(\Delta m^2_{\text{sol}}/\Delta m^2_{\text{atm}})_{\text{LA}} \sim 1/40$

In fact: 12 \rightarrow $\begin{bmatrix} 0 & m \\ m & 0 \end{bmatrix} \rightarrow$ Pseudodirac θ_{12} maximal 23 \rightarrow $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \rightarrow \theta_{23} \sim o(1)$

With perturbations: $\begin{bmatrix} 0 & m & m \\ m & 0 & 0 \\ m & 0 & 0 \end{bmatrix} \rightarrow m \begin{bmatrix} \delta & 1 & 1 \\ 1 & \eta & \eta \\ 1 & \eta & \eta \end{bmatrix}$

$$\text{tg}^2 \theta_{12} \sim 1 + o(\delta + \eta) \quad (\Delta m^2_{\text{sol}}/\Delta m^2_{\text{atm}})_{\text{LA}} \sim o(\delta + \eta)$$

- In principle one can use the charged lepton mixing to go away from θ_{12} maximal.
In practice constraints from θ_{13} small ($\delta\theta_{12} \sim \theta_{13}$)

For the corrections to bimixing from
the charged lepton sector,
typically $|\sin\theta_{13}| \sim (1 - \tan^2\theta_{12})/4$

GA, Feruglio, Masina '04





Figure 1: Taking an upper bound on $|U_{e3}|$ respectively equal to 0.23, 0.1, 0.05, 0.01, we show (from yellow to red) the allowed regions of the plane $[s_{12}^e, s_{13}^e]$. Each plot is obtained by setting α_1 to a particular value, while leaving $\alpha_2 + \delta_e$ free. We keep the present 3σ window for δ_{sol} [10].

- In general more θ_{12} is close to maximal, more is IH likely
G. Altarelli

Normal Hierarchy



- Assume 3 widely split light neutrinos.
- For u, d and l Dirac matrices the 3rd generation eigenvalue is dominant.
- May be this is also true for $m_{\nu D}$: $\text{diag } m_{\nu D} \sim (0, 0, m_{D3})$.
- Assume see-saw is dominant: $m_\nu \sim m_D^T M^{-1} m_D$
See-saw quadratic in m_D : tends to enhance hierarchy
- Maximally constraining: GUT's relate q, l, ν masses!

- A crucial point: in the 2-3 sector we need both large m_3 - m_2 splitting and large mixing.

$$m_3 \sim (\Delta m_{\text{atm}}^2)^{1/2} \sim 5 \cdot 10^{-2} \text{ eV}$$

$$m_2 \sim (\Delta m_{\text{sol}}^2)^{1/2} \sim 8 \cdot 10^{-3} \text{ eV for LA}$$

- The "theorem" that large Δm_{32} implies small mixing (pert. th.: $\theta_{ij} \sim 1/|E_i - E_j|$) is not true in general: all we need is $(\text{sub})\det[23] \sim 0$

- Example: $m_{23} \sim \begin{bmatrix} x^2 & x \\ x & 1 \end{bmatrix}$

Det = 0; Eigenvl's: 0, $1+x^2$
 Mixing: $\sin^2 2\theta = 4x^2/(1+x^2)^2$



So all we need are natural mechanisms for $\det[23]=0$

For $x \sim 1$
 large splitting
 and large mixing!

Examples of mechanisms for $\text{Det}[23] \sim 0$

see-saw $m_\nu \sim m_D^T M^{-1} m_D$

1) A ν_R is lightest and coupled to μ and τ

King; Allanach; Barbieri et al.....

$$M \sim \begin{bmatrix} \varepsilon & 0 \\ 0 & 1 \end{bmatrix} \longrightarrow M^{-1} \sim \begin{bmatrix} 1/\varepsilon & 0 \\ 0 & 1 \end{bmatrix} \approx \begin{bmatrix} 1/\varepsilon & 0 \\ 0 & 0 \end{bmatrix}$$

$$m_\nu \sim \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1/\varepsilon & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & c \\ b & d \end{bmatrix} \approx 1/\varepsilon \begin{bmatrix} a^2 & ac \\ ac & c^2 \end{bmatrix}$$

2) M generic but m_D "lopsided"

$$m_D \sim \begin{bmatrix} 0 & 0 \\ x & 1 \end{bmatrix}$$

Albright, Barr; GA, Feruglio,

$$m_\nu \sim \begin{bmatrix} 0 & x \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ b & c \end{bmatrix} \begin{bmatrix} 0 & 0 \\ x & 1 \end{bmatrix} = c \begin{bmatrix} x^2 & x \\ x & 1 \end{bmatrix}$$

Caution: if $0 \rightarrow 0(\varepsilon)$, $\text{det}23=0$ could be spoiled by suitable $1/\varepsilon$ terms in M^{-1}

An important property of SU(5)

Left-handed quarks have small mixings (V_{CKM}),
but right-handed quarks can have large mixings (unknown).

In SU(5):
LH for d quarks \longleftrightarrow RH for l- leptons

$$\bar{5} \rightarrow \bar{d}_R \leftarrow 10$$

$$m_d \sim \bar{d}_R d_L$$

$$10 \rightarrow \bar{e}_R \leftarrow \bar{5}$$

$$m_e \sim \bar{e}_R e_L$$

$$\bar{5} : (\underbrace{\bar{d}, \bar{d}, \bar{d}}_R, \underbrace{\nu, e^-}_L)$$

$$m_d = m_e^T$$

cannot be exact, but approx.

Most "lopsided" models are based on this fact. In these models large atmospheric mixing arises (at least in part) from the charged lepton sector.

G. Altarelli

- Hierarchical ν 's and see-saw dominance

$$L^T m_\nu L \rightarrow m_\nu \sim m_D^2/M$$

allow to relate q , l , ν masses and mixings in GUT models.
For dominance of dim-5 operators \rightarrow less constraints

$$\lambda^2/M L^T L H H \rightarrow m_\nu \sim \lambda^2 v^2/M$$


- The correct pattern of masses and mixings, also including ν 's, is obtained in simple models based on

$$SU(5) \times U(1)_{\text{flavour}}$$

Ramond et al; GA, Feruglio+Masina; Buchmuller et al;
King et al; Yanagida et al, Berezhiani et al; Lola et al.....

- $SO(10)$ models could be more predictive, as are non abelian flavour symmetries, eg $O(3)$

Albright, Barr; Babu et al; Buccella et al; Barbieri et al;
Raby et al; King, Ross

- The non trivial pattern of fermion masses and mixing demands a flavour structure (symmetry)
- (SUSY) $SU(5)XU(1)_F$ models offer a minimal description of flavour symmetry 
- A flexible enough framework used to realize and compare models with anarchy or hierarchy (direct or inverse) in ν sector, with see-saw dominance or not.

- On this basis we found that for LA there is still a significant preference for hierarchy vs anarchy

G.A., F. Feruglio, I. Masina, hep-ph/0210342 (v2 Nov '03)

Previous related work: Haba,Murayama; Hirsch,King; Vissani; Rosenfeld,Rosner; Antonelli et al....

Hierarchy for masses and mixings via horizontal U(1) charges.

Froggatt, Nielsen '79

Principle:

A generic mass term

$$\bar{R}_1 m_{12} L_2 H$$

is forbidden by U(1)
if $q_1 + q_2 + q_H$ not 0

q_1, q_2, q_H :
U(1) charges of
 \bar{R}_1, L_2, H

U(1) broken by vev of "flavon" field θ with U(1) charge $q_\theta = -1$.
The coupling is allowed: if $\text{vev } \theta = w$, and $w/M = \lambda$ we get:

$$\bar{R}_1 m_{12} L_2 H (\theta/M)^{\Delta_{\text{charge}}} \quad m_{12} \rightarrow m_{12} \lambda^{q_1 + q_2 + q_H}$$

Hierarchy: More $\Delta_{\text{charge}} \rightarrow$ more suppression (λ small)

One can have more flavons (λ, λ', \dots)
with different charges (>0 or <0) etc \rightarrow many versions

G. Altarelli

With suitable charge assignments all relevant patterns can be obtained

Recall: $u \sim 10 \ 10$
 $d=e^T \sim \bar{5} \ 10$
 $\nu_D \sim \bar{5} \ 1; M_{RR} \sim 1 \ 1$

No structure for leptons

No automatic $\det 23 = 0$

Automatic $\det 23 = 0$

G. Altarelli

1st fam. 2nd 3rd

$\Psi_{10}: (5, 3, 0)$
 $\Psi_5: (2, 0, 0)$
 $\Psi_1: (1, -1, 0)$

Equal 2,3 ch. for lopsided

Model	Ψ_{10}	Ψ_5	Ψ_1	(H_u, H_d)
Anarchical (<i>A</i>)	(3,2,0)	(0,0,0)	(0,0,0)	(0,0)
Semi-Anarchical (<i>SA</i>)	(2,1,0)	(1,0,0)	(2,1,0)	(0,0)
Hierarchical (<i>H_I</i>)	(6,4,0)	(2,0,0)	(1,-1,0)	(0,0)
Hierarchical (<i>H_{II}</i>)	(5,3,0)	(2,0,0)	(1,-1,0)	(0,0)
Inversely Hierarchical (<i>IH_I</i>)	(3,2,0)	(1,-1,-1)	(-1,+1,0)	(0,+1)
Inversely Hierarchical (<i>IH_{II}</i>)	(6,4,0)	(1,-1,-1)	(-1,+1,0)	(0,+1)

all charges positive

not all charges positive

All entries are a given power of λ times a free $o(1)$ coefficient

$$m_u \sim v_u \begin{bmatrix} \lambda^{10} & \lambda^8 & \lambda^5 \\ \lambda^8 & \lambda^6 & \lambda^3 \\ \lambda^5 & \lambda^3 & 1 \end{bmatrix}$$

In a statistical approach we generate these coeff.s as random complex numbers $\rho e^{i\phi}$ with $\phi = [0, 2\pi]$ and $\rho = [0.5, 2]$ (default) or $[0.8, 1.2]$, or $[0.95, 1.05]$ or $[0, 1]$ (real numbers also considered for comparison)

For each model we evaluate the success rate (over many trials) for falling in the exp. allowed window:

(boundaries $\sim 3\sigma$ limits)


Maltoni et al, hep-ph/0309130

$$r \sim \Delta m_{\text{sol}}^2 / \Delta m_{\text{atm}}^2$$

G. Altarelli

$$\begin{aligned} 0.018 < r < 0.053 \\ |U_{e3}| < 0.23 \\ 0.30 < \tan^2 \theta_{12} < 0.64 \\ 0.45 < \tan^2 \theta_{23} < 2.57 \end{aligned}$$

for each model the λ, λ' values are optimised




The optimised values of λ are of the order of λ_C or a bit larger (moderate hierarchy)

model	$\lambda(= \lambda')$
A_{SS}	0.2
SA_{SS}	0.25
$H_{(SS,II)}$	0.35
$H_{(SS,I)}$	0.45
$IH_{(SS,II)}$	0.45
$IH_{(SS,I)}$	0.25

Results with see-saw dominance (updated in Nov. '03):

1 or 2 refer to models with 1 or 2 flavons of opposite ch.

With charges of both signs and 1 flavon some entries are zero



Errors are linear comb. of stat. and syst. errors (varying the extraction procedure: interval of ρ , real or complex)

H2 is better than SA, better than A, better than IH

G. Altarelli

Example: Normal Hierarchy

G.A., Feruglio, Masina

Note: not all charges positive
 \rightarrow det23 suppression

1st fam. 2nd 3rd

$$\begin{aligned} q(10): & (5, 3, 0) \\ q(\bar{5}): & (2, 0, 0) \\ q(1): & (1, -1, 0) \end{aligned}$$

$$\begin{aligned} q(H) &= 0, \quad q(\bar{H}) = 0 \\ q(\theta) &= -1, \quad q(\theta') = +1 \end{aligned}$$

In first approx., with $\langle \theta \rangle / M \sim \lambda \sim \lambda' \sim 0.35 \sim o(\lambda_c)$

$10_i 10_j$

$$m_u \sim v_u \begin{pmatrix} \lambda^{10} & \lambda^8 & \lambda^5 \\ \lambda^8 & \lambda^6 & \lambda^3 \\ \lambda^5 & \lambda^3 & 1 \end{pmatrix},$$

$10_i \bar{5}_j$

$$m_d = m_e^T \sim v_d \begin{pmatrix} \lambda^7 & \lambda^5 & \lambda^5 \\ \lambda^5 & \lambda^3 & \lambda^3 \\ \lambda^2 & 1 & 1 \end{pmatrix}$$

"lopsided"

$\bar{5}_i 1_j$

$$m_{\nu D} \sim v_u \begin{pmatrix} \lambda^3 & \lambda & \lambda^2 \\ \lambda & \lambda' & 1 \\ \lambda & \lambda' & 1 \end{pmatrix},$$

$1_i 1_j$


$$M_{RR} \sim M \begin{pmatrix} \lambda^2 & 1 & \lambda \\ 1 & \lambda'^2 & \lambda' \\ \lambda & \lambda' & 1 \end{pmatrix}$$

G. Altarelli **Note:** coeffs. 0(1) omitted, only orders of magnitude predicted

With no see-saw (m_ν generated directly from $L^T m_\nu L \sim$
is better than A

$\bar{5} \bar{H}$

[With no-see-saw H coincide with SA]



Note: we always include the effect of diagonalising charged leptons

G. Altarelli


Some distributions

IH2 NO-SS

$\lambda = \lambda' = 0.3$

We see that IH tends to predict maximal solar mixing angle θ_{12}

Only compatible because of ch. lepton diagonalisation



With data drifting away from maximal θ_{12} , IH is rapidly disfavoured (in U(1) models)




G. Altarelli

ch. lepton mixing small because m_e small



The main problem of Anarchy is U_{e3} (as expected)

In all models the distr. for $\tan^2\theta_{23}$ is flat

$\lambda=\lambda'=0.2$



$\lambda=\lambda'=0.35$



G.

The main advantage of SA vs A is for U_{e3}

$$r \equiv \frac{\Delta m_{sun}^2}{\Delta m_{atm}^2}$$





$$\Psi_5 \sim (2,0,0)$$

$$m_\nu \sim \begin{pmatrix} \lambda^4 & \lambda^2 & \lambda^2 \\ \lambda^2 & 1 & 1 \\ \lambda^2 & 1 & 1 \end{pmatrix}$$

$$\text{Det}_{23} \sim 0(1)$$

$$SA_{(NOSS)}, \lambda = 0.2$$

works when r is small enough by chance



Summing up:

- ν masses very small \rightarrow Majorana ν 's and see-saw mechanism
- ν masses are consistent with the standard way beyond the SM: SUSY and GUT's
- Recent exp progress:
 - Δm_{sol}^2 went closer to Δm_{atm}^2 \longrightarrow less hierarchy
 - smaller upper limit on absolute mass: $|m_3/m_2| \sim 6$
- Crucial issues:
 - LSND??
 - WMAP: $\sum m_\nu < 0.69$ eV
 - s_{13} small (how small?) disfavors anarchy
 - $s_{23} \sim$ maximal (too maximal?),
 $s_{12} \sim$ large not maximal disfavors inv. hierarchy
 - $0\nu\beta\beta$:
 - near bound? \longrightarrow degenerate ν 's
 - intermediate? \longrightarrow inverted hierarchy
 - small? \longrightarrow normal hierarchy
 - CP violation: still in the future \longrightarrow Looks simplest and fine

G. Altarelli