

Laboratori Nazionali di Frascati

LNF-62/93 (1962)

G. Barbiellini, G. Bologna, G. Diambrini, G. P. Murtas: MEASUREMENT OF THE POLARIZATION OF THE FRASCATI 1 GeV ELECTRON SYNCHROTRON γ -RAY BEAM FROM A DIAMOND CRYSTAL RADIATOR.

Estratto dal: Phys. Rev. Letters, 9, 396 (1962)

The other approximate method of solving Eq. (5) is that of Chew-Wick impulse approximation. The strong short-range interactions in configuration space will yield extremely slowly varying functions, $\tau_V(\vec{k})$, in momentum space. From Eq. (5a) we see then that if the momentum functions $X(\vec{k})$ (which in the impulse approximation are the transforms of the asymptotic functions φ , i.e., $\Omega = 1$) are peaked at $k = 0$, the major contributions to the integral come only from this region. Thus for large momentum transfers we can approximate $\tau_V(\vec{k} - \vec{\Delta})$ by $\tau_V(\vec{\Delta})$, and take it outside the integral as a multiplicative factor. The resulting integral over only the momentum functions $X(\vec{k})$ give the effects on the scattering amplitude of the "spectator" particle as in the usual formulation of the impulse approximation. Another advantage of our formalism is that it offers a relatively simple way of calculating corrections to the impulse approximation. Such corrections appear in a multiplicative rather than additive way if one uses $\chi = \Omega\varphi$ ($\Omega \neq 1$) in a distorted-wave approximation.

A forthcoming paper will give results for other applications of the theory (in particular to the scattering of electrons by hydrogen atoms) as well as expand on both the details of these calculations and the general formalism.

The author would like to thank Professor G.

Breit for bringing up the specific problem from which this approximation evolved, and Professor L. Durand, III, and Professor G. Rawitscher for many helpful discussions and comments.

*This research was supported by the U. S. Atomic Energy Commission and by the Air Force Office of Scientific Research.

¹G. F. Chew, Phys. Rev. 80, 196 (1952); G. F. Chew and G. C. Wick, Phys. Rev. 85, 636 (1952).

²M. Gell-Mann and M. L. Goldberger, Phys. Rev. 91, 398 (1953), Eq. (4.4). It should be noted that $\chi_a^{(+)}$ of their first term of this equation should read $\psi_a^{(+)}$. This example of a nonrearrangement process is used to illustrate conveniently the formulation of the two-potential problem in terms of the single-potential scattering amplitudes, and this formulation is easily extended to describe also the scattering of a particle by a bound two-body system.

³B. A. Lippmann, Phys. Rev. 102, 254 (1956).

⁴N. C. Francis and K. M. Watson, Phys. Rev. 93, 313 (1954).

⁵G. Breit and M. E. Ebel, Phys. Rev. 103, 679 (1956).

⁶F. C. Jobes, thesis, Yale University, 1962 (unpublished), and F. C. Jobes and J. A. McIntyre (to be published). The author would like to thank Professor McIntyre for advance communication of the data.

⁷W. Selove, Phys. Rev. 101, 231 (1956).

⁸G. F. Chew and M. L. Goldberger, Phys. Rev. 77, 470 (1950).

⁹K. R. Greider, Phys. Rev. 114, 786 (1959).

LNF-62/53

MEASUREMENT OF THE POLARIZATION OF THE FRASCATI 1-GeV ELECTRON SYNCHROTRON γ -RAY BEAM FROM A DIAMOND CRYSTAL RADIATOR

G. Barbiellini, G. Bologna, G. Diambrini, and G. P. Murtas

Laboratori Nazionali di Frascati del Comitato Nazionale per l'Energia Nucleare, Roma, Italy
(Received October 8, 1962)

In a previous Letter¹ we showed experimental evidence for a line spectrum in the 1-GeV bremsstrahlung γ -ray beam obtained using a diamond radiator. In this Letter we give some results of the calculations and measurements concerning the polarization of a 150-MeV photon line.

In high-energy electron bremsstrahlung, the radiated photons are emitted preferentially with a state of linear polarization parallel to the plane determined by the direction of the primary electrons and the recoil momentum of the nucleus. In a crystal only those recoil momenta are allowed which, in suitable units, are coincident with the reciprocal lattice vectors. Therefore the process is not symmetrical around the direction of the primary electron. It follows that the entire brems-

strahlung beam has a net polarization, with respect to a crystal plane.

The first calculation of this polarization was performed by Überall²; we have repeated it in order to take into account the discrete structure of the lattice planes. The importance of this structure has already been stated in other works.^{1, 3} A few numerical results of this calculation were given by us in reference 1. At the same time Überall also had recalculated the polarization, but his numerical data⁴ are not comparable with our experiment.

We define the polarization of the entire γ -ray beam as

$$P = (I_H - I_V) / (I_H + I_V), \quad (1)$$

where I_H and I_V represent the bremsstrahlung intensities for photons with polarization perpendicular and parallel, respectively, to the plane (\vec{p}_1, \vec{b}_1) , determined by the incident electron momentum \vec{p}_1 and a reciprocal lattice axis \vec{b}_1 . The choice of the indices H and V is due to the fact that the (\vec{p}_1, \vec{b}_1) plane is held vertical.

We obtained

$$I_H(\theta, x, E_1) - I_V(\theta, x, E_1) = 2(1-x)\psi_3^{0*}(\theta, \delta); \quad (2)$$

$$\psi_3^{0*}(\theta, \delta) = [(2\pi)^2/\Delta]4\delta^3 \sum_{\vec{g}} |F|^2 [\exp(-Ag^2)/(\beta^{-2} + g^2)^2] \times [(g_2^2 - g_3^2)/g_3^4\theta^4], \quad (3)$$

where E_1 and k are the electron and photon energy, respectively, $x = k/E_1$, and θ is the angle between the vectors \vec{p}_1 and \vec{b}_1 . The sum in (3) is extended over the reciprocal lattice vectors $\vec{g} = (0, g_2, g_3)$ which lie in the (\vec{b}_2, \vec{b}_3) plane; \vec{p}_1 lies in the (\vec{b}_1, \vec{b}_3) plane. The other quantities in Formula (3) were already defined in reference 1. In Formula (1) one has

$$I_H(\theta, x, E_1) + I_V(\theta, x, E_1) = I(\theta, x, E_1). \quad (4)$$

$I(\theta, x, E_1)$ is the bremsstrahlung intensity, already defined in reference 1.

The numerical values of Formula (1) were calculated for diamond at room temperature with the same input data as in reference 1; the only difference is that now \vec{p}_1 is parallel to the plane of the axes $\vec{b}_1 = (110)$ and $\vec{b}_3 = (001)$. The reason is that in this situation the polarization is very large. The continuous curve of Fig. 1 represents Formula (1) as a function of θ , for $E_1 = 1$ GeV and $k = 150$ MeV.

The measurement of the polarization depends upon the fact that in electron pair production from linearly polarized photons, the electrons are emitted preferentially in the plane of polarization (i.e., the plane determined by the direction of the incident photon \vec{k} , and the polarization vector $\vec{\epsilon}$). The experimental apparatus is about the same as described in reference 1. In the present experiment we chose a collimation of 1 mrad. The collimators were arranged in such a way to eliminate that part of the electron multiple traversals of the diamond radiator which spread out the γ -ray source in the horizontal direction. The source extension obtained in this way was ≈ 6 mm². The beam intensity after collimation was $\approx 10^{10}$ equivalent quanta/minute. The aluminum converter was 10^{-4} radiation length thick; in this way the mean square scattering angle of the electrons was much less than their most probable emission angle.




FIG. 1. Polarization of $k = 150$ -MeV photons in the entire bremsstrahlung beam of $E_1 = 1$ -GeV electrons, striking a diamond at room temperature. θ (in milliradians) is the angle between the momentum \vec{p}_1 of the incoming electron and the crystal axis [110]; \vec{p}_1 is parallel to the plane of the crystal axes [110], [001] (which is a vertical plane). $P = (I_H - I_V)/(I_H + I_V)$. The points represent the experimental results of Table I. They are normalized to the theoretical curve at $\theta = 11.6$ mrad. The error is statistical.

The principal change with respect to the old apparatus was made in the system of scintillation counters which detected symmetrical pairs. The present disposition is shown in Fig. 2; the scintillators A_2 and A_3 , 10 cm in height, were placed in the e^- branch of the pair, while the scintillators A_1 and S_1 were placed in the e^+ branch. A_1 has the same height as A_2 and A_3 ; S_1 is only 0.5 cm in height. All the scintillators accepted an electron momentum spread $\Delta p/p = 8.5\%$.

By means of the triple coincidence $S_1 A_2 A_3$ we detected the symmetrical pairs in which one branch is emitted at any angle with respect to the incident photon, and the other branch is emitted in the neighborhood of the spectrometer median plane MP , which is horizontal and contains S_1 . On the other hand, by means of the triple coincidence A_1, A_2, A_3 we detected the symmetrical pairs with the two branches emitted at any angle.




FIG. 2. Sketch of the analyzing device. \vec{k} is the direction of the incoming photon. $\vec{\epsilon}$ is the direction of the polarization vector. C is the aluminum converter. e^+ and e^- are the two branches of the symmetrical pair. \vec{B} is the direction of the magnetic field of the pair spectrometer. The height of the scintillators A_1, A_2, A_3 is 10 cm; that of S_1 is 0.5 cm. MP is the median plane of the pair spectrometer.

The counting rate of $S_1 A_2 A_3$ depends on the position of the polarization plane; if this is horizontal, the counting rate is larger than if it is vertical. Olsen and Maximon⁵ calculated the pair production cross section from linearly polarized photons, integrated over the emission angles of one branch and differential in the emission angles of the other branch and, of course, in the electron energy. In order to adjust this cross section to our experimental situation, we integrated it over the emission angle in the horizontal plane. We took into account also the electron multiple scattering in the converter as well as the finite size of the beam at the converter (its diameter was 1.3 cm), and the finite height of the scintillator S_1 . We obtained the asymmetry ratio

$$R = (d\sigma_{||} - d\sigma_{\perp}) / (d\sigma_{||} + d\sigma_{\perp}) = 10.1\%, \quad (5)$$

for the photon energy $k = 150$ MeV; in Formula (5) $d\sigma_{||}$ and $d\sigma_{\perp}$ represent the pair production cross section in the given experimental situation from photons polarized parallel and perpendicular to the spectrometer median plane, respectively. R

approaches zero when the converter thickness is more than 10^{-2} radiation length. R is also decreased when the size of the γ -ray beam is increased. For this reason the 1-mrad angle of acceptance of the collimators was an upper limit.

In order to measure the polarization of the γ -ray beam we fixed the nominal energy of the photons at $k = 150$ MeV, by the proper choice of the spectrometer magnetic field, and we measured the ratio of coincidence counts

$$\rho = S_1 A_2 A_3 / A_1 A_2 A_3, \quad (6)$$

taking into account also delayed coincidences; background is negligible. We considered the coincidence $A_1 A_2 A_3$ only as a monitor in order to measure at constant number of photons. ρ was measured for several values of θ , with \vec{p}_1 held parallel to the plane of the crystal axes $\vec{b}_1 = (110)$ and $\vec{b}_3 = (001)$, which is a vertical plane.

The relationship between Formulas (6) and (1) is

$$\rho = \rho_0 (1 + RP), \quad (7)$$

from which we see that the constant ρ_0 represents the values of ρ for $P = 0$. Since for several reasons we could not yet measure the absolute value of the polarization, we assumed that the γ -ray beam was unpolarized at $\theta = \theta_0 = 5.8$ mrad, where the theoretical curve of Fig. 1 passes through the zero value. At this angle we measured ρ_0 . Thus we made relative measurements, normalized at $\theta = \theta_0$.

Since the maximum value of P , on the basis of the theoretical results, is 50% and the value of R is 10%, the maximum relative difference between ρ and ρ_0 is 5%. Therefore we had to take particular care in measuring this small effect. In order to eliminate the bias introduced by the instability of the electronics, each measurement of ρ was followed by a measurement of ρ_0 ; we rejected those measurements in which the values of ρ_0 differed more than the statistical error of $\approx 1\%$. We encountered another difficulty in the fact that the energies of the photons detected from the two triple coincidences $S_1 A_2 A_3$ and $A_1 A_2 A_3$ must be coincident within 1 MeV. In fact, we see from Fig. 1(b) of reference 1 the large rate with which the number of photons in the spectrum varies by changing the energy in the neighborhood of the first discontinuity; a change of 1 MeV in the photon energy will cause a change of some percent in the number of counts, and this is just of the order of the variation of ρ .

If we could have rotated the crystal by 90° in its

plane it would have been possible to measure an effect twice as large and, besides, we would have been able to make an absolute measurement. However, due to experimental difficulties, this was not possible. We hope to overcome this difficulty shortly.

The experimental results are summarized in Table I. In the first column the value of the angle θ in milliradians is given; in the second column we give the weighted average

$$\langle \rho/\rho_0 \rangle = \sum \frac{1}{\mu_i^2} \frac{\rho^{(i)}}{\rho_0^{(i)}} / \sum \frac{1}{\mu_i^2}, \quad (8)$$

where μ_i is the error propagated in $\rho^{(i)}/\rho_0^{(i)}$ by the statistical errors of the counts $S_1 A_2 A_3$ and $A_1 A_2 A_3$. In the third column we give the error of (8), i.e.,

$$\mu_\theta = [\sum (1/\mu_i^2)]^{-1/2},$$

multiplied by 10^3 . In the fourth column we give the polarization P , calculated by means of the formula

$$\frac{\langle \rho/\rho_0 \rangle}{\langle \rho_1/\rho_0 \rangle} = \frac{1+RP}{1+RP_1}, \quad (9)$$

ρ_1 being the value of ρ obtained experimentally for $\theta = \theta_1 = 11.6$ mrad; $P_1 = 16.0\%$ is the theoretical value of the polarization at the same angle. The reason for which, in presenting the results, we prefer to renormalize the data at $\theta = \theta_1$ is that at

Table I. Experimental results obtained in the measurement of the polarization.

θ (mrad)	$\langle \rho/\rho_0 \rangle$	$10^3 \mu_\theta$	P (%)
2.3	0.998	7.0	2 ± 7
2.9	1.002	9.6	6 ± 10
4.9	0.990	8.9	-6 ± 9
5.4	1.001	9.5	5 ± 9
5.8	1.000	4.5	4 ± 4
7.0	1.026	5.9	30 ± 6
7.4	1.049	4.8	52 ± 5
7.9	1.033	6.8	37 ± 7
9.0	1.029	7.1	33 ± 7
11.6	1.012	6.1	16 ± 6

this angle the influence of the electron multiple scattering in the diamond is negligible.

In the same column, following the value of the polarization, its error is quoted. This is $\approx 10\mu_\theta$; i.e., in propagating the error by means of (9), we disregard the error of $\langle \rho_1/\rho_0 \rangle$. This is because $\langle \rho_1/\rho_0 \rangle$ is used only to normalize the theoretical and experimental behavior of P .

The data of the fourth column are displayed also in Fig. 1. The theoretical curve is not yet corrected for the electron multiple scattering in the diamond radiator. Nevertheless, one can see how the variations of the polarization, as measured between an angle θ and the angle θ_1 , make a good fit with the theoretical curve, at least for $\theta > 6$ mrad; below this angle multiple scattering is important, so the experimental values do not agree well with the theoretical ones.

By inspection of Fig. 1 we see that the polarizing device, as well as the analyzing one, works satisfactorily, because for a change of only $\Delta\theta = 1.6$ mrad in the neighborhood of the discontinuity, we detected a change in polarization of

$$\Delta P = (48 \pm 7)\%.$$

On the other hand, after replacement of the converter 10^{-4} radiation length thick with another one, 10^{-2} radiation length thick, the values of ρ and ρ_0 , relative to the preceding $\Delta\theta$, appeared coincident within the statistical error.

The big advantage in using a crystal instead of an amorphous radiator as a polarizer was indicated by Überall in his first work.² At this date we can say that the advantage is even greater than that expected at the time of the initial work.

We wish to thank Professor J. DeWire for many valuable discussions and Dr. B. Antonini for the integration of the cross section in Formula (5).

¹G. Barbiellini, B. Bologna, G. Diambrini, and G. P. Murtas, Phys. Rev. Letters 8, 454 (1962); 9, 46(E) (1962).

²H. Überall, Phys. Rev. 107, 223 (1957).

³G. Barbiellini, G. Bologna, G. Diambrini, and G. P. Murtas, Phys. Rev. Letters 8, 112 (1962).

⁴H. Überall, Z. Naturforsch. 17, 332 (1962). Überall's numerical data are relative to only one reciprocal lattice row; thus they are different from ours, which were obtained by a sum over all reciprocal lattice points.

⁵H. Olsen and L. C. Maximon, Phys. Rev. 114, 887 (1959).