

COMITATO NAZIONALE PER L'ENERGIA NUCLEARE
Laboratori Nazionali di Frascati

LNF-71/66
4 Ottobre 1971

G. Parisi and F. Zirilli: ANGULAR CORRELATIONS OF THE
DECAY PRODUCTS OF TWO HEAVY LEPTONS.

LNF-71/66
4 Ottobre 1971

G. Parisi and F. Zirilli^(x): ANGULAR CORRELATIONS OF THE
DECAY PRODUCTS OF TWO HEAVY LEPTONS.

It is well known that there is no experimental evidence against the existence of a third heavy lepton, if its mass is greater than the kaon mass.

If this particle is not too heavy it can be produced very easily in colliding beams experiments: the cross section for $e^+e^- \rightarrow X^+X^-$ is:

$$(1) \quad \sigma(E) = \frac{\alpha^2 \pi}{3 E^2} \left(1 + \frac{M_x^2}{2 E^2}\right) \sqrt{1 - \frac{M_x^2}{E^2}}$$

If the heavy lepton has the same weak interaction as the electrons or the muon, its branching ratio can be computed.

The one prong decays: $X^+ \rightarrow \mu^+ \nu \nu$, $e^+ \nu \nu$, $\pi^+ \nu$, $K^+ \nu$, turn out to be dominant⁽¹⁾, so the main process in colliding beams is: $e^+e^- \rightarrow 2$ charged + neutral.

For analyzing the data it is interesting to know the angular distribution of the produced charged particles. We denote by E the beam energy, by θ the angle between the direction of the produced heavy lepton and the beam, by ω^+ the angle between the direction of X^+ and of its charged decay product in the c.m. frame of the heavy lepton and by ω^- the same angle for X^- .

(x) - Istituto di Matematica e Fisica dell'Università di Roma.

2.

One finds that :

$$\begin{aligned}
 & \frac{d\sigma}{d\cos\theta, d\cos\omega^- d\cos\omega^+} \propto \\
 (2) \quad & \propto [1 + \cos^2\theta] \left\{ 1 + \alpha^+ \alpha^- \cos(\omega^+ + \omega^-) \right\} + \\
 & + \frac{M}{E} 2 \sin\theta \cos\theta \alpha^+ \alpha^- \sin[\omega^+ + \omega^-] + \\
 & + \left(\frac{M}{E}\right)^2 \sin^2\theta \left\{ 1 - \alpha^+ \alpha^- \cos(\omega^+ + \omega^-) \right\}
 \end{aligned}$$

where α^+ and α^- are the asymmetry parameters of the decay of X^+ and X^- .

Their values are:

$$\begin{aligned}
 (3) \quad & \alpha = 1 && \text{for the decays:} \\
 & && X^+ \rightarrow \pi^+ \nu, \quad X^- \rightarrow K^- \nu \\
 & \alpha = \frac{4 E_1 - M_x}{3 M_x - 4 E_1} && \text{for the decays:} \\
 & && X^+ \rightarrow \mu^+ + \nu + \nu, \quad X^- \rightarrow e^- + \nu + \nu
 \end{aligned}$$

where E_1 is the energy of the e or μ in the center of mass of the X.

Formula (2) can be computed from the helicity amplitudes for the production and for the decay of an heavy lepton, using the methods of ref. (2).

We thank Dr. M. Testa for discussions.

REFERENCES. -

- (1) - H. B. Thacker and J. J. Sakurai, Phys. Letters 36 B, 103 (1971).
- (2) - G. Parisi and F. Zirilli, A simple method for computing electrodynamic processes of high order, Frascati Internal Report (1971), to be published.