

LNF-72/51
6 Giugno 1972

G. Parisi and L. Peliti: CRITICAL INDICES FOR THE SPHERICAL
MODEL FROM CONFORMAL COVARIANT SELF CONSISTENCY
CONDITIONS, -

LNF-72/51
6 Giugno 1972

G. Parisi and L. Peliti^(x): CRITICAL INDICES FOR THE SPHERICAL
MODEL FROM CONFORMAL COVARIANT SELF CONSISTENCY
CONDITIONS^(o).

ABSTRACT. -

Critical indices for the spherical model are computed in arbitrary dimension space. Well known exact results are recovered in three dimensional space. The key tools used are conformal covariant self consistency conditions.

(x) - Queen Mary College, Mile End Rd., London E. 1. (U. K.); Present address: Istituto di Fisica dell'Università, Roma.

(o) - This reserach has been partially sponsored by CNR GNSM.

2.

In recent times it has been suggested that changing the dimension of the space may be an useful tool to compute critical indices⁽¹⁾. One finds that the critical indices for the spherical model in a $4-\epsilon$ dimensional space are $\nu = \frac{1}{2} + \frac{\epsilon}{4} + \frac{\epsilon^2}{8} + O(\epsilon^3)$ $\eta = 0 + O(\epsilon^4)$.

In this letter we extend this result to arbitrary dimension space using conformal covariant self consistency conditions^(2,3). The equivalence between our and Wilson's approach was recently proved by G. Mack⁽⁴⁾. The critical indices turn out to be: $\nu = \frac{1}{2-\epsilon}$; $\eta = 0$.

This expression is equal to Wilson's one for small $\epsilon = 4-d$ and yields the right values $\eta = 0$, $\nu = 1$ for $\epsilon = 1$ ⁽⁵⁾.

The self consistency conditions are written in Fig. 1: G is the "one particle" propagator, D is the "photon" propagator, V is the pho-




FIG. 1 - Self consistency conditions for the vertex, the one¹particle and the "photon" propagators.

non-particle vertex, K_1 and K_2 are the two different Bethe Salpeter kernels and the dotted lines mean momentum differentiation^(2, 6). These self consistency conditions are derived following ref. (1) using the Landau Ginsburg functional for the free energy as a starting point and introducing an auxiliary "phonon" field⁽⁷⁾. We need the solution of the equations in the limit $N \rightarrow \infty$. The factor N plays the rôle of a multiplicity factor.

The form of the propagator and vertices at the critical temperature follows from conformal invariance requirements:

$$G(x) = \frac{1}{|x|^{d+\eta-2}}, \quad D(x) = \frac{1}{|x|^{2d-2/\nu}}$$

(1)

$$V(x, y, z) = \frac{g \Gamma^{-1}(2 + \frac{1}{\nu} - \eta - d)}{|x-z|^{1/\nu} |z-y|^{1/\nu} |x-y|^{d+2-\eta-1/\nu}}$$

The normalization of the coupling constant is chosen in such a way that the Fourier transform of the vertex is always finite, for any value of the indices.

The kernels K_1 and K_2 have a standard diagrammatical expansion in terms of G , D and V .

Letting (1) in the self consistency equations we find a system of three equations in three unknowns whose solution yields the value of the critical indices.

We look now for solution of the type $\frac{1}{\nu} = 2 - \varepsilon + \Delta_\nu$, where η and Δ_ν are small parameters.

In this limit we find that the three self consistency equations are respectively:

4.

$$(2) \quad \frac{g^2}{\Delta \nu - \eta} + \dots = C_1; \frac{g^4}{\Delta \nu - \eta} + \dots = C_2 \eta; \frac{Ng^4}{\Delta \nu - \eta} + \dots = C_3$$

where C_1, C_2, C_3 are some constants dependent on the dimension of the space. These equations are obtained taking only the simplest diagrams in the self consistency conditions; a more careful analysis shows that neglected terms do not change our conclusions⁽⁷⁾.

In the $N \rightarrow \infty$ limit we find that the solution is $\nu = \frac{1}{2 - \epsilon'}$ $\eta = 0$. Any ambiguity in the solution of the equations is removed by the condition that in the 4 dimensional limit we recover the classical indices.

The simplicity of our result is due to the compensation of the singularities of the denominator with the large N limit.

The authors are grateful to Prof. G. Mack for useful discussions.

REFERENCES. -

- (1) - K. Wilson and M. E. Fisher, Phys. Rev. Letters 28, 240 (1972); K. Wilson, Phys. Rev. Letters 28, 548 (1972).
- (2) - G. Parisi and L. Peliti, Lett. Nuovo Cimento 2, 627 (1971).
- (3) - A. A. Migdal, Phys. Letters 37B, 386 (1971); G. Mack and I. T. Todorov, Trieste preprint IC/71/139 (1971); G. Mack and K. Symanzik, Desy preprint 72/19 (1972).
- (4) - G. Mack, Expansion around canonical dimensions in conformal invariant quantum field theory, Bern preprint (1972).
- (5) - H. E. Stanley, Phys. Rev. 176, 718 (1968), and references therein.
- (6) - G. Parisi, On self consistency conditions, Frascati preprint LNF-72/35 (1972); Lett. Nuovo Cimento, to be published.
- (7) - A. A. Migdal, Soviet Phys. -JETP 28, 1036 (1969).
- (8) - G. Parisi and L. Peliti, Conformal invariance and second order phase transitions, in preparation.