

To be submitted to  
Physics Letters B

ISTITUTO NAZIONALE DI FISICA NUCLEARE  
Laboratori Nazionali di Frascati

LNF-76/39(P)  
6 Luglio 1976

M. E. Biagini, D. Bisello, B. Esposito, F. Felicetti, A. Marini,  
P. Monacelli, A. Nigro, M. Nigro, L. Paoluzi, L. Pescara, G.  
Piano Mortari, F. Ronga, A. Sciubba, F. Sebastiani, and F.  
Vanoli: SEARCH FOR NARROW RESONANCES IN  $e^+e^-$   
ANNIHILATION INTO HADRONS AT ADONE IN THE MASS  
REGION  $2.5-3.0 \text{ GeV}/c^2$ .

Laboratori Nazionali di Frascati  
Servizio Documentazione

LNF-76/39(P)

6 Luglio 1976

M. E. Biagini<sup>(x)</sup>, D. Bisello<sup>(o)</sup>, B. Esposito, F. Felicetti, A. Mari-  
ni, P. Monacelli<sup>(x)</sup>, A. Nigro<sup>(+)</sup>, M. Nigro<sup>(o)</sup>, L. Paoluzi<sup>(x)</sup>, L. Pe-  
scara<sup>(o)</sup>, G. Piano Mortari<sup>(x)</sup>, F. Ronga, A. Sciubba<sup>(x)</sup>, F. Sebastiani<sup>(x)</sup>  
and F. Vanoli<sup>(+)</sup>: SEARCH FOR NARROW RESONANCES IN  
 $e^+e^-$  ANNIHILATION INTO HADRONS AT ADONE IN THE MASS  
REGION 2.5 - 3.0 GeV/c<sup>2</sup>.

ABSTRACT. -

The mass region 2.5 to 3.0 GeV/c<sup>2</sup> has been searched for evidence of  $\psi$ -like narrow resonances in  $e^+e^-$  annihilation into hadrons. No evidence for new narrow resonances has been found with a sensitivity of about the 10% of the integrated cross section of the  $J/\psi(3100)$ .

A systematic search for narrow resonances in the total hadronic cross section in  $e^+e^-$  annihilation over the center - of - mass energy range  $W = 1.9 - 3.1$  GeV has been performed at Adone, the Frascati 2x1.5 GeV storage ring, with the MEA magnetic detector<sup>(1)</sup>. In this paper we present the results for the energy region  $W = 2.5 - 3.0$  GeV. Previous data at  $W = 1.9 - 2.5$  GeV and  $W = 3.0 - 3.1$  GeV have already been published<sup>(2)</sup>.

---

(x) - Istituto di Fisica dell'Università di Roma, and Istituto Nazionale di Fisica Nucleare, Sezione di Roma.

(o) - Istituto di Fisica dell'Università di Padova, and Istituto Nazionale di Fisica Nucleare, Sezione di Padova.

(+) - Istituto di Fisica dell'Università di Napoli, and Istituto Nazionale di Fisica Nucleare, Sezione di Napoli.

A resonance in the reaction:



is said to be narrow if its total width is negligible with respect to the FWHM machine energy resolution  $\Gamma_W$ . At Adone  $\Gamma_W$  is given by the following relation<sup>(3)</sup>:

$$\Gamma_W (\text{MeV}) = 0.32 W^2 (\text{GeV}^2), \quad (2)$$

and consequently we have choosen to explore our energy region by 1 MeV steps.

The total collected luminosity was  $\mathcal{L} = 86.6 \text{ nb}^{-1}$ . No statistically significant evidence for new narrow resonances has been found. The  $e^+ e^- \rightarrow e^+ e^-$  Bhabha scattering at small angles ( $3^\circ - 6^\circ$ ) as measured by the Adone machine group in a different interaction region has been used to provide a fast relative luminosity monitor. The  $e^+ e^- \rightarrow e^+ e^-$  Bhabha scattering at large angles, as seen by the MEA apparatus, has been used to provide an absolute normalization of the luminosity.

The MEA apparatus is described in some detail in ref. (1) and it consists basically of a set of cilindrical optical spark-chambers inserted into a soleinoidal magnet transverse to the beam orbit, and of a number of hodoscopes of scintillation counters. The trigger of the apparatus requires at least two charged particles, with a minimum kinetic energy of 130 MeV (if pions) in the two opposite halves of the apparatus. We call multihadronic events those events which show the following features:

- a) more than two particles are detected;
  - b) two particles are observed with an acoplanarity angle  $\Delta\varphi \geq 10^\circ$ .
- In addition, all events must fulfil the following requirements:
- c) correct timing with respect to the bunch-bunch collisions;
  - d) the charged prongs must converge in the interaction region; an acceptance of  $\pm 1 \text{ cm}$  in the radial dimensions of the source, as measured by a set of multiwire proportional chambers<sup>(1)</sup>, has been imposed.


Condition b) practically eliminates all contamination from collinear events and cosmic rays, and strongly reduces the contribution from the

two photon processes ( $e^+e^- \rightarrow e^+e^-e^+e^-$  and  $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ ). The background from beam-gas interaction is also strongly reduced by the acoplanarity cut. It should be observed, in addition, that such background is basically energy independent and thus it does not affect the results concerning fine structures in the energy dependence of the cross section. The same argument is valid as far as the detection efficiency of the apparatus for multihadronic events is concerned. Indeed, the detection efficiencies for each final state and the average efficiency are almost independent of energy in the mass range being considered.

In Fig. 1 the detected yield vs. total energy is shown. The  $J/\psi$  (3100) peak as detected by the apparatus is also reported for comparison. Quoted errors are statistical only.

No significant structures can be seen in the total mass<sup>range</sup> explored. In order to derive an upper limit for the production cross-section of new resonances times their branching ratio into more than two hadrons, we have proceeded as follows.

A narrow resonance would be observed in the excitation curve as a gaussian bump with a width equal to the energy spread of the machine, distorted by radiative effects. Since in the explored energy range ( $W = 2.5 - 3.0$  GeV) the machine spread  $\Gamma_W$  varies from 2.1 to 2.8 MeV, the evaluation of the upper limit of the integrated yield ( $Y_R^{Int}$ ) from a narrow resonance at the energy  $W_o$  has been performed based on the number of events contained in all possible three consecutive energy bins, viz. in any 3 MeV wide energy intervals centered at  $W_o; (N(W_o))$ . Over such energy interval the fractional yield from a narrow resonance gives a contribution which can range from 60% to 74% of the total yield, depending on  $\Gamma_W$ . From the observed rate  $N(W_o)$ , and using Poisson statistics we have evaluated  $N^*(W_o)$ , the upper limit (with 90% probability) to the true rate. The maximum integrated yield from a possible narrow resonance  $Y_R^{Int}$  is related to  $N^*(W_o)$  by the relation:



$$N^*(W_0) = \sum_{i=1}^3 \left[ Y_{NR} + \frac{Y_R^{Int}}{\sqrt{2\pi} \sigma_W} e^{-\frac{(W_i - W_0)^2}{2\sigma_W^2}} \right] \mathcal{L}_i \quad (3)$$

where  $W_1 = W_0 - \Delta W$ ,  $W_2 = W_0$ ,  $W_3 = W_0 + \Delta W$  with  $\Delta W$  equal to the step of the energy scan (1 MeV);  $Y_{NR}$  is the mean value of the observed yield due to non resonant hadronic production and  $\mathcal{L}_i$  is the luminosity corresponding to the energy bin involved. From eq. (3) one can derive  $Y_R^{Int}$  and work out the ratio  $Y_R^{Int}/Y_{J/\psi}^{Int}$  relative to the measured integrated yield of the  $J/\psi$ . Assuming the detection efficiency of the apparatus to be equal for the  $J/\psi$  and for the hypothetical resonance at  $W_0$ , this ratio is equal to the ratio between the integrated cross sections, and it can be used to derive an upper limit for the hadronic cross-section of the new resonance.

The results of this procedure are quoted in Table I. The region  $2520 \leq W \leq 3000$  MeV has been split for convenience in five intervals, for which all data have been analysed together to produce single upper limits.

#### Acknowledgements:

We would like to express our deep gratitude to the Adone Staff Group for the very efficient running of the machine achieved in the delicate conditions required by this experiment.

TABLE I

Energy region (MeV)	$\sigma_R^{\text{Int}} / \sigma_{J/\psi}^{\text{Int}}$ upper limits (90% c.l.)
2520 - 2599	0.06
2600 - 2699	0.12
2700 - 2799	0.07
2800 - 2899	0.07
2900 - 3000	0.14

**REFERENCES. -**

- (1) - W. W. Ash et al., Lett. Nuovo Cimento 11, 705 (1974).
- (2) - B. Esposito et al., Phys. Letters 58B, 478 (1975).
- (3) - M. Bassetti, Adone Internal Report E-15 (1974).