

Presented at the Rome
Meeting of ECFA-LEP
Working Group (1978)

ISTITUTO NAZIONALE DI FISICA NUCLEARE
Laboratori Nazionali di Frascati

LNF-78/53(P)
25 Novembre 1978

A. Turrin: GENERALIZED FROISSART-STORA FORMULA FOR
FAST TUNE SHIFTING DURING ACCELERATION.

INFN Laboratori Nazionali di Frascati
Servizio Documentazione

LNF-78/53(P)
25 Novembre 1978

A. Turrin: GENERALIZED FROISSART-STORA FORMULA FOR
FAST TUNE SHIFTING DURING ACCELERATION

(Presented at the Rome Meeting of ECFA-LEP Working Group, to
the Specialized Study Group on Beams Polarization - Rome, Novem-
ber 23-24, 1978).

Abstract:

We have derived a generalized version of the Froissart - Stora
formula, which allows one to calculate the effect, on the polarization,
of the finite Q-jump magnitude, ΔQ , in crossing rapidly an intrinsic
depolarization resonance during acceleration.

1. - INTRODUCTION.

The forthcoming possibility of routinely accelerating polarized electron beams in the Bonn Synchrotron ¹ makes it worth-while to reanalyze the dynamics of depolarization in crossing the so-called intrinsic spin-resonances. Jumping these resonances for protons by rapidly changing at proper times the vertical betatron wavenumber, Q , is presently unique, standard practice at the Argonne ZGS ², and the same technique has been considered for the CERN PS ³, the AGS ⁴, the FNAL booster ⁴ and the KEK Proton Synchrotron ⁵.

Since it seems desirable to treat the effect, on polarization, of the finite Q -jump magnitude, ΔQ , in crossing rapidly an intrinsic depolarization resonance during acceleration, in the present note we give the derivation of a generalized version of the well known Froissart-Stora formula ⁶. This generalized formula is anticipated here below:

$$S_{z(t \rightarrow + \infty)} = 2 \left\{ \sinh(p\pi) / \sinh(r\pi) \right\}^2 - 1. \quad (1)$$

Here,

$$p = (1/2) (\Delta Q/2)^2 / Q' \quad \text{and} \quad r = p \left[1 + (\varepsilon/(\Delta Q/2))^2 \right]^{1/2}. \quad (2a, b)$$

In eqs. (2a, b), $Q' = \dot{Q}/\omega_0$ is the rate of change (per radian around the accelerator) of Q at the resonance crossing ($\dot{} \equiv d/dt$; ω_0 denotes the particle's angular velocity), ΔQ is the range of the Q -jump, $\varepsilon = \omega/\omega_0$ (ω is the width ⁶ of the resonance for a selected group of particles having the same amplitude of vertical betatron oscillations), and the assumption $S_{x(t \rightarrow - \infty)} = S_{y(t \rightarrow - \infty)} = 0$, $S_{z(t \rightarrow - \infty)} = 1$ is implicit.

Eq. (1) is valid for any magnitude of ΔQ , and allows one to calculate the resulting depolarization in cases where the Froissart and Stora model is inadequate, namely when Teng's condition ⁷ is violated. We recall that Teng's condition demands that \dot{Q} remains constant during the whole range of time $-(\omega_0 \dot{Q})^{-1/2} < t < +(\omega_0 \dot{Q})^{-1/2}$ where the resonance is effective, so that there exist cases where the applied range ΔQ may be too small for Teng's condition to be considered satisfied.

We also recall that the effective residue polarization can be cal-

culated by averaging S_z over the vertical betatron amplitude distribution. $(t \rightarrow + \infty)$

2. - ADIABATIC FOLLOWING SOLUTIONS BEFORE AND AFTER THE PASSAGE OF THE RESONANCE TAIL.

Fig. 1 shows the direction of the polarization vector \bar{S} (for a selected group of particles having the same amplitude of vertical betatron oscillations) at different times during the acceleration stages and the Q-jump stage. In the rotating reference frame, one may expect the following sequence of events to happen during normal acceleration, as the pulsed quadrupoles are subsequently turned on in the vicinity of the resonance, and eventually, i. e., at $t \rightarrow + \infty$, after the resonance is passed.

Sufficiently far from resonance, $\bar{S}_{(t \rightarrow - \infty)}$ is oriented along the main field direction, \bar{k} . When approaching the resonance, the effective field, $\omega \bar{j} + \dot{\chi} \bar{k}$ ($\dot{\chi} \gg \omega$), around which \bar{S} precesses, starts turning slowly away from the vertical, and \bar{S} will follow this change in orientation adiabatically⁸. Thus, before the occurrence of the Q-jump pulse, the vertical component, S_z , of \bar{S} is given by⁸

$$S_z(\gamma < \gamma_{res}) = 1 / \left[1 + (\omega / \dot{\chi})^2 \right]^{1/2}, \quad (3)$$

where $\dot{\chi} = \omega_0 G(\gamma - \gamma_{res})$ is the detuning⁶ ($\gamma = E/(mc^2)$ and G is the gyromagnetic anomaly).

Suppose now that when $\dot{\chi}$ becomes equal to $\omega_0 \Delta Q/2$, and, consequently, when

$$S_z(\gamma < \gamma_{res}) = (S_-)_z, \quad (3a)$$

with

$$(S_-)_z = p/r \quad (4)$$




FIG. 1 - Vector diagram, in the (y, z) plane, showing the variations of \bar{S} during the whole time interval $-\infty < t < +\infty$ in the case where a Q-jump is applied in the vicinity of an intrinsic spin-resonance. See text for the meaning of the symbols.

$$|\bar{S}(t \rightarrow -\infty)| = |\bar{S}_-| = |\bar{S}_+| = 1. \quad |\bar{S}_{+av}| = S_z(t \rightarrow +\infty)$$

(see Fig. 1), the pulsed quadrupoles are switched on. Clearly \bar{S}_- will start to precess around a rapidly varying effective field. Thus, after the resonance tail is passed through, the overall effect of this process is that it will lead to a polarization vector \bar{S}_+ precessing around the effective field $\omega \bar{j} - \omega_0 (\Delta Q/2) \bar{k}$. Let be $\bar{S}_{+ \text{av}}$ the time-average value of \bar{S}_+ . $(S_{+ \text{av}})_z$ will be expressed in subsequent paragraph 3, on the basis of a suitable model.

Eventually, i. e., over the remaining part of the acceleration cycle, $\bar{S}_{+ \text{av}}$ undergoes an adiabatic process of following the direction of the effective field, and slowly aligns along the \bar{k} direction as $\dot{\chi} \rightarrow -\infty$. Thus, we may write

$$S_{z(t \rightarrow +\infty)} = (S_{+ \text{av}})_z \quad r/p \quad . \quad (5)$$

3. - JOINING TOGETHER OF THE ADIABATIC FOLLOWING SOLUTIONS.

We now look for a relation between $(S_{+ \text{av}})_z$ and $(S_-)_z$ in order to derive a convenient asymptotic form for $S_{z(t \rightarrow +\infty)}$, i. e., eq. (1).

As a simple model which exhibits sweeping of $\dot{\chi}$ through zero, let

$$\omega = \text{constant} \quad \text{and} \quad \dot{\chi} = -\delta \tanh(\alpha t), \quad (6a, b)$$

where α and δ are assumed to be constant parameters. Clearly,

$$|\dot{\chi}_{t \rightarrow +\infty}| = \delta = \omega_0 \Delta Q/2 \quad (7a)$$

and

$$|\ddot{\chi}(0)| = \alpha \delta = \omega_0 \dot{Q}. \quad (7b)$$

Once the model above is adopted, the object is to solve the con-

ventional coupled equations of motion ⁶

$$2i\dot{f} = \omega g \exp(i\chi), \quad 2i\dot{g} = \omega f \exp(-i\chi) \quad (8a, b)$$

(which can be derived by considering the Schrödinger equation for a spin 1/2 particle interacting with an external field) written, as done above, in the rotating perturbing field approximation. ($gg^* + ff^* = 1$ is our normalization condition).

Thus, in the following, the time-average value of the transition probability,

$$(gg^*)_{av} \equiv (1/2) \left\{ 1 - (S_{+av})_z \right\}, \quad (9)$$

in the long-time limit will be calculated for a particle initially in the state given by eq. (4), by inserting eqs. (6) in eqs. (8) and solving these equations.

The first step consists of decoupling the spin-motion equations. We have, e. g., for g ,

$$\ddot{g} - i\delta \tanh(\alpha t) \dot{g} + (\omega/2)^2 g = 0. \quad (10)$$

Eq. (10) is easily transformed into the equation

$$x^2(1-x)g'' + x \left\{ (1+ip) + (-1+ip)x \right\} g' + (q/2)^2(1-x)g = 0 \quad (11)$$

by the transformation

$$x = -\exp(2\alpha t), \quad (' \equiv d/dx; \quad '' \equiv d^2/dx^2). \quad (12)$$

The dimensionless constants p and q are given by

$$p = \delta/(2\alpha) \quad \text{and} \quad q = \omega/(2\alpha). \quad (13a, b)$$

On introduction of the new function y by the substitution ⁹

$$g = x^\lambda y \quad , \quad (14)$$

Eq. (11) becomes ⁹

$$x(1-x)y'' + \{c - (a+b+1)x\} y' - aby = 0 \quad , \quad (15)$$

The constants λ , c , a and b are

$$2\lambda_{\pm} = -ip \pm ir; \quad c_{\pm} = 1 \pm ir; \quad r = (p^2 + q^2)^{1/2}; \quad (16a, b, c)$$

$$a_+ = -ip + ir; \quad b_+ = -ip; \quad (16d, e)$$

$$a_- = -ip; \quad b_- = -ip - ir. \quad (16f, g)$$

Eq. (15) is the well know hypergeometric differential equation ¹⁰. Thus, in the vicinity of $x = -0$ (i. e. at $t \rightarrow -\infty$) the solution of Eq.(11) which satisfies the boundary condition (4) can be written ¹¹ as follows:

$$g = D(-x)^{-ip/2-ir/2} F(-ip-ir, -ip; 1-ir; x) \quad . \quad (17)$$

Here, D is the proper integration constant, namely,

$$DD^* = gg^*_{t \rightarrow -\infty} \equiv (gg^*)_- \equiv (1/2) \{1 - (S_-)_z\} = (1-p/r)/2 \quad (18)$$

and $F(-ip-ir, -ip; 1-ir; x)$ is a hypergeometric function ($F(a, b; c; x) \equiv F(b, a; c; x)$; $F(a, b; c; 0) = 1$).

Note that solution (17) is consistent with the coupled nature of the original Eqs. (8). This can be easily verified by considering separately both Eqs. (8) at $x = -0$ (use the differentiation formula ¹² for the F functions and remember that $ff^*_{t \rightarrow -\infty} = (1+p/r)/2$).

The asymptotic form for g as $t \rightarrow +\infty$ (i. e. $x \rightarrow -\infty$) can be found by making use of the appropriate linear transformation formula ¹³

for the $F(-ip-ir, -ip; 1-ir; x \rightarrow -\infty)$ function appearing in Eq. (17). Replacing in the resulting linearly-transformed¹³ form of $g_t \rightarrow +\infty$ the $F(a, b; c; 1/x)$ functions by unity and performing a fair amount of Γ -algebra, one finds

$$g_t \rightarrow +\infty = D(-x)^{ip/2-ir/2} \left\{ \Sigma - (-x)^{ir} \sinh(p\pi) / \sinh(r\pi) \right\}, \quad (19)$$

where

$$\Sigma = r \Gamma^2(-ir) \left[(r-p) \Gamma(-ip-ir) \Gamma(ip-ir) \right]^{-1}. \quad (19')$$

Here, the Γ 's are gamma functions.

Finally, we develop the squared modulus $gg^* \rightarrow +\infty$ and use Eq. (18). We get for the time-average value of the transition probability, $(gg^*)_{+av}$,

$$2(gg^*)_{+av} \equiv 1 - (S_{+av})_z = (1+p/r)s_- s_+ + (1-p/r) \left[\sinh(p\pi) / \sinh(r\pi) \right]^2, \quad (20)$$

where

$$s_{\pm} = \sinh \left\{ (r \pm p)\pi \right\} / \sinh(r\pi). \quad (20')$$

In writing Eq. (20) we have dropped an (expected) rapidly oscillating term which enters (its time-average value is zero) since we are interested only in the mean value of the residue polarization.

By some algebraic manipulation, eq. (20) may be written as¹⁴

$$(S_{+av})_z = (p/r) \left\{ 2 \sinh^2(p\pi) / \sinh^2(r\pi) - 1 \right\}. \quad (21)$$

Thus it follows from eq. (5) that $S_z(t \rightarrow +\infty)$ is expressed by eq. (1).

Finally, we note that eq. (1) reduces to the Froissart and Stora formula⁶ in the case where Teng's condition,⁷ i. e.

$$p \gg 2 ,$$

(22)

is satisfied.

REFERENCES.

- (1) - W. Brefeld, V. Burkert, K. P. Crone, W. von Drachenfels, E. Ehse, M. Hofmann, D. Husmann, G. Knop, Th. M. Müller, W. Paul and H. R. Schaefer, 3-rd Intern. Symposium on High-Energy Physics with Polarized Beams and Polarized Targets, ANL, Oct. 25-28, 1978. Proceedings to be published in AIP Conference Proceedings. See also D. Husmann, Rome Meeting of ECFA-LEP Working Group, Rome, Nov. 23-24, 1978. Proceedings to be published.
- (2) - T. Khoe, R. L. Kustom, R. L. Martin, E. F. Parker, C. W. Potts, L. G. Ratner, R. E. Timm, A. D. Krisch, J. B. Roberts and J. R. O'Fallon, Part. Acc. 6, 213 (1975).
- (3) - M. Bell, P. Germain, W. Hardt, W. Kubischta, P. Lefèvre and D. Möhl, High-Energy Physics with Polarized Beams and Targets, Argonne, 1976. Edited by M. L. Marshak (AIP Conference Proceedings No. 35, 1976) p. 405.
- (4) - E. D. Courant, Workshop on Possibilities for Higher Energy Polarized Proton Beams, Ann Arbor, Oct. 18-27, 1977. Edited by A. D. Krisch and A. J. Salthouse (AIP Conference Proceedings No. 42, 1978) p. 94. See also E. D. Courant and L. G. Ratner (Co-Chairmen of the Working Group on Accelerator Problems), ibidem, p. 41; E. D. Courant, 3-rd Polarization Symposium (Ref. 1); L. C. Teng, 3-rd Polarization Symposium (Ref. 1).
- (5) - S. Suwa, 3-rd Polarization Symposium (Ref. 1).
- (6) - M. Froissart and R. Stora, Nuclear Instr. and Meth. 7, 297 (1960).
- (7) - L. C. Teng, Proceedings of the Summer Studies on High-Energy Physics with Polarized Beams, Argonne, July 1974. Edited by J. B. Roberts (ANL/HEP 75-02, 1975), p. XIII-1. Eq. (22).
- (8) - A. Turrin, Proceedings 10th Intern. Conf. on High-Energy Accelerators, Protvino, Moscow, July 1977, Vol. 2, p. 81.
- (9) - G. M. Murphy, Ordinary Differential Equations and Their Solutions (Van Nostrand, 1960), Eq. 518, pag. 370.
- (10) - Handbook of Mathematical Functions, Edited by M. Abramowitz and I. A. Stegun (National Bur. Stand. Appl. Math. Series 55, U. S. G. P. O. , Washington, 1965), Chap. 15.
- (11) - NBS Handbook (Ref. 10), Eq. 15. 5. 3. of Chap. 15.
- (12) - NBS Handbook (Ref. 10), Eq. 15. 2. 1 of Chap. 15.
- (13) - NBS Handbook (Ref. 10), Eq. 15. 3. 7 of Chap. 15.
- (14) - The author is grateful to R. D. Ruth (BNL) for pointing this out.