

Presented at the "1979  
intern. Symp. on Lepton  
and Photon Interactions  
at High Energies",  
Batavia, August 1979.

ISTITUTO NAZIONALE DI FISICA NUCLEARE  
Laboratori Nazionali di Frascati

LNF-79/65(P)  
26 Settembre 1979

M. Spinetti: MULTIHADRON PRODUCTION AT ADONE.

MULTIHADRON PRODUCTION AT ADONE

M. Spinetti  
 INFN, Laboratori Nazionali di Frascati  
 Frascati, Italy

SUMMARY

A short review on recent experimental results obtained at Adone, on multihadronic  $e^+e^-$  annihilation is given.

Results and comments concern:

- 1) R values, compared QCD and EVMD predictions;
- 2) Charged and neutral multiplicities, threshold for "energy crisis", G parities and  $SU_3$  checks;
- 3)  $\sigma(2\pi^+, 2\pi^-)$ : evidence for a large  $\rho(1600)$ ;
- 4)  $\sigma(\pi^+\pi^-2\pi^0)$ : lack of evidence for a  $\rho'(1250)$ ;
- 5)  $\sigma(2\pi^+, 2\pi^-, n\pi^0)$ : possible interpretation of the 1.82 GeV structure.

INTRODUCTION

The results I'll report about are some of the latest results from the Frascati  $e^+e^-$  storage ring Adone. Recently interesting results have been obtained on QED tests, search for a heavy electron,  $\gamma\gamma$  interactions. I'll concentrate only on results on multihadronic  $e^+e^-$  annihilation, mainly coming from the  $\gamma\gamma 2$  experiment. Results on total cross sections and two body reactions will be available also from MEA and  $B\bar{B}$  experiments in the next months.

Adone is an  $e^+e^-$  storage ring covering interval  $1.4 \leq W \leq 3.1$  GeV with a typical luminosity  $L \sim 3 \times 10^{29}$   $\text{cm}^{-2} \text{sec}^{-1}$  at  $W = 3$  GeV, which drops down to  $L \sim 2 \times 10^{28}$  at  $W = 1.5$  GeV.

Three of the four interaction regions of Adone are equipped with experimental set-ups; in the fourth a small angle ( $3^\circ + 6^\circ$ ) Bhabha scattering detector, which serves as a high rate luminosity monitor, is installed. The luminosity used by the experiments is directly measured by counting wide angle electron pairs detected in each apparatus.

The three set-ups are somewhat complementary since MEA has magnetic analysis ( $\Delta\Omega = 40\%$ ),  $\gamma\gamma 2$  is oriented toward  $\gamma$  detection ( $\Delta\Omega_\pi 90\%$ ,  $\Delta\Omega_\gamma 66\%$ ) and  $B\bar{B}$  uses calorimetric information from large liquid scintillators ( $\Delta\Omega = 70\%$ ). They are described in detail in refs. (2, 3, 4) respectively. The three experiments ran at Adone up to June 1978.

NARROW RESONANCES

All data have been collected in scanning mode. As first, combining all the results of the three Adone experiments, we obtain for the  $J/\psi$ -like particle existence, in the full energy range ( $1.42 \leq W \leq 3.1$  GeV) an upper-limit of  $\sim 0.05 \Gamma_{ee}(J/\psi)$  with 90% C.L..

R VALUES

In order to obtain R, multiplicities and exclusive cross sections, using the standard likelihood method, the following assumptions were made:

- a) all particles in the final states are  $\pi$ ;
- b) invariant phase space (IPS) momenta distributions;
- c) isospin relation  $\langle n^\pm \rangle = 2\langle n^0 \rangle$  for  $n^\pm + n^0 = 5$ ;
- d) maximum multiplicity  $n^\pm + n^0 = 6$  up to 2 GeV and  $n^\pm + n^0 = 8$  up to 3.1 GeV.

The first assumption is reasonable especially at lower energies. In a more quantitative way the MEA<sup>5</sup> group has measured the percentage of kaons respect to the total prongs in a given momentum window (see Fig. 1). This K fraction is lower than 20%.




Fig. 1. Ratio between the numbers of K and prongs whose momentum falls in the range  $400 \leq p \leq 600$  MeV/c.

Using a special category of events the  $\gamma\gamma$  group has also estimated an upper limit at 1.5 GeV for  $KK^*$  production of 8 nb (95% c.l.), that is 20% of the total cross section<sup>6</sup>.

The DCI-DM1 experiment<sup>7</sup> gives also the ratio  $\sigma(KK^*)/\sigma_{TOT} \approx 20-30\%$ . In conclusion this assumption leads to a systematic error on R of 10-20% at the lowest energies, that becomes negligible at higher energies.

The rationale for the second assumption can be found in the energy distribution of produced hadrons (Fig. 2) well fitted by a thermodynamical spectrum with




Fig. 2. Invariant distribution for  $\pi$  and K as function of the total energy of the particle.

$KT = 145 \pm 3$  MeV. This distribution can be independently reproduced summing the IPS momentum distributions for each channel weighted with their own exclusive cross sections we will see after.

The values of R for at least  $3\pi$  in the final states are reported in Fig. 3 together with a large panorama of already published values<sup>8-13</sup>. The values of VEPP2M are calculated summing the two most important cross sections that are  $\sigma(2\pi^-, 2\pi^0)$  and  $\sigma(4\pi^\pm)$ . The agreement between all the machines is very good.

of R, calculated by formula  $R = 16\pi^2 / \Delta m_0^2 f_0^2 \approx 2.5^{16}$  fits very well the value of R above 2.4 GeV.

The same contradictory situation comes from QCD corrections<sup>17</sup> to the asymptotic value  $R = 3 \sum_q Q_q^2 = 2$  which gives a  $R \approx 2.3$  between 2.4 and 3.6 GeV.




Fig. 3. R versus total energy. The dotted and dashed lines represent the QCD and EVMD expectations respectively.

The quoted errors are statistical only while the systematical ones on  $\gamma\gamma$  group data are  $\sim 20\%$  (due to luminosity, efficiencies, K presence).

Data from DCI-M3N<sup>14</sup>, not quoted, are in general good agreement with the Adone  $\gamma\gamma$  data.

The energy behaviour of the ratio R is now rather well established above 1 GeV.

In the low energy region after the  $\phi$ , there is a sharp rise whose pattern is typical of the opening of new channels. In this behaviour the more important contribution comes from  $\sigma(4\pi^\pm)$  as we will see later.

Theoretical calculations give good asymptotic values but are not able to reproduce the low energy behaviour. The asymptotic value is expected to be reached from above at the lowest energies where, on the contrary, the measured R is growing. More precisely, the EVMD, with the help of the local duality<sup>15</sup>, predicts, around the asymptotic value, a series of oscillations due to the vector meson recurrences. Using the mass formula  $\Delta m_i^2 \approx 1 \text{ GeV}^2$  and scaling  $\Gamma_{ee}^{-1}$  and  $\Gamma_{tot}$  with the masses the expected values of R do not fit the data concluding that, as we will see also later, some of these recurrences do not exist or have  $\Gamma_{ee}$  much smaller. On the contrary, the asymptotic value

### MULTIPLICITIES, G-PARITIES

Mean charged and neutral multiplicities are reported in Fig. 4. New  $\ln W$  fits are reported also.




Fig. 4. Charged and neutral multiplicity.

Comparing neutral multiplicity with 1/2 of the charged multiplicity clearly the so called "energy crisis" appears starting from  $\sim 2$  GeV. A possible interpretation for such a phenomenon is  $\eta$ 's production.

Before considering the various reactions which contribute to the total cross section, let us divide them in two categories, according to the G-parities. The total cross section for even,  $G^+$ , and odd,  $G^-$ , number of pions are reported in Fig. 5 ( $n \geq 3\pi$ ).  $G^+$  dominant-




Fig. 5.  $G^+$  (sum of  $\sigma(n\pi)$  with  $n$  even) and  $G^-$  (sum of  $\sigma(n\pi)$  with  $n$  odd) versus total energy.

ce is expected. Taking into account the quarks content of the  $\rho$ -like and  $\omega$ -like states, one finds:

$$\frac{|A(e^+e^- \rightarrow \omega\text{-like} \rightarrow G^-)|^2}{|A(e^+e^- \rightarrow \rho\text{-like} \rightarrow G^+)|^2} = \frac{|Q_u + Q_d|^2}{|Q_u - Q_d|^2} = \frac{1}{9} = \frac{\Gamma_e(\omega\text{-like})}{\Gamma_e(\rho\text{-like})} = \frac{\int_{M_\phi}^W G^-(W) dW}{\int_{M_\phi}^W G^+(W) dW}$$

being the  $\Gamma_e$  proportional to the integrated cross section over the resonance. Data have been collected at Adone in scanning mode, so a real energy integration can be done. The result is reported in Fig. 6.




Fig. 6. Ratio between  $G^+$  and  $G^-$  after the energy integration starting from  $W = 1.42$  GeV.

Adone- $\gamma\gamma$  data are in fair agreement with  $SU_3$  prediction, if no dramatic contribution to  $G^-$  is present for  $M_\phi < W < 1.42$  GeV.

### EXCLUSIVE CROSS SECTIONS

$$\sigma(\pi^+\pi^-\pi^+\pi^-)$$

The better established channel is the 4 charged pions. The  $\gamma\gamma$  and MEA values are reported in Fig. 7




Fig. 7.  $\sigma(\pi^+\pi^-\pi^+\pi^-)$ . The dashed line represents the Breit-Wigner best fit.

together with those already published<sup>10, 18, 19, 20</sup>. The general behaviour shows a broad resonance centered at  $\sim 1.5$  GeV interpreted as the  $\rho$ -recurrence  $\rho''(1.6)$ . The dashed line represents a best fit over all the values with only one relativistic Breit-Wigner with an energy dependent width for threshold effect (for sake of simplicity assumed as a simple function of the IPS of  $\rho''(1.6) \rightarrow \rho\pi\pi$  in S wave) according to the following formulas:

$$\sigma = \frac{12\pi}{s} \frac{M_v^2 \Gamma_v \Gamma_{ee}}{(s-M_v^2)^2 + M_v^2 \Gamma_v^2}, \quad \Gamma_v = \Gamma_0 \frac{\alpha K}{1+\alpha K}$$

where  $K = \text{IPS}(\rho\pi\pi)/s$  and  $\alpha$  a free parameter.

The result of the fit gives the following parameters:

$$\begin{aligned} M_v &= 1650 \pm 20 \text{ MeV}, \\ \Gamma_v(M_v) &= 710 \pm 160 \text{ MeV}, \\ \Gamma_{ee} &= 2.8 \pm 0.2 \text{ keV}. \end{aligned}$$

All these parameters are in a rather good agreement with those known of the  $\rho''(1.6)$ . The  $\Gamma_e$  value expected by local duality is  $\Gamma_e \simeq 3.1$ .

Dynamical correlations in the final state have been investigated by the MEA group<sup>10</sup>.

Selecting the 4 prongs events the corresponding Dalitz plot is shown in Fig. 8.




Fig. 8. Scatter plot of the invariant mass of oppositely charged pion pairs,  $M(\pi_1\pi_2)$  vs the invariant mass of the remaining pair  $M(\pi_3\pi_4)$ .

The result largely favours a  $\rho\pi\pi$  intermediate state with the two pions in S-wave.

$$\sigma(\pi^+\pi^-\pi^0\pi^0).$$

Another important channel in the total cross section is that with 2 charged plus 2 neutral pions in the final state. The  $\gamma\gamma$  and MEA values are reported in Fig. 9 together with those from VEPP2-M<sup>9</sup>. This cross section shows a rather different behaviour respect to the  $\sigma(4\pi^\pm)$ .




Fig. 9.  $\sigma(\pi^+\pi^-\pi^0\pi^0)$ . The dashed-dotted line is the calculation of the  $\rho_{TAIL} \rightarrow \omega\pi^0$ . The dashed line is the incoherent sum of the  $\rho_{TAIL} \rightarrow \omega\pi^0$  and the  $\rho''(1.6) \rightarrow \rho^0\pi^0\pi^0$ .

Two contributions must be taken into account. First, the  $\rho$ -tail going into  $\omega\pi$ , computed by Renard<sup>22</sup> starting from the  $\omega \rightarrow \rho\pi$  decay (dashed-dotted line). Second, the  $\rho''(1.6)$  decay via  $\rho^0\pi^0\pi^0$  whose cross section can be taken as  $(\sigma(4\pi^\pm))/2$ , if the two  $\pi$  are in S-wave. An incoherent sum of these two contributions is reported (dashed line) also in Fig. 9.

The measured values are systematically higher

with respect to this curve so that other contributions should be present.

In the low energy region ( $W \lesssim 1.5$  GeV) integrating the cross section exceeding the dashed line of Fig. 9, we get a value  $\Gamma_e \lesssim 0.6$  keV. On the other hand around 1.25 GeV the first  $\rho$ -recurrence is expected. Local duality, that works rather well on  $J/\psi$  and  $Y$  families, gives<sup>15</sup>:

$$\left(\frac{m\Gamma_{ee}}{\Delta m^2}\right)_\rho = \left(\frac{m\Gamma_{ee}}{\Delta m^2}\right)_{\rho'} = \left(\frac{m\Gamma_{ee}}{\Delta m^2}\right)_{\rho''}$$

Applying this formula to the  $\rho''(1.6)$ , we obtain, within a factor 2, the values quoted above.

For the  $\rho'(1.25)$  we obtain, using  $\Delta m^2 \approx 1$  GeV<sup>2</sup>,  $\Gamma_e \approx 4$  keV that is far from the value  $\Gamma_e \lesssim 0.6$  quoted above.

$$\sigma(2\pi^+, 2\pi^-, 1\pi^0) \text{ and } \sigma(2\pi^+, 2\pi^-, 2\pi^0).$$

The separation between these two channels is a very critical one: it depends essentially on how well the photons detection efficiency is known. For the  $\gamma\gamma$  apparatus this efficiency has been checked on the observed photons taking in account also the optical spark chamber efficiency for multisparks. However the sum of these two cross sections does not depend on the photon efficiency and is reported in Fig. 10. The high point at  $W = 1.82$  GeV represents, integrated in energy, the resonance ( $\Gamma \approx 30$  MeV) already published by the 3 Adone experiments<sup>23</sup>.





Fig. 10. Sum of  $\sigma(2\pi^+, 2\pi^-, 2\pi^0)$  and  $\sigma(2\pi^+, 2\pi^-, 1\pi^0)$ .

A nice interpretation of this structure is a decay of a baryonium state near  $N\bar{N}$  threshold according to the following scheme:



The small value  $\Gamma_e \approx 60$  eV favours this interpretation instead of a standard vector meson recurrence.

Unlikely, in separating these two channels the errors become larger (Figg. 11 and 12) so that it is impossible to determine the resonant channel.

The values of  $\sigma(2\pi^+, 2\pi^-, 1\pi^0)$  agree rather well with the DM1 data<sup>21</sup>.




Fig. 11.  $\sigma(2\pi^+, 2\pi^-, 1\pi^0)$ .




Fig. 12.  $\sigma(2\pi^+, 2\pi^-, 2\pi^0)$ .

$\sigma(\pi^+\pi^-\pi^0)$ .

For completeness in Fig. 13 the  $\sigma(\pi^+\pi^-\pi^0)$  is reported.




Fig. 13.  $\sigma(\pi^+\pi^-\pi^0)$ .

$\gamma\gamma 2$  Group (Frascati-Bologna-Rome Collaboration):  
C. Bacci, R. Baldini-Celio, R. Battistoni, D. Bollini, G. Capon, R. Del Fabbro, G. De Zorzi, E. Iarocci, M. Massai, S. Moriggi, G. P. Murtagh, G. Pensò, M. Spinetti, L. Trasatti and B. Stella.

MEA Group (Frascati-Maryland-Naples-Padua-Rome Collaboration):

- R. Bernabei, S. D'Angelo, B. Esposito, A. Marini, P. Monacelli, N. Moricca, M. Nigro, L. Paoluzi, L. Pescara, G. Piano Mortari, F. Ronga, R. Santonico, A. Sebastiani, B. Sechi-Zorn and G. Zorn.
2. C. Bacci et al., Phys. Letters 64B, 356 (1976).
  3. W. W. Ash et al., Frascati report LNF-77/18 (1977); Lett. Nuovo Cimento 11, 705 (1974).
  4. G. Barbiellini et al., Phys. Letters 68B, 397 (1977).
  5. B. Esposito et al., Accepted by Nuovo Cimento.
  6. C. Bacci et al., Frascati report LNF-79/34 (1979); Phys. Letters 86B, 234 (1979).
  7. B. Delcourt, Talk in these Proceedings.
  8. G. Cosme et al., Phys. Letters 63B 349 (1976); G. Parroul et al., Phys. Letters 63B, 357 (1976).
  9. V. A. Sidorov, Talk in these Proceedings.
  10. B. Esposito et al., Lett. Nuovo Cimento 19, 21 (1977); 25, 5 (1979).
  11. J. Siegrist et al., to be submitted to Phys. Rev.
  12. S. Yamada, Proc. of Intern. Symp. on Lepton and Photon Interactions at High Energies, Hamburg (1977).
  13. G. Knies, *ibid.*
  14. G. Cosme et al., Orsay report LAL 78/32 (1978).
  15. M. Greco, Phys. Letters 77B, 84 (1978).
  16. M. Greco, Nuclear Phys. 63B, 398 (1978).
  17. T. Appelquist et al., Phys. Rev. 12, 43 (1975).
  18. A. Cordier et al., Orsay report LAL 78/31 (1978).
  19. R. F. Schwitters, Proc. of 1975 Intern. Symp. on Lepton and Photon Interaction, Stanford (1975).
  20. V. A. Sidorov, Proc. of the XVIII Intern. Conf. on High Energy Physics, Tbilisi (1976).
  21. B. Delcourt, Talk in these Proceedings.
  22. F. M. Renard, Nuovo Cimento 64A, 979 (1969).
  23. B. Esposito et al., Phys. Letters 68B, 389 (1978); C. Bacci et al., Phys. Letters 68B, 393 (1978); G. Barbiellini et al., Phys. Letters 68B, 397 (1978).

REFERENCES

1. List of authors:

$B\bar{B}$  Group (Frascati-Naples-Pisa-Rome Collaboration):

M. Ambrosio, C. Bemporad, R. Biancastelli, G. Bro sco, M. Calvetti, L. Cerrito, F. Costantini, G. Gian nini, P. Lariccia, G. Paternoster, L. Tortora and U. Troya.