

ISTITUTO NAZIONALE DI FISICA NUCLEARE
Laboratori Nazionali di Frascati

LNf-81/81

M. Basile et al. :
THE METHOD OF REMOVING THE LEADING PROTONS
IN THE STUDY OF HIGH-ENERGY pp REACTIONS,
COMPARED WITH THE STANDARD ANALYSIS

Estratto da :

Nuovo Cimento 65A, 414 (1981)

The Method of Removing the Leading Protons in the Study of High-Energy pp Reactions, Compared with the Standard Analysis.

M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ,
P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, R. NANIA,
F. PALMONARI, V. ROSSI, G. SARTORELLI, M. SPINETTI, G. SUSINNO,
G. VALENTI, L. VOTANO and A. ZICHICHI

CERN - Geneva, Switzerland

Istituto di Fisica dell'Università - Bologna, Italia

Istituto Nazionale di Fisica Nucleare - Laboratori Nazionali di Frascati, Italia

Istituto Nazionale di Fisica Nucleare - Sezione di Bologna, Italia

Istituto di Fisica dell'Università - Perugia, Italia

Istituto di Fisica dell'Università - Roma, Italia

(ricevuto il 14 Agosto 1981)

Summary. — The method of removing the leading protons in order to study the properties of multiparticle systems produced in pp interactions is compared, in a direct way, with the standard method. The experimental results show that the remarkable agreement between pp interactions and e^+e^- annihilations, found by removing the leading protons, is lost if the standard analysis is adopted. This shows the validity and the relevance of our method of subtracting the leading-proton effects in order to understand the properties of multiparticle systems produced in strong interactions. The experiment has been performed using two values of the pp c.m. energy, $(\sqrt{s})_{pp} = 30$ and 62 GeV.

1. — Introduction.

The method ⁽¹⁾ of removing the leading-proton effects, in the study of the multiparticle systems produced in pp interactions, has provided a series of

⁽¹⁾ M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ, P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, F. PALMONARI, G. SARTORELLI, G. VALENTI and A. ZICHICHI: *Phys. Lett. B*, **92**, 367 (1980).

remarkable similarities⁽¹⁻⁷⁾ with the simple and well-understood multiparticle production mechanism of e^+e^- annihilation.

The purpose of the present paper is to present a straightforward comparison of our new method⁽¹⁾ with the standard one. The latter method is based on the notion that the energy in the pp c.m. system, *i.e.* \sqrt{s} , is the basic parameter which characterizes the properties of our event. The new method⁽¹⁾ is based on the notion that the basic parameter in a pp interaction is the effective hadronic energy available for particle production. This is obtained taking into account the quadrimomentum carried by the leading protons.

2. - The experimental apparatus and data taking.

The experiment was performed at the CERN Intersecting Storage Rings (ISR) using pp interactions at two c.m. energies, $(\sqrt{s})_{pp} = 30$ GeV and $(\sqrt{s})_{pp} = 62$ GeV. The apparatus consisted mainly of a large-volume magnetic field, the so-called Split-Field Magnet (SFM), coupled to a powerful system of multiwire proportional chambers (MWPCs). A description of the apparatus has been given elsewhere⁽⁸⁾. The set-up was used in the simplest possible mode, *i.e.* the « minimum » bias trigger⁽¹⁻⁷⁾. For details on data taking we refer the reader to our previous papers⁽¹⁻⁷⁾.

⁽²⁾ M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ, P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, R. NANIA, F. PALMONARI, G. SARTORELLI, G. VALENTI and A. ZICHICHI: *Phys. Lett. B*, **95**, 311 (1980).

⁽³⁾ M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ, P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, R. NANIA, F. PALMONARI, G. SARTORELLI, G. VALENTI and A. ZICHICHI: *Nuovo Cimento A*, **58**, 193 (1980).

⁽⁴⁾ M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ, P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, R. NANIA, F. PALMONARI, G. SARTORELLI, G. VALENTI and A. ZICHICHI: *Lett. Nuovo Cimento*, **29**, 491 (1980).

⁽⁵⁾ M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ, P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, R. NANIA, F. PALMONARI, G. SARTORELLI, M. SPINETTI, G. SUSINNO, G. VALENTI and A. ZICHICHI: *Phys. Lett. B*, **99**, 247 (1981).

⁽⁶⁾ M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ, P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, R. NANIA, F. PALMONARI, G. SARTORELLI, M. SPINETTI, G. SUSINNO, G. VALENTI, L. VOTANO and A. ZICHICHI: *Lett. Nuovo Cimento*, **30**, 389 (1981).

⁽⁷⁾ M. BASILE, G. CARA ROMEO, L. CIFARELLI, A. CONTIN, G. D'ALÍ, P. DI CESARE, B. ESPOSITO, P. GIUSTI, T. MASSAM, R. NANIA, F. PALMONARI, G. SARTORELLI, M. SPINETTI, G. SUSINNO, G. VALENTI, L. VOTANO and A. ZICHICHI: *Lett. Nuovo Cimento*, **31**, 273 (1981).

⁽⁸⁾ R. BOUCLIER, R. C. A. BROWN, E. CHESI, L. DUMPS, H. G. FISCHER, P. G. INNOCENTI, G. MAURIN, A. MINTEN, L. NAUMANN, F. PIUZ and O. ULLALAND: *Nucl. Instrum. Methods*, **125**, 19 (1975).

3. - Data analysis and results.

Two sets of data have been taken at two c.m. energies: $(\sqrt{s})_{pp} = 30$ GeV and $(\sqrt{s})_{pp} = 62$ GeV.

3'1. Data at $(\sqrt{s})_{pp} = 30$ GeV. - The number of « minimum bias » events at $(\sqrt{s})_{pp} = 30$ GeV was 6473. These events have been analysed in the standard way, *i.e.* the fractional momentum of a particle, produced in a pp interaction, has been expressed by the standard variable

$$(1) \quad x_R = \frac{2|\mathbf{p}|}{\sqrt{s}},$$

where $|\mathbf{p}|$ is the modulus of the momentum of that particle.

In order to avoid any contamination from the unidentified protons, the inclusive distribution has been done using only negative particles. This is in fact the safest way to avoid a leakage of protons from the initial to the final state.

The inclusive distribution for negative particles is

$$(2) \quad \frac{1}{N_{ev}} \frac{dN(-)}{dx_R},$$

where N_{ev} is the number of events and $dN(-)$ is the number of negative particles in the interval dx_R . The data are reported in fig. 1. This distribution has been compared with one half of the distribution for charged particles produced in e^+e^- annihilation, *i.e.*

$$(3) \quad \frac{1}{2} \frac{1}{\sigma} \frac{d\sigma(e^+e^- \rightarrow h^\pm + \text{anything})}{dx_R}.$$

The factor $\frac{1}{2}$ is there because only one sign of charge is considered. The e^+e^- data must, of course, be taken at the same c.m. energy, *i.e.* $(\sqrt{s})_{e^+e^-} = (27.4 \div 31.6)$ GeV. The e^+e^- data ⁽⁹⁾, reported in terms of

$$(s)_{e^+e^-} \frac{d\sigma}{dx_R},$$

have been transformed into

$$\frac{1}{\sigma} \frac{d\sigma}{dx_R}$$

using for the ratio R the value ⁽⁹⁾

$$R = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = 4.$$

⁽⁹⁾ R. H. WILK: DESY Report 80/124 (1980); R. BRANDELIK *et al.* (TASSO COLLABORATION): *Phys. Lett. B*, **89**, 418 (1980).




Fig. 1. - The inclusive fractional-momentum distributions of negative particles, for pp interactions by using the standard analysis at $(\sqrt{s})_{pp} = 30$ GeV (\circ), by using the method of removing the leading protons at $(\sqrt{s})_{pp} = 62$ GeV and $2E_{had} = (28 \div 32)$ GeV (\bullet); the e^+e^- data at $(\sqrt{s})_{e^+e^-} = (27.4 \div 31.6)$ GeV (\blacktriangle).

3'2. Data at $(\sqrt{s})_{pp} = 62$ GeV. - The number of « minimum bias » events at $(\sqrt{s})_{pp} = 62$ GeV was 62270. The analysis of these data has been done

using the method of removing the leading protons. The method has already been described in previous publications (¹⁻⁷), where also the details about the overall procedure concerning event selection, track selection and corrections for detection efficiencies can be found.

The basic quantity, in this method of analysis, is the effective hadronic energy available for multiparticle production, E_{had} . If we call E_{leading} the energy of the leading proton in one hemisphere of the pp event, the quantity E_{had} is given by

$$(4) \quad E_{\text{had}} = \frac{(\sqrt{s})_{\text{pp}}}{2} - E_{\text{leading}}.$$

The fractional-momentum variable is now

$$(5) \quad x_R^* = \frac{|\mathbf{P}|}{E_{\text{had}}}.$$

The inclusive distribution for negative particles

$$(6) \quad \frac{2}{N_{\text{ev}}} \frac{dN(-)}{dx_R^*}$$

is determined in the energy range $2E_{\text{had}} = (28 \div 32)$ GeV. This is the value required in order to compare pp and e^+e^- data at equivalent energies. In this effective hadronic energy range the number of pp events is 1471. The factor two in (6) arises because the analysis is made in the hemisphere where we can identify the « leading » proton, and it is a trivial normalization factor to the full event.

The results (see fig. 1) are compared with the same distribution

$$(7) \quad \frac{1}{2} \frac{1}{\sigma} \frac{d\sigma}{dx_R},$$

from e^+e^- at $(\sqrt{s})_{e^+e^-} = (27.4 \div 31.6)$ GeV, already used to compare the $(\sqrt{s})_{\text{pp}} = 30$ GeV pp data.

4. - Conclusions.

The results of the inclusive fractional-momentum distribution, obtained using the standard method and our new method of analysis, are reported in fig. 1 together with the e^+e^- data.

The standard method of analysis, based on the c.m. energy in the pp system, $(\sqrt{s})_{\text{pp}}$, produces an inclusive fractional-momentum distribution which clearly differs from the e^+e^- data.

The new method of analysis, based on the effective hadronic energy available for multiparticle production, E_{had} , shows a remarkable agreement with e^+e^- data, at $(\sqrt{s})_{e^+e^-} = 2E_{\text{had}}$. This is a direct proof of the relevance and validity of the method of subtracting the leading-proton effects in pp interactions.

● RIASSUNTO

Il metodo, da noi introdotto, di rimuovere i protoni «leading» nello studio delle proprietà dei sistemi di particelle prodotti nelle interazioni pp è confrontato, in modo diretto, con l'analisi ordinaria. I risultati sperimentali mostrano che il notevole accordo, tra interazioni pp e annichilazioni e^+e^- , che si ottiene rimuovendo i protoni «leading», non si ha più quando si esegue l'analisi ordinaria. Questo risultato mostra la validità e l'importanza del nostro metodo per capire le proprietà dei sistemi di particelle prodotti nelle interazioni forti. L'esperimento è stato eseguito usando due valori dell'energia nel sistema c.m. pp, $(\sqrt{s})_{pp} = 30$ e 62 GeV.

Сравнение метода удаления лидирующих протонов при исследовании pp реакций при высоких энергиях со стандартным анализом.

Резюме (*). — Метод удаления лидирующих протонов при исследовании свойств многочастичных систем, образованных в pp взаимодействиях, непосредственно сравнивается со стандартным методом. Экспериментальные результаты показывают, что удивительное согласие между pp взаимодействиями и e^+e^- аннигиляцией, обнаруженное с помощью метода удаления лидирующих протонов, исчезает, если используется стандартный метод. Этот факт указывает на справедливость и уместность предложенного нами метода, извлечения эффектов лидирующих протонов для интерпретации свойств многочастичных систем, образованных в сильных взаимодействиях. Проведен эксперимент с использованием двух значений pp энергии в системе центра масс, $(\sqrt{s})_{pp} = 30$ и 62 ГэВ.

(*). *Переведено редакцией.*