

ISTITUTO NAZIONALE DI FISICA NUCLEARE

Laboratori Nazionali di Frascati

Submitted to
Phys. Rev. C

LNF-84/65(P)
30 Ottobre 1984

D. Babusci, R. Bernabei, O.M. Bilaniuk, L. Casano, S. d'Angelo, M.P.
De Pascale, S. Frullani, G. Giordano, B. Girolami, G. Matone, M. Mattioli,
P. Picozza, D. Prospero and C. Schärfer:
POLARIZATION ASYMMETRY IN THE PHOTODISINTEGRATION OF
THE DEUTERON

POLARIZATION ASYMMETRY IN THE PHOTODISINTEGRATION OF THE DEUTERON

M.P. De Pascale, G. Giordano, G. Matone,
INFN - Laboratori Nazionali di Frascati, Frascati, Italy

D. Babuści, R. Bernabei, O.M. Bilaniuk, L. Casano, S. d'Angelo, M. Mattioli, P. Picozza, D. Prospero
and C. Schaerf
Dipartimenti di Fisica, Università di Roma and INFN - Sezione di Roma - Roma, Italy

S. Frullani and B. Girolami
Istituto Superiore di Sanità, Roma, and INFN - Sezione Sanità - Roma, Italy

ABSTRACT

The reaction $d(\gamma, p)n$ has been studied using a monochromatic and polarized gamma ray beam at energies $E_\gamma = 19.8, 29.0, 38.6$ and 60.8 MeV.

The beam of an intensity $\sim 4 \times 10^5$ γ /sec was obtained by Compton back-scattering of mode-locked laser light off electron bunches in the Adone storage ring. Photoneutron yields were measured at nine neutron angles $\theta_n = 15, 30, 45, 60, 90, 120, 135, 150$ and 165 degrees in the center of mass (CM) for $E_\gamma = 19.8, 29.0$ and 38.6 MeV, and at $\theta_n = 30, 60, 90, 120$ and 150 deg. CM for $E_\gamma = 60.8$ MeV. The polarization independent component $I_0(\theta)$ of the differential cross section and the polarization dependent component $PI_1(\theta)$ were deduced and the angular distribution of the azimuthal asymmetry factor $\Sigma(\theta) = I_1(\theta)/I_0(\theta)$ was obtained. An extensive comparison with theory has been carried out and the inclusion of corrections due to meson exchange currents (MEC) and to Δ -isobar configurations (IC) have been shown to be mandatory at energies $E_\gamma \gg 40$ MeV.

Theoretical fits argue in favor of potentials, such as the Reid soft-core or Paris, that provide for a high D-state percentage in the deuteron wave functions. Theoretical and experimental implications of intermediate energy deuteron photodisintegration studies are discussed in some detail.

1. - INTRODUCTION

The deuteron photodisintegration $d(\gamma, p)n$ is a basic nuclear process and as such it has been extensively used for studies of electromagnetic properties of nucleons and of the nucleon-nucleon (N-N) interactions. As of recent, the development of advanced computational capabilities and of new experimental techniques has thrust the $d(\gamma, p)n$ reaction into the forefront of renewed experimental and theoretical interest.

In particular, it has become evident that non-nucleonic degrees of freedom play an essential role in the dynamics of the deuteron break-up at intermediate energies. Because of their relatively long range, the π -meson exchange currents (π -MEC) are of prime importance, but as the γ -ray energy increases, the sub-nucleonic degrees of freedom due to internal excitation of the isobar configurations (IC) are expected to come into play.

A number of refined calculations of the deuteron photodisintegration cross section below the pion production threshold have become available in the last few years¹⁻²⁵. Among other things, these calculations enlighten the importance of mesonic degrees of freedom within the deuteron at gamma energies as low as 30 or 40 MeV. The experimental implications of these calculations point to a need for new cross section measurements using monochromatic and polarized gamma beams in the energy region below the pion threshold, a range for which so far very few data are available. Particularly important are the measurements of the polarization independent part of the differential cross section $I_0(\theta)$, of the polarization dependent part $PI_1(\theta)$ and of the angular distribution of the azimuthal asymmetry factor $\Sigma(\theta) = I_1(\theta)/I_0(\theta)$ (see Sect. 3 below) which turns out to be remarkably sensitive to meson exchange currents (MEC) and to virtual excitation of nucleonic Δ -isobar configurations (IC).

The present work constitutes the first stage of a more extensive research program that addresses this need. The source of γ -rays, developed at Frascati and named LADON^{26,27}, is laser light Compton back-scattered off ultrarelativistic electrons circulating in the Adone storage ring.

Besides the LADON project, there have been to date three major experimental efforts that resulted in Compton back-scattered photon beams (Lebedev Institute²⁸, Cambridge²⁹, Stanford³⁰). Because in all three cases an external laser beam has been used, the resulting gamma beam intensity was very low and only the Stanford beam had been used in an actual experiment³⁰. On the other hand, in the LADON process the Compton scattering takes place within an optical cavity that is superposed onto a straight section of the storage ring and where the interaction takes place between compact bunches of electrons and phase-locked travelling pulses of concentrated laser power (see Sect. 2 below). This "photon bunching" and the use of a storage ring are the two most important aspects of the LADON approach (Fig. 1). After some stages of development, the LADON facility can now provide gamma ray beams of an intensity of $(3 \text{ to } 5) \times 10^5$ photons/sec. This has permitted the accumulation of a considerable amount of new data on deuteron photodisintegration, particularly on the azimuthal asymmetry.

Reported here are measurements of photo-neutron yields, carried out at nine CM neutron angles ($\theta_n = 15, 30, 45, 60, 90, 120, 135, 150$ and 165 deg.), for three gamma energies ($E_\gamma = 19.8, 29.0$ and 38.6 MeV) and at $\theta_n = 30, 60, 90, 120$ and 150 deg. for $E_\gamma = 60.8$ MeV. Each set of measurements consisted of a series of runs, with the angle φ between the reaction plane and the gamma polarization plane alternating repeatedly between $\varphi = 0^\circ$ and $\varphi = 90^\circ$ so as to insure identical experimental conditions for both polarizations. From these measurements the angular distributions of both the polarization independent component $I_0(\theta)$ and the polarization dependent component $PI_1(\theta)$ of the




FIG. 1 - Schematic representation of the LADON process (not to scale). The effective circumference (i.e. the perimeter consisting of circular and straight sections) is $2\pi R_{\text{eff}}=105.3$ m, so that a bunch of ultrarelativistic electrons takes about 351 ns to make the round. Under normal operation there are three such bunches circulating in the ring, so that they succeed one another every 117 nsec. This time interval defines the length $L=17.55$ m of the optical cavity in which a laser power packet bounces back and forth between the cavity end mirrors. Compton back scattering of the laser photons off the electrons takes place in a well defined portion of a straight segment of the storage ring free from magnetic field (Refs. 26,27,32).

photodisintegration cross section have been deduced and the angular distribution of the asymmetry factor $\Sigma(\theta)$ obtained.

The experimental set-up and our data acquisition system are described in Sect. 2. In Sect. 3 a brief review of the pertinent theory is presented and its experimental relevance discussed. Sect. 4 contains a summary of our experimental results, a critical analysis of these results with respect to current theories and a comparison with earlier experiments. Finally, our main conclusions are summarized in Sect. 5.

2. - EXPERIMENTAL SET-UP

There are essentially two independent experimental set-ups involved in this work: the LADON gamma source on the Adone ring and the actual detection system in an experimental area 42 meters away. The LADON facility will be shortly described first.

2.1. - The Gamma Beam

The basic technical features of the LADON system are shown schematically in Fig. 1 (see Refs. 26 and 27 for further details).

Under normal laser operation, the harmonic modes - resonating at frequencies $f_n=nc/2L$ (n integer) in an optical cavity of length L that comprises the laser tube - have random phases and the

laser power distribution is uniform over the entire cavity. Because the electron-photon scattering takes place only during the passage of an electron bunch through the 5.7 m long Adone straight section (see Fig. 1), this uniformly distributed laser power remains unused for about 100 ns of each 117 ns. The effective laser power can be increased by "compacting" the uniformly distributed laser power into a travelling pulse of phase-locked harmonics³¹. If the length of the cavity is so adjusted that this laser power packet finds itself going to the right just as an electron bunch crosses it going in the opposite direction (see Fig. 1), the yield of back-scattered photon will increase²⁷.

Moreover in this way we define the laser-electron interaction region in the center of the straight section, where there are no quadrupoles or others ring optical components. In this magnetic field free region, the angular divergence of the stored electron beam is minimum, yielding the best monochromaticity for the scattered gamma ray beam.

The cavity mode locking is effected using the piezoelectric property of a quartz crystal for acousto-optic modulation of cavity losses at a frequency $f=c/2L$, which is equal to the difference in frequency, $f_{n+1}-f_n$, between any two neighboring harmonics³². While the losses associated with the mode-locking process lower the average internal power of the laser cavity by about a factor of two (from 60 watts to about 35 watts in the Coherent Radiation Model CR-18 Argon-Ion laser we use), the effective laser power overlapping the electron bunch in the electron-photon interaction region is increased by about a factor of six. It is this spatial and temporal superposition of the electron bunches and the "compacted" laser power which has raised the LADON beam intensity to the level of $\approx 10^5$ photons/sec and which thus made possible to take advantage of its unique properties of tunability and polarization for photonuclear research, particularly for polarization asymmetry measurements.

Other salient features of the LADON gamma source are:

- a) its ready tunability in the energy range from $E_\gamma=5$ to 78.7 MeV, effected by varying the energy of the electrons circulating in the storage ring from 370 MeV to 1.5 GeV;
- b) its nearly perfect polarization (as an example, at $E_\gamma=15$ MeV the degree of linear polarization is $P=0.997$ theoretically and $P=0.99\pm 0.02$ as actually measured²⁶), and a straightforward way to change the plane of polarization to any desired angle relative to the reaction plane;
- c) a very low background of unwanted photons, due essentially only to electron bremsstrahlung off the residual gas molecules in the 10^{-9} Torr vacuum of the Adone ring.

The energy resolution varies from $\sim 2\%$ FWHM at lower gamma energies, increasing up to 8% at $E_\gamma=70$ MeV. The reasons for the decrease of resolution with increasing energy are twofold: the relativistic squeezing of a wider energy range of scattered photons into the forward solid angle of $\Omega=2.5 \times 10^{-8}$ sr defined by the beam collimation system, and the increasing divergence of the electron beam. These two effects set a practical upper limit to the "monochromaticity" of gamma beams produced by Compton back-scattering. In Table

TABLE I - Intensity (in units of 10^5 gammas/sec), resolution (%) and linear polarization degree P of the LADON beam at the γ energies of our experiment.

E_γ (MeV)	γ /sec (10^5)	$\Delta E/E$ (%)	P
19.8	1.0	3.00	0.999
29.0	1.1	5.00	0.998
38.6	1.3	5.74	0.997
60.8	1.8	7.55	0.994

I the γ beam characteristics at the energies of our experiment are reported.

2.2. - Experiment

The lay-out of the experimental area is shown in Fig. 2. After leaving the interaction region, the gamma beam passes a system of collimators which limits it to the forward solid angle of 2.5×10^{-8} sr, and a set of three sweeping magnets that clean the beam of charged particles. During the entire experiment, the beam energy spectrum is analyzed by a magnetic pair spectrometer, the output of which is evaluated on-line by a microprocessor. The target consists of a C_6D_6 deuterated liquid scintillator NE-230 in a cylindrical container 3.81 cm in diameter and 10.16 cm in height, which serves at the same time as proton detector, optically coupled to a 56-AVP photomultiplier.




FIG. 2 - Layout of the experimental area. The laser beam is aligned to $1 \mu\text{m}$ by on-line remote control via CAMAC using a PDP 11/04 minicomputer and position transducers. Three sweeping magnets clear the gamma beam of all charged particles. The principal collimator restricts the beam to a solid angle of 2.5×10^{-8} sr. The energy profile of the gamma beam is continuously monitored by a magnetic pair spectrometer. A cylinder of deuterated scintillation liquid NE-230 serves both as target and proton detector. Neutron time of flight spectra are recorded at five angles using NE-213 scintillators and pulse-shape analysis.

Five neutron detectors, independently movable in the reaction plane, cover an angular range from ~ 15 to ~ 165 deg. Each neutron detector consists of a cylindrical container 30 cm in diameter and 15 cm deep filled with a NE-213 liquid scintillator (C_6H_6) and each coupled to a 58-DVP photomultiplier. The beam intensity is continuously monitored by a lead glass counter which also serves as gamma beam stopper.

The following data are collected via Camac and recorded by a PDP 11/34 for each event:

- The "head" integral H of each pulse from the proton counter (obtained by integrating the entire pulse) and the corresponding "tail" integral T (obtained by starting a second integration 25 ns after the pulse start). These data, processed subsequently off line, permit to suppress most of the considerable gamma background of the proton detectors by effective pulse-shape discrimination.
- Similar "head" and "tail" integrals are recorded for each pulse from any of the five neutron counters, together with the corresponding time of flight and the code number of the particular counter. The time of flight records constitute the essential part of our experimental data.

The information from the proton counter and from the neutron counters is stored only if two coincidence conditions are satisfied:

- A fast coincidence (~ 20 ns window) between the electron beam pulse and a signal from the proton counter.
- A slower coincidence (~ 80 ns window) between the output of the fast coincidence and any of the neutron counters.

The most important contribution to the background are due to electromagnetic processes (multiple Compton scattering, pair-production and annihilation) involving atomic electrons. These processes give mainly rise to narrow peaks in the time of flight spectra of the neutron counters. By using a TPHC-SCA (time to pulse height conversion single channel analyzer) to clear the Camac modules whenever the time of flight is in the "gamma-like" range, the sharp peaks are suppressed.

The first step in the off-line data processing involves reconstruction of the "raw" neutron time of flight spectra for each of the five neutron detectors. For each energy and each neutron angle there are two sets of such spectra, one for $\varphi=0^\circ$ and one for $\varphi=90^\circ$, each obtained by adding the results of a number of corresponding runs. Each "raw" time of flight spectrum includes a residual gamma background, having a flat time structure which is effectively suppressed by pulse shape discrimination. Still further background reduction is effected by imposing a pulse height window onto the signals from the proton ADC (analog to digital converter) and by rejecting signals below a fixed threshold from the neutron ADC. A typical result is shown in Fig. 3. A residual background remains, mostly due to the extreme tails of the gamma peak, the carbon in the target, bremsstrahlung events and cosmic rays.




FIG. 3 - A typical time of flight spectrum at $E_\gamma = 19.8$ MeV (0.8 nsec per channel), obtained after background reduction by pulse shape discrimination and rejection of events that fall outside of the kinematic limits for the proton energy. The neutron peak n is clearly shown; the window indicated by the γ symbol corresponds to the region of the γ background, cut-out by a hardware system.

3. - THEORETICAL CONSIDERATIONS

The general expression of the differential cross section for the $d(\gamma,p)n$ process is given in perturbation theory by:

$$d\sigma/d\Omega = (E_\gamma M/8\pi^2) \left| \langle \psi_f | H_{int} | \psi_i \rangle \right|^2, \quad (1)$$

where ψ_i is the non-relativistic deuteron wave function and ψ_f represents the n-p final state with its interaction. For linearly polarized photons, the above expression is usually reduced¹¹, in the center of mass (CM) system, to the form:

$$\begin{aligned} d\sigma/d\Omega &= I_0(\theta) + P I_1(\theta) \cos 2\varphi = \\ &= I_0(\theta) \left[1 + P \Sigma(\theta) \cos 2\varphi \right], \end{aligned} \quad (2)$$

where φ is the angle between the $d(\gamma,p)n$ reaction and polarization planes and θ is the CM angle between the proton and photon momenta (if neutrons rather than protons are detected, $\theta = \theta_p$ must be replaced by $\theta_n = \pi - \theta$, but otherwise the expression remain unchanged); P represents the degree of linear polarization of the photon beam and

$$\Sigma(\theta) = I_1(\theta)/I_0(\theta) \quad (3)$$

is a parameter which describes the azimuthal asymmetry in the differential cross section due to the polarization of the incident beam. $\Sigma(\theta)$ has been shown¹⁰ to be markedly affected by mesonic degrees of freedom. It should be noted that $I_1(\theta)$ and $\Sigma(\theta)$ are accessible to experimental determination only with polarized γ beams.

The quantities $I_0(\theta)$ and $I_1(\theta)$ of Eq. (2) may be expanded in terms of Legendre polynomials $P_J(\cos \theta)$ and associated Legendre functions $P_J^m(\cos \theta)$ ¹¹, respectively. If one includes multipole contributions up to the dipole-octupole interference, one has:

$$I_0(\theta) = \sum_i^4 A_i(E_\gamma) P_i(\cos \theta), \quad (4a)$$

$$I_1(\theta) = \sum_i^4 B_i(E_\gamma) P_i^2(\cos \theta), \quad (4b)$$

where $A_i(E_\gamma)$ and $B_i(E_\gamma)$ are energy dependent coefficients.

An alternate way in which the function $I_0(\theta)$ and $I_1(\theta)$ may be written^{11,49} is:

$$I_0(\theta) = a + b \sin^2 \theta + c \cos \theta + d \sin^2 \theta \cos \theta + e \sin^4 \theta, \quad (5a)$$

$$I_1(\theta) = f \sin^2 \theta + g \sin^2 \theta \cos \theta + h \sin^4 \theta, \quad (5b)$$

where the energy dependent coefficients are now a, b, c, ..., h. The relationship between these and the previous coefficients is given by¹¹:

$$\begin{aligned}
 a &= A_0 + A_2 + A_4, & b &= (3A_2 + 10A_4)/2, \\
 c &= A_1 + A_3, & d &= -5A_3/2, \\
 e &= 35A_4/8, & f &= 3(B_2 + 15B_4), \\
 g &= 15B_3, & h &= -105B_4/2.
 \end{aligned} \tag{6}$$

While the two forms of expressions for $I_0(\theta)$ and $I_1(\theta)$ are equivalent, the coefficients A_i and B_i of Eqs. (4) are better suited for comparison between theory and experiment because of the orthogonality of the Legendre functions³⁴. As far as the coefficients a, b, c, \dots, h of Eqs. (5) are concerned, the experimental data are in most cases insufficient for their unequivocal determination. Still, the expressions of Eqs. (5) can be useful for extracting additional information regarding nuclear potentials, as will be seen in Sect. 4 below.

In the present experiment, angular distributions in θ_n of neutrons were taken and measurements were made for two polarization angles, $\varphi=0$ and $\varphi=\pi/2$. Under these conditions, Eq. (2) reduces to

$$d\sigma/d\Omega = I_0(\theta_n) + P I_1(\theta_n), \tag{7}$$

for $\varphi=0$, and to

$$d\sigma/d\Omega = I_0(\theta_n) - P I_1(\theta_n) \tag{8}$$

for $\varphi = \pi/2$, respectively.

Thus the values of $I_0(\theta_n)$ and $I_1(\theta_n)$ may be obtained from the experimental data by the following relations:

$$I_0(\theta_n) = \frac{1}{2} \alpha \left[K_1 Y(\theta_n, \varphi=0) + K_2 Y(\theta_n, \varphi=\pi/2) \right], \tag{9}$$

$$I_1(\theta_n) = \frac{1}{2} \alpha \left[K_1 Y(\theta_n, \varphi=0) - K_2 Y(\theta_n, \varphi=\pi/2) \right], \tag{10}$$

where $Y(\theta_n, \varphi)$ is the experimental neutron yield (ratio between the number of photoneutrons detected at θ_n and the gamma-ray flux). The corrective coefficients K_1 and K_2 , evaluated by Monte Carlo methods, account for neutron absorption, multiple scattering and for finite solid angle. In the present work the α constant has been fixed to normalize our data in such a way that $\int I_0(\theta_n) d\Omega_n = 1$; consequently, both $I_0(\theta_n)$ and $I_1(\theta_n)$ are given in units of the total cross section for unpolarized photons ($\sigma_t = \int I_0(\theta_n) d\Omega_n$).

By defining

$$R(\theta_n) = Y(\theta_n, \pi/2) / Y(\theta_n, 0) \tag{11}$$

and writing $K=K_2/K_1$, we obtain a form of Eq. (3) which is directly linked with the neutron yields obtained in the experiment:

$$\Sigma(\theta_n) = \frac{1 - KR(\theta_n)}{P [1 + KR(\theta_n)]}. \tag{12}$$

As previously pointed-out, all available calculations clearly indicate that, to account accurately for the experimental data at photon energies above some 30 or 40 MeV, meson exchange currents (MEC) and nucleon isobar contributions (IC) must be taken into account in the evaluation of the A_i

and B_1 coefficients. The functions $I_1(\theta)$ and $\Sigma(\theta)$ are particularly sensitive to the MEC and IC corrections. For this reason, the $d(\gamma, p)n$ reaction with polarized gammas turns out to be a sensitive probe for checking the standard perturbative calculations of the MEC and IC effects.

Another important aspect of $d(\gamma, p)n$ studies involves the forward and backward differential cross section for unpolarized photons, respectively given by $I_0(\theta=0)$ and $I_0(\theta=\pi)$. These values are strongly dependent on the percentage of the P_D admixture^{19,20,23} in the deuteron wave function. Consequently experimental determination may offer a way for better defining the medium range shape of the tensor component of the N-N interaction.

The calculations of $I_0(\theta)$ are usually carried out under the assumption that the charge density operator may be expressed as a sum of standard one-body terms, and that two-body and relativistic corrections may be neglected (Siegert hypothesis). Because the values of the forward and backward differential cross sections are highly sensitive to the presence of relativistic and exchange contributions to the nucleon charge density operator^{15,21,22,24,25}, corresponding experimental data may help to ascertain the validity limits of the Siegert hypothesis.

Unfortunately, the above theoretical framework cannot be accurately tested by existing experimental data. In almost all experiments carried out so far, values of $I_0(\theta)$ have been measured only in the angular region of $25^\circ \leq \theta \leq 150^\circ$ (for a complete listing see Ref. 35). The notable exceptions are the three measurements of $I_0(0)$ and $I_0(\pi)$ (carried out by Hughes et al.³⁶, by Gilot et al.³⁷ and by Meyer et al.³⁸, and a few measurements of $\Sigma(\theta)$ ³⁹⁻⁴⁴. Moreover, all the available measurements of $I_0(\theta)$, except those carried out at the LADON facility in Frascati, have been produced by non-monochromatic beams with results that are not always consistent. Moreover the measurements of $\Sigma(\theta)$ are affected by uncertainties stemming from a poor knowledge of the degree of polarization P . Particular problems for the theoretical examination of the $d(\gamma, p)n$ reaction derive from the lack of reliable and firm experimental values for $I_0(\theta)$ and from the insufficient experimental information on $\Sigma(\theta)$ at energy $E_\gamma \lesssim 70$ MeV. It is the purpose of our research program to remedy this situation, at least partially.

4. - EXPERIMENTAL RESULTS AND DISCUSSION

Our experimental data are listed as a function of energy and angle in Tables II to IV. All reported

TABLE II - $[I_0(\theta_n)/\sigma_t]10^2$ as a function of the neutron CM angle $\theta_n = (\pi - \theta)$ and of the laboratory gamma-ray energy E_γ (MeV).

E_γ (MeV) \n θ_n (deg.)	19.8	29.0	38.6	60.8
15	1.71 \pm 0.26	1.96 \pm 0.11	2.42 \pm 0.21	-
30	2.81 \pm 0.22	2.96 \pm 0.20	3.99 \pm 0.34	5.21 \pm 0.38
45	5.79 \pm 0.55	5.03 \pm 0.22	5.04 \pm 0.29	-
60	8.36 \pm 0.42	7.82 \pm 0.34	8.44 \pm 0.78	6.36 \pm 0.41
90	11.72 \pm 0.44	10.92 \pm 0.23	10.99 \pm 0.39	9.99 \pm 0.48
120	9.16 \pm 0.44	10.32 \pm 0.41	8.73 \pm 0.50	10.22 \pm 0.50
135	6.91 \pm 0.69	7.57 \pm 0.25	8.08 \pm 0.52	-
150	4.08 \pm 0.30	4.96 \pm 0.29	6.03 \pm 0.44	6.22 \pm 0.36
165	2.15 \pm 0.30	2.82 \pm 0.15	3.75 \pm 0.29	-

TABLE III - $[I_1(\theta_n)/\sigma_t]10^2$ as a function of the neutron CM angle $\theta_n = (\pi - \theta)$ and of the laboratory gamma-ray energy E_γ (MeV).

E_γ (MeV) \ θ_n (deg.)	19.8	29.0	38.6	60.8
15	0.63 \pm 0.22	0.62 \pm 0.12	0.32 \pm 0.26	-
30	2.11 \pm 0.21	1.66 \pm 0.21	1.81 \pm 0.35	0.80 \pm 0.37
45	4.72 \pm 0.51	3.65 \pm 0.23	2.54 \pm 0.29	-
60	7.11 \pm 0.41	6.06 \pm 0.35	5.62 \pm 0.79	2.07 \pm 0.40
90	10.74 \pm 0.44	8.94 \pm 0.23	7.74 \pm 0.40	4.97 \pm 0.46
120	8.01 \pm 0.44	8.31 \pm 0.41	6.29 \pm 0.50	5.08 \pm 0.49
135	5.66 \pm 0.62	5.57 \pm 0.27	5.05 \pm 0.53	-
150	3.00 \pm 0.29	3.05 \pm 0.29	3.40 \pm 0.44	1.56 \pm 0.39
165	1.04 \pm 0.25	0.90 \pm 0.15	1.29 \pm 0.42	-

TABLE IV - The azimuthal asymmetry factor $\Sigma(\theta_n) = I_1(\theta_n)/I_0(\theta_n)$ as a function of the neutron CM angle $\theta_n = (\pi - \theta)$ and of the laboratory gamma-ray energy E_γ (MeV).

E_γ (MeV) \ θ_n (deg.)	19.8	29.0	38.6	60.8
15	0.370 \pm 0.054	0.316 \pm 0.053	0.134 \pm 0.104	-
30	0.752 \pm 0.036	0.562 \pm 0.051	0.453 \pm 0.076	0.154 \pm 0.064
45	0.816 \pm 0.020	0.725 \pm 0.023	0.504 \pm 0.050	-
60	0.851 \pm 0.022	0.775 \pm 0.024	0.666 \pm 0.072	0.326 \pm 0.053
90	0.917 \pm 0.007	0.819 \pm 0.013	0.704 \pm 0.020	0.497 \pm 0.039
120	0.874 \pm 0.019	0.805 \pm 0.021	0.721 \pm 0.036	0.497 \pm 0.043
135	0.819 \pm 0.019	0.736 \pm 0.018	0.625 \pm 0.050	-
150	0.737 \pm 0.030	0.615 \pm 0.041	0.564 \pm 0.057	0.251 \pm 0.067
165	0.482 \pm 0.062	0.320 \pm 0.046	0.344 \pm 0.076	-

results represent the statistically weighted mean of a number (from two to five) of separated measurements. Quoted errors are mainly due to counting statistics; in a few cases (a few extreme θ_n angles) the errors deriving from uncertainties in the subtraction of the residual flat background in the time of flight spectra had also to be taken into account. Systematic errors essentially derive from the Montecarlo evaluation of the K_1 and K_2 correction factors (see Eqs. (9, 10)). Conservative estimates lead us to conclude that, assuming $\Delta K_1/K_1 = \Delta K_2/K_2$:

$$(\Delta I_0)_{\text{syst}}/I_0 \approx \frac{\Delta K_1}{K_1} \left(\frac{1 + \Sigma^2}{2} \right)^{\frac{1}{2}},$$

$$(\Delta I_1)_{\text{syst}}/I_1 \sim \frac{(\Delta I_0)_{\text{syst}}}{\Sigma I_0} ,$$

$$(\Delta \Sigma)_{\text{syst}} \sim 2^{-\frac{1}{2}} \frac{\Delta K_1}{K_1} (1 - \Sigma^2).$$

Thus, by assuming $\Delta K_1/K_1 \approx 5\%$, systematic errors are expected to be of the same order as the statistical errors around $\theta = \pi/2$, and considerably lower elsewhere.

Our results for $(I_0/\sigma_t)10^2$ and $(I_1/\sigma_t)10^2$ as a function of the CM neutron angle θ_n , for the four laboratory photon energies used in our experiment, are plotted in Fig. 4, a through d. The behaviour of the azimuthal asymmetry factor $\Sigma(\theta_n)$ is shown in Fig. 5, a through d.

For comparison purposes, two theoretical curves are presented in all figures: the dashed lines have been obtained by Arenhövel^{5,10}, and independently by Cambi et al.^{11,16}, using the Reid soft-core (RSC) N-N potential⁴⁵. According to Ref. 11, no appreciable difference with respect to this result is obtained by using the Hamada-Johnston⁴⁶ or the Paris potential⁴⁷. On the other hand, the




FIG. 4 - Plots of $[I_0(\theta_n)/\sigma_t]10^2$ and of $[I_1(\theta_n)/\sigma_t]10^2$ as a function of the CM neutron angles $\theta_n = (\pi - \theta)$ at: (a) $E_\gamma = 19.8$ MeV; (b) 29.0 MeV; (c) 38.6 MeV; (d) 60.8 MeV. Dashed and solid lines represent theoretical calculations of Refs. 10 and 11 with the Reid soft-core (RSC) and the De Tourreil-Sprung (DTS-B) potentials, respectively.




FIG. 5 - Plots of the azimuthal asymmetry factor $\Sigma(\theta_n)$ as a function of the CM neutron angles $\theta_n = \pi - \theta$. The meaning of symbols is the same as in Fig. 4.

full lines^{11,16} obtained by using the version B of the DeTourreil-Sprung (DTS-B) potential⁴⁸ are higher as far as $I_1(\theta_n)$ and $\Sigma(\theta_n)$ are concerned, the difference increasing with the energy. In all these calculations, multipoles up to $L=4$ have been included and MEC (and where indicated also IC) contributions have been added to the standard (or "normal") Partovi theory⁴⁹. These MEC effects are more pronounced in $I_1(\theta)$ and $\Sigma(\theta)$ than in $I_0(\theta)$ and rapidly increase with E_γ energy. Relativistic or two-body corrections have not been added to the usual one-body charge operator (Siebert hypothesis).

The role of the mesonic and isobar degrees of freedom in this electromagnetic process is clearly shown in Fig. 6, where a plot of the asymmetry $\Sigma(\theta_n = \pi/2)$ versus the laboratory γ -ray energy is given⁴². Our data are compared with the results of earlier experiments^{39,41,43,44} and with the theoretical RSC calculation carried out by Arenhövel¹⁰. The dashed line (N=normal) has been obtained in the standard Partovi approximation, the solid line (N + MEC + IC) contains MEC + IC corrections to the current operator. Moreover, as shown in Ref. 13, relativistic charge corrections may also sensibly affect the theoretical curves at the highest energies.

In Fig. 7, a through d, our data are compared with those of other authors. Fig. 7a shows our data at 19.8 MeV in comparison with the 20 MeV results of Halpern and Weinstock⁵⁰ and Shin et al.⁵¹. In




FIG. 6 - Plot of the asymmetry factor $\Sigma(\theta_n = \pi/2)$ vs. laboratory gamma-ray energy (MeV). Our present data (solid circles) are compared with the results of earlier experiments of Liu (solid and open triangles, Ref. 39), Del Bianco et al. (inverted open triangle, Ref. 41), our previous data (open circles, Ref. 43), Gorbenko et al. (open squares, Ref. 44). The theoretical curves have been obtained in Ref. 10 with the RSC potential. The dashed line corresponds to the standard Pärtovi approximation (Ref. 49); the solid line reflects the inclusion of MEC and IC corrections.




FIG. 7 - A survey of the experimental determinations of $[I_0(\theta_n)/\sigma_t] 10^2$ for: (a) $E_\gamma = 19.8$ MeV; (b) 29.0 MeV; (c) 38.6 MeV; (d) 60.8 MeV. Our data (solid circles) are compared with results of earlier measurements of Halpern and Weinstock (stars, Ref. 50); Shin et al. (open triangles, Ref. 51); Weissman et al. (open circles, Ref. 33); Allen (inverted solid triangles, Ref. 52); Galey (solid triangles, Ref. 53). Corresponding theoretical curves obtained in Refs. 10 and 11 with the RSC (dashed lines) and the DTS-B (solid lines) potentials are also shown.

Fig. 7b the 30 MeV data of Shin⁵¹ and Weissman and Shultz³³ are reported, together with Allen⁵² results interpolated for 29.0 MeV; Fig. 7c shows the 40 MeV results of Refs. 51 and 33. Finally, Fig. 7d displays the 60 MeV data obtained by Galey⁵³ and the data of Shin⁵¹ interpolated to 60.8 MeV. As can be readily checked, these data are in substantial agreement.

A standard least-squares procedure was used to fit the experimental points $I_0(\theta)/\sigma_t$ with the sum of Legendre-polynomials $P_l(\cos \theta)$, and those of $I_1(\theta)/\sigma_t$ with the sum of associated Legendre-functions $P_l^1(\cos \theta)$, as shown in Eqs. (4).

The results obtained for the coefficients A_i/σ_t ($i=1, 2, 3$) are displayed in Fig. 8 versus the laboratory γ -ray energy, together with the results obtained from a reexamination of the earlier measurements. The error bars shown are derived from the least-squares procedure. The coefficient A_4 was fixed in all cases at its theoretical DTS-B estimate because its value is small and the available data are not sufficiently precise to give a meaningful value for A_4 . It was checked that generally this procedure introduces negligible changes in A_1 , A_2 and A_3 ; in particular, no significant change resulted from substituting the RSC for the DTS predictions for A_4 .




FIG. 8 - Plots of a representative sampling of coefficients A_i/σ_t (see Eq. 4a) vs. the laboratory gamma-ray energy (MeV). Our data (solid circles) are compared with results of measurements by other groups: all symbols have the same meaning as in Fig. 7. In addition, the open squares represent the data of Alexandrov et al., Ref. 56. Corresponding theoretical curves obtained in Refs. 10 and 11 with the RSC (dashed lines) and the DTS-B (solid lines) potentials are also shown. All original data have been reanalyzed in the present work by fixing A_4/σ_t to its theoretical value, shown in part (d) of this figure.

Theoretical curves obtained in Refs. 5 and 10 and Refs. 11 and 16 with the RSC and DTS-B potentials, including MEC + IC effects in the Siegert approximation, are also shown in Fig. 8. The ratio A_1/σ_t seems to be consistent with the theory, while the value of A_2/σ_t displays a significant difference at $E_\gamma \gtrsim 50$ MeV. This result has already been pointed-out in Ref. 35. Preliminary estimates¹¹ of the relativistic and pion corrections to the charge density operator seem to enhance $|A_2|$ ($A_2 < 0$) by a substantial amount. Similar arguments hold for A_3 , but here the situation appears to be less definite, owing to the size of the quoted errors.

The results obtained for the coefficients B_i/σ_t ($i=2, 3$) are reported in Fig. 9, together with the RSC and DTS-B theoretical predictions. The coefficient B_4 was fixed at its theoretical value. In the Siegert approximation the 29.0 and 38.6 MeV data seem to give a preference to the RSC interaction. This point will be examined further on.





FIG. 9 - Plots of the coefficients B_i/σ_t (see Eq. (4b)) vs. the laboratory gamma-ray energy (MeV). Our data for B_2/σ_t and B_3/σ_t (solid circles) are compared with results of theoretical calculations of Refs. 10 and 11 based on the RSC (dashed lines) and DTS-B (solid lines) potentials, respectively. The coefficient B_4/σ_t was fixed at its DTS-B theoretical value.

Additional information that can shed some more light on the details of the problem may be obtained from the coefficients a, \dots, h of the angular distributions formulation introduced by Partovi⁴⁹ (see Eqs. (5)).



Least-square fitted coefficients $a/\sigma_t, \dots, d/\sigma_t$ are compared directly with their predicted values in Fig. 10. The coefficient e was kept fixed. Furthermore, at $E_\gamma = 60.8$ MeV, the coefficient c also remained completely undetermined and had to be fixed at its theoretical value (cross in Fig. 10). In this case only the determination of d is sensibly affected

FIG. 10 - Plots of the coefficients $a/\sigma_t, b/\sigma_t, c/\sigma_t, d/\sigma_t$ (see Eq. 5a) vs. the laboratory gamma-ray energy (MeV). Our data (solid circles) are compared with theoretical curves obtained in Refs. 10 and 11 with the RSC (dashed lines) and the DTS-B (solid lines) potentials, respectively. The coefficient e/σ_t , which is not shown, was fixed at its DTS-B theoretical value.

by the assumed c -value. The bulk of the obtained data, in particular the a/σ_t values at 19.8, 29.0, 38.6 MeV and the b/σ_t , d/σ_t coefficients at 29.0 MeV, seem again to favour the RSC potential, at least in a framework of the Siegert hypothesis. However, our estimate of a and c coefficients must be regarded with a particular caution, owing to the following arguments:

- a) Our estimates of $I_0(\theta=0) = a+c$ (obtained by multiplying our results for the σ_t estimates of Ref. 11) do not entirely agree with the corresponding Mainz³⁶ and Louvain³⁷ results.
- b) Although, the two extreme angular points $\theta_n=15^\circ$ and 165° are of particular importance, the values obtained for a/σ_t (and for c/σ_t) are still critically dependent on the random and systematic errors of all the measured points.

Thus, there is clearly a need for new independent measurements of a (and c), that is of $I_0(\theta=0)$ and $I_0(\theta=\pi)$. This point will be discussed further in the next section. Also the disagreement observed for d/σ_t must be further explored, both on the theoretical and experimental side.

Our experimental estimates for the f/σ_t and g/σ_t coefficients is reported in Fig. 11 together with the corresponding theoretical RSC and DTS-B estimates. The small coefficient h , practically without influence in the fit, has been fixed. Just as the corresponding B_2/σ_t and B_3/σ_t coefficients, the estimates of f/σ_t and g/σ_t at 29.0 and 38.6 MeV also seem to favor the RSC potential.




FIG. 11 - Plots of the coefficients f/σ_t , g/σ_t , h/σ_t (see Eq. 5b) vs. the laboratory gamma-ray energy (MeV). Our data for f/σ_t and g/σ_t are compared with theoretical curves obtained in Refs. 10 and 11 with the RSC (dashed lines) and the DTS-B (solid lines) potentials, respectively. The coefficient h/σ_t was fixed at its DTS-B theoretical values.

A summary of all coefficients obtained in our fits is shown in Table V and VI. Only statistical errors are appended.

TABLE V - Coefficients of the expansion for $[I_0(\theta)/\sigma_t] 10^2$ given by Eqs. (4a) and (5a).

E_γ (MeV)	$(A_1/\sigma_t)10^2$	$(A_2/\sigma_t)10^2$	$(A_3/\sigma_t)10^2$	$(a/\sigma_t)10^2$	$(b/\sigma_t)10^2$	$(c/\sigma_t)10^2$	$(d/\sigma_t)10^2$
19.8	$1.44^{+0.21}$	$6.75^{+0.26}$	$1.20^{+0.33}$	$1.02^{+0.15}$	$10.86^{+0.37}$	$0.25^{+0.16}$	$2.99^{+0.64}$
29.0	$1.54^{+0.14}$	$6.10^{+0.10}$	$1.30^{+0.19}$	$1.65^{+0.09}$	$10.14^{+0.22}$	$0.24^{+0.03}$	$3.22^{+0.25}$
38.6	$1.47^{+0.22}$	$5.21^{+0.18}$	$0.89^{+0.31}$	$2.49^{+0.18}$	$9.09^{+0.39}$	$0.58^{+0.21}$	$2.20^{+0.77}$
60.8	$1.60^{+0.29}$	$3.66^{+0.42}$	$2.69^{+0.59}$	$3.90^{+0.38}$	$7.32^{+0.62}$	-	$2.93^{+0.41}$

TABLE VI - Coefficients of the expansion for $[I_1(\theta)/\sigma_t] 10^2$ given by Eqs. (4b) and (5b).

E_γ (MeV)	$(B_2/\sigma_t)10^2$	$(B_3/\sigma_t)10^2$	$(f/\sigma_t)10^2$	$(g/\sigma_t)10^2$
19.8	$3.43^{+0.09}$	$0.22^{+0.03}$	$10.83^{+0.26}$	$3.29^{+0.43}$
29.0	$3.04^{+0.06}$	$0.19^{+0.02}$	$9.91^{+0.16}$	$2.85^{+0.35}$
38.6	$2.55^{+0.07}$	$0.21^{+0.03}$	$8.64^{+0.25}$	$3.19^{+0.55}$
60.8	$1.59^{+0.10}$	$0.23^{+0.05}$	$6.19^{+0.30}$	$3.38^{+0.69}$

5. - CONCLUSIONS

The bulk of the existing data (see, i.e. Fig. 6) clearly shows the role of the non nucleonic degrees of freedom in the deuteron photodisintegration. Moreover, the measured values of the A_1/σ_t , B_1/σ_t and a/σ_t , ..., g/σ_t coefficients lie generally in the theoretically predicted ranges, with a few exceptions. However, some points require a particular attention. The present data show a better agreement with predictions that use the RSC and Paris potentials, rather than with those that use DTS-B, at least in the framework of the Siegert approximation. Furthermore, the coefficients d/σ_t at 38.6 and 60.8 MeV and g/σ_t at 29.0 MeV seem to be lower than both the DTS-B and RSC (Paris) estimates.

On the other hand, all theoretical predictions at higher energies could be appreciably modified by the inclusion of relativistic and two-body exchange contributions to the charge density operator. In addition, the two-body charge effects, particularly as far as the a and c coefficients are concerned, are markedly different if pseudoscalar (PS) or pseudovector (PV) coupling are assumed^{10,15,16}. Thus a definitive analysis of the data presently on hand must be postponed until extensive theoretical calculations for different N-N potentials become available. In particular, one can hope that a careful examination of the a and c coefficients (and perhaps of the other ones) will permit an unequivocal choice between PS and PV couplings and a new determination of the D-wave asymptotic A_D amplitude⁵⁴, strictly connected to the deuteron D-state percentage P_D .

Our final remarks concern the a, ..., g coefficients. Their extraction from the experimental data is considerably more uncertain than the determination of the A_1 and B_1 coefficients because in the fitting procedure of $I_0(\theta)$ and $I_1(\theta)$ they appear to be strongly correlated. Keeping this point in mind, let us further recall that our estimates of a/σ_t seem to be higher than the corresponding Mainz results³⁶, normalized by assuming standard σ_t values³⁵. Recent calculations including relativistic and two-body charge contributions^{11,15,16} seem to suggest that our results can be almost partially reproduced by assuming a PV coupling and using potentials that give P_D percentages in the range (6-8)%. However, our estimates of a/σ_t and of c/σ_t are rather indirect and must be tested by further direct measurements of $I_0(\theta=0)$ and $I_0(\theta=\pi)$.

As a concluding remark, let us recall that besides the well known dependence on the standard parameters characterizing the N-N potentials, the deuteron photodisintegration angular distributions $I_0(\theta)$ and $I_1(\theta)$ could be explicitly affected by the internal nucleonic degrees of freedom, that is by their internal quark structure. To the best of our knowledge only the influence of the nucleon structure on the total photodisintegration cross section σ_t has been so far studied, and in a preliminary way⁵⁵ at that. This point too must be investigated further before any definitive conclusions on the role of non-nucleonic degrees of freedom in deuteron photodisintegration may be extracted from our data.

ACKNOWLEDGEMENTS

We wish to thank the technical staff of the Ladon Group, E. Cima, A. Dante, L. Ruggieri and E. Turri, for their dedicated help throughout this experiment. The expert operation of the storage ring by the Adone staff is warmly acknowledged.

REFERENCES

- (1) M.L. Rustgi, O.P. Rustgi and T.S. Sandhu, *Can J. Phys.* 55, 158 (1977).
- (2) J.M. Laget, *Nucl. Phys.* A312, 265 (1978).
- (3) H. Arenhövel, in "Few body systems and electromagnetic interactions", ed. by C. Ciofi degli Atti and E. De Sanctis, *Lecture Notes in Physics*, Vol. 86 (Springer Verlag 1978).
- (4) K. Ogawa, T. Kamae and K. Nakamura, *Nucl. Phys.* A340, 451 (1980).
- (5) H. Arenhövel, *Z. Phys.* A302, 25 (1981).
- (6) M. Anastasio and M. Chemtob, *Nucl. Phys.* A364, 219 (1981).
- (7) A. Cambi, B. Mosconi and P. Ricci, *Phys. Rev.* C23, 992 (1981).
- (8) Reeta Vyas, Manoj Chopra and M.L. Rustgi, *Phys. Rev.* C25, 1801 (1982).
- (9) Reeta Vyas and M.L. Rustgi, *Phys. Rev.* C26, 1399 (1982).
- (10) H. Arenhövel, *Nucl. Phys.* A374, 521c (1982); *Nuovo Cimento* 76A, 256 (1983); Contribution to the Workshop on "Perspectives in Nuclear Physics at Intermediate Energies", Trieste, October 1983 and private communication.
- (11) A. Cambi, B. Mosconi and P. Ricci, *Phys. Rev.* C26, 2358 (1982); *Journ. of Phys.* G10, L11 (1984) and private communication.
- (12) M.L. Rustgi, Reeta Vyas and Manoj Chopra, *Phys. Rev. Letters* 50, 236 (1983).
- (13) M.L. Rustgi and Reeta Vyas, *Phys. Letters* 121B, 365 (1983).
- (14) M.L. Rustgi, R.D. Nunemaket and R.D. Sharma, *Nuovo Cimento* 77A, 317 (1983).
- (15) J.L. Friar, B.F. Gibson and G.L. Payne, *Phys. Rev.* C30, 441 (1984).
- (16) A. Cambi, B. Mosconi and P. Ricci, Contribution to the Workshop on "Perspectives in Nuclear Physics at Intermediate Energies", Trieste, October 1983 and private communication.
- (17) W.Y.P. Hwang, J.T. Londergan and G.E. Walker, *Ann. of Phys.* 149, 335 (1983).
- (18) M.L. Rustgi, Reeta Vyas and O.P. Rustgi, *Phys. Rev.* C29, 785 (1984).
- (19) M.L. Rustgi, T.S. Sandhu and O.P. Rustgi, *Phys. Letters* 70B, 145 (1977); *Phys. Rev.* C20, 24 (1979).
- (20) E.L. Lomon, *Phys. Letters* 68B, 419 (1977).
- (21) M. Gari and B. Sommer, *Phys. Rev. Letters* 41, 22 (1978).
- (22) E. Hadjimichael, *Phys. Letters* 85B, 17 (1979).
- (23) H. Arenhövel and W. Fabian, *Phys. Rev. Letters* 47, 34 (1981).
- (24) W. Jaus and W.S. Woolcock, *Nucl. Phys.* A365, 477 (1981).
- (25) A. Cambi, B. Mosconi and P. Ricci, *Phys. Rev. Letters* 48, 462 (1982).
- (26) L. Federici, G. Giordano, G. Matone, G. Pasquariello, P.G. Picozza, R. Caloi, L. Casano, M.P. De Pascale, M. Mattioli, E. Poldi, C. Schaerf, M. Vanni, P. Pelfer, D. Prospero, S. Frullani and B. Girolami; *Nuovo Cimento* B59, 247 (1980).
- (27) M.P. De Pascale, G. Giordano, G. Matone, P. Picozza, R. Caloi, L. Casano, M. Mattioli, E. Poldi, D. Prospero, and C. Schaerf, *Appl. Phys.* B28, 151 (1982).
- (28) O.F. Kulikov, Y.Y. Telnov, E.I. Filippov and M.N. Yakimenko, *Phys. Letters* 13, 344 (1964).
- (29) C. Bemporad, R.H. Milburn, N. Tanaka and M. Fotino, *Phys. Rev.* 138, 1546 (1965).
- (30) J. Ballam, G.B. Chadwick, R. Gearhart, Z.G.T. Guiragossian, P.R. Klein, A. Levy, M. Menke, J.J. Murray, P. Seyboth, G. Wolf, C.K. Sinclair, H.H. Bingham, W.B. Fretter, K.C. Moffeit, W.J. Podolsky, M.S. Rabin, A.H. Rosenfeld and R. Windmolders, *Phys. Rev. Letters* 23, 498 (1969).
- (31) G. Matone and A. Tranquilli, *LNF-75/21* (1975).
- (32) M.P. De Pascale, G. Giordano, G. Matone, P. Picozza, R. Caloi, L. Casano, M. Mattioli, E. Poldi, D. Prospero and C. Schaerf, *Appl. Optics* 21, 2660 (1982).
- (33) B. Weissman and H.L. Schultz, *Nucl. Phys.* A174, 129 (1971).
- (34) J.E.E. Baglin, R.W. Carr, E.J. Bentz Jr. and C.P. Wu, *Nucl. Phys.* A201, 593 (1973).
- (35) M.P. De Pascale, G. Giordano, G. Matone, P. Picozza, L. Azario, R. Caloi, L. Casano, L. Inghrosso, M. Mattioli, E. Poldi, D. Prospero and C. Schaerf, *Phys. Letters* 119B, 30 (1982).
- (36) R. Hughes, A. Zieger, H. Wäffler and B. Ziegler, *Nucl. Phys.* A267, 329 (1976).
- (37) J.F. Gilot, A. Bol, P. Leleux, P. Lipnik and P. Macq, *Phys. Rev. Letters* 47, 304 (1984).
- (38) H.O. Meyer, J.R. Hall, H. Hugi, H.J. Karwowski, R.E. Pollock and P. Schwandt, *Phys. Rev. Letters* 52, 1759 (1984).
- (39) F.F. Liu, *Phys. Rev.* 138B, 1443 (1965).
- (40) G. Barbiellini, C. Bernardini, F. Felicetti and G.P. Murtas, *Phys. Rev.* 154, 988 (1967).
- (41) W. Del Bianco, H. Jeremie, M. Irshad and G. Kajrys, *Nucl. Phys.* A343, 121 (1980).
- (42) W. Del Bianco, L. Federici, G. Giordano, G. Matone, G. Pasquariello, P. Picozza, R. Caloi, L. Casano, M.P. De Pascale, L. Inghrosso, M. Mattioli, E. Poldi, C. Schaerf, P. Pelfer, D. Prospero, S. Frullani, B. Girolami and H. Jeremie, *Phys. Rev. Letters* 47, 1118 (1981).
- (43) M.P. De Pascale, L. Federici, G. Giordano, G. Matone, P. Picozza, L. Azario, R. Caloi, L. Casano, M. Mattioli, E. Poldi, D. Prospero and C. Schaerf, *Phys. Letters* 114B, 11 (1982).

- (44) V.G. Gorbenko, Yu.V. Zhebrovskij, L.Ya. Kolesnikov, A.L. Rubashkin and P.V. Sorokin, Nucl. Phys. A381, 330 (1982).
- (45) R.V. Reid Jr., Ann. Phys. 50, 411 (1968).
- (46) T. Hamada and J.D. Johnston, Nucl. Phys. B34, 382 (1962).
- (47) M. Lacombe, B. Løiseau, J.M. Richard, R. Vinh Mau, J. Côté, P. Pire's and R. De Turreil, Phys. Rev. C21, 861 (1980); Phys. Rev. C23, 2405 (1981).
- (48) R. De Turreil and D.W.L. Sprung, Nucl. Phys. A201, 193 (1973).
- (49) F. Partovi, Ann. Phys. 27, 79 (1964).
- (50) J. Halpern and E.V. Weinstöck, Phys. Rev. 91, 934 (1953).
- (51) Y.M. Shin, J.A. Rawlins, W. Buss and A.O. Ewwaraye, Nucl. Phys. A154, 482 (1970).
- (52) L. Allen Jr., Phys. Rev. 98, 705 (1955).
- (53) J.A. Galey, Phys. Rev. 117, 763 (1960).
- (54) M.E. Schulze, D.P. Saylor and R. Goloskie, Phys. Rev. C24, 2435 (1981).
- (55) E. Hadjimichael and D.P. Saylor, Phys. Rev. Letters 45, 1776 (1980).
- (56) Yu.A.A. Aleksandrov, N.B. Delone, L.I. Slovokhotov, G.A. Sokol and L.N. Shtarkov, JETP (Sov. Phys.) 6, 472 (1958).