

Strong dynamics in the large- N limit from string theory and from the lattice

Marco Panero

Institute for Theoretical Physics
ETH Zürich
Zürich, Switzerland

Frascati,
10 June 2010

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Outline

Physical motivation

The large- N limit

Lattice QCD

Results

Conclusions and outlook

Based on:

- ▶ M.P., *Thermodynamics of the QCD plasma and the large- N limit*, Phys. Rev. Lett. **103** 232001 (2009), [arXiv:0907.3719 [hep-lat]]

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Outline

Physical motivation

The large- N limit

Lattice QCD

Results

Conclusions and outlook

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The physical problem - I

- ▶ Due to asymptotic freedom in non-Abelian gauge theories [**Gross and Wilczek, 1973; Politzer, 1973**], hadronic matter is expected to undergo a change of state to a deconfined phase at sufficiently high temperatures or densities [**Cabibbo and Parisi, 1975; Collins and Perry, 1974**].
- ▶ Extensive experimental investigation through heavy ion collisions since the Eighties: first at AGS (BNL) and SPS (CERN), then at RHIC (BNL)
- ▶ Present experimental evidence from SPS and RHIC: a 'A new state of matter' has been created [**Heinz and Jacob, 2000, Arsene et al., 2004; Back et al., 2004; Adcox et al., 2004; Adams et al., 2005**]

The physical problem - II

- ▶ The plasma behaves as an almost ideal fluid [**Kolb and Heinz, 2003**] ('The most perfect liquid observed in Nature')

The physical problem - II

- ▶ The plasma behaves as an almost ideal fluid [**Kolb and Heinz, 2003**] ('The most perfect liquid observed in Nature')
- ▶ Forthcoming experiments at LHC (CERN) and FAIR (GSI) to provide a more detailed picture
- ▶ However, the theoretical understanding of the QCD plasma [**Rischke, 2003**] is still far from complete . . .

Theoretical approaches - I

- ▶ Relativistic fluidodynamics is a successful phenomenological description [**Romatschke, 2009**], but is not derived from QCD first principles
- ▶ The perturbative approach in thermal gauge theory has a non-trivial mathematical structure, involving odd powers of the coupling [**Kapusta, 1979**], as well as contributions from diagrams involving arbitrarily large numbers of loops [**Linde, 1980; Gross, Pisarski and Yaffe, 1980**] . . .
- ▶ . . . and shows poor convergence at the temperatures probed in experiments [**Kajantie, Laine, Rummukainen and Schröder, 2002**]
- ▶ Dimensional reduction [**Ginsparg, 1980; Appelquist and Pisarski, 1981**] to EQCD and MQCD [**Braaten and Nieto, 1995**], hard-thermal loop resummations [**Blaizot and Iancu, 2002**], and other effective theory approaches [**Kraemmer and Rebhan, 2004**]

Theoretical approaches - II

- ▶ Analytical progress in strongly interacting gauge theories: the AdS/CFT conjecture [**Maldacena, 1997**] and related theories as possible models for the non-perturbative features of QCD, including spectral [**Erdmenger, Evans, Kirsch and Threlfall, 2007**] and thermal properties [**Gubser and Karch, 2009**]
- ▶ In the large- N limit, the Maldacena conjecture relates a strongly interacting gauge theory to the classical limit of a gravity model

Theoretical approaches - III

- ▶ Numerical approach: Computer simulations of QCD regularized on a lattice allow first-principle, non-perturbative studies of the finite-temperature plasma
- ▶ The lattice determination of equilibrium thermodynamic properties in $SU(3)$ gauge theory is regarded as a solved problem [**Boyd et al., 1996**]
- ▶ In recent years, finite-temperature lattice QCD has steadily progressed towards parameters corresponding to the physical point [**Karsch et al., 2000; Ali Khan et al., 2001; Aoki et al., 2005; Bernard et al., 2006; Cheng et al., 2007; Bazavov et al., 2009**]**—see also [DeTar and Heller, 2009]** for a review of recent results

Goals of this work

- ▶ High-precision determination of the equilibrium thermodynamic properties in $SU(N \geq 3)$ Yang-Mills theories
- ▶ Comparison with holographic predictions
- ▶ Entropy deficit: comparison with a supergravity model in a strongly interacting, nearly conformal regime
- ▶ Investigation of possible non-perturbative contributions to the trace anomaly
- ▶ Extrapolation to the large- N limit

Related works: **[Bringoltz and Teper, 2005]** and **[Datta and Gupta, 2010]**

Outline

Physical motivation

The large- N limit

Lattice QCD

Results

Conclusions and outlook

ETH



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**'t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory . . .
- ▶ . . . in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], . . .
- ▶ . . . which is characterized by planar diagrams' dominance . . .

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**'t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory ...
- ▶ ... in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], ...
- ▶ ... which is characterized by planar diagrams' dominance ...



ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory ...
- ▶ ... in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], ...
- ▶ ... which is characterized by planar diagrams' dominance ...
- ▶ ... and amenable to a topological expansion in terms of surfaces of increasing genus

$$\mathcal{A} =$$

- ▶ Formal connection to string theory: loop expansion in Riemann surfaces for closed string theory with coupling constant $g_{\text{string}} \sim 1/N$ [**Aharony, Gubser, Maldacena, Ooguri and Oz, 1999; Mateos, 2007**]

$$\mathcal{A} = \sum_{G=0}^{\infty} N^{2-2G} \sum_{n=0}^{\infty} c_{G,n} \lambda^n$$

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory ...
- ▶ ... in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], ...
- ▶ ... which is characterized by planar diagrams' dominance ...
- ▶ ... and amenable to a topological expansion in terms of surfaces of increasing genus

$$\mathcal{A} = \textcircled{8}$$

- ▶ Formal connection to string theory: loop expansion in Riemann surfaces for closed string theory with coupling constant $g_{\text{string}} \sim 1/N$ [**Aharony, Gubser, Maldacena, Ooguri and Oz, 1999; Mateos, 2007**]

$$\mathcal{A} = \sum_{G=0}^{\infty} N^{2-2G} \sum_{n=0}^{\infty} c_{G,n} \lambda^n$$

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**'t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory ...
- ▶ ... in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], ...
- ▶ ... which is characterized by planar diagrams' dominance ...
- ▶ ... and amenable to a topological expansion in terms of surfaces of increasing genus

$$\mathcal{A} = \text{[torus]} + \text{[donut with Homer Simpson]}$$

- ▶ Formal connection to string theory: loop expansion in Riemann surfaces for closed string theory with coupling constant $g_{\text{string}} \sim 1/N$ [**Aharony, Gubser, Maldacena, Ooguri and Oz, 1999; Mateos, 2007**]

$$\mathcal{A} = \sum_{G=0}^{\infty} N^{2-2G} \sum_{n=0}^{\infty} c_{G,n} \lambda^n$$

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**'t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory ...
- ▶ ... in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], ...
- ▶ ... which is characterized by planar diagrams' dominance ...
- ▶ ... and amenable to a topological expansion in terms of surfaces of increasing genus



- ▶ Formal connection to string theory: loop expansion in Riemann surfaces for closed string theory with coupling constant $g_{\text{string}} \sim 1/N$ [**Aharony, Gubser, Maldacena, Ooguri and Oz, 1999; Mateos, 2007**]

$$\mathcal{A} = \sum_{G=0}^{\infty} N^{2-2G} \sum_{n=0}^{\infty} c_{G,n} \lambda^n$$

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory ...
- ▶ ... in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], ...
- ▶ ... which is characterized by planar diagrams' dominance ...
- ▶ ... and amenable to a topological expansion in terms of surfaces of increasing genus



- ▶ Formal connection to string theory: loop expansion in Riemann surfaces for closed string theory with coupling constant $g_{\text{string}} \sim 1/N$ [**Aharony, Gubser, Maldacena, Ooguri and Oz, 1999; Mateos, 2007**]

$$\mathcal{A} = \sum_{G=0}^{\infty} N^{2-2G} \sum_{n=0}^{\infty} c_{G,n} \lambda^n$$

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**'t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory ...
- ▶ ... in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], ...
- ▶ ... which is characterized by planar diagrams' dominance ...
- ▶ ... and amenable to a topological expansion in terms of surfaces of increasing genus



- ▶ Formal connection to string theory: loop expansion in Riemann surfaces for closed string theory with coupling constant $g_{\text{string}} \sim 1/N$ [**Aharony, Gubser, Maldacena, Ooguri and Oz, 1999; Mateos, 2007**]

$$\mathcal{A} = \sum_{G=0}^{\infty} N^{2-2G} \sum_{n=0}^{\infty} c_{G,n} \lambda^n$$

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

The old perspective: QCD at large N

- ▶ QCD has no obvious dimensionless expansion parameter (the coupling is used to set the scale)
- ▶ 't Hooft proposed to use $1/N$ (N being the number of colors) as an expansion parameter [**t Hooft, 1974**]
- ▶ Generically, a large- N limit can be interpreted as a 'classical limit'; identification of coherent states and construction of a classical Hamiltonian [**Yaffe, 1982**]
- ▶ The large- N limit of QCD, at fixed 't Hooft coupling $\lambda = g^2 N$ and fixed number of flavors N_f , is a simpler theory . . .
- ▶ . . . in which certain non-trivial non-perturbative features of QCD can be easily explained in terms of combinatorics [**Witten, 1979; Manohar, 1998**], . . .
- ▶ . . . which is characterized by planar diagrams' dominance . . .
- ▶ . . . and amenable to a topological expansion in terms of surfaces of increasing genus
- ▶ Formal connection to string theory: loop expansion in Riemann surfaces for closed string theory with coupling constant $g_{\text{string}} \sim 1/N$ [**Aharony, Gubser, Maldacena, Ooguri and Oz, 1999; Mateos, 2007**]

$$\mathcal{A} = \sum_{G=0}^{\infty} N^{2-2G} \sum_{n=0}^{\infty} c_{G,n} \lambda^n$$

The AdS/CFT correspondence

- ▶ Maldacena conjectured that the large- N limit of the maximally supersymmetric $\mathcal{N} = 4$ supersymmetric YM (SYM) theory in four dimensions is dual to type IIB string theory in a $AdS_5 \times S^5$ space **[Maldacena, 1997]**


$$ds^2 = \frac{r^2}{R^2} \left(-dt^2 + d\mathbf{x}^2 \right) + \frac{R^2}{r^2} dr^2 + R^2 d\Omega_5^2$$

- ▶ The conjecture arises from the observation that the low-energy dynamics of open strings ending on a stack of N D3 branes in $AdS_5 \times S^5$ can be described in terms of $\mathcal{N} = 4$ SYM
- ▶ Geometric interpretation: There exists a correspondence of symmetries in the two theories
- ▶ A highly non-trivial correspondence, linking the strongly coupled regime of field theory to the weak-coupling limit of a gravity model
- ▶ Identification of the generating functional of connected Green's functions in the gauge theory with the minimum of the supergravity action with given boundary conditions: correlation functions of gauge theory operators from perturbative calculations in the gravity theory **[Gubser, Klebanov and Polyakov, 1998]**
- ▶ A stringy realization of the holographic principle: the description of dynamics within a volume of space is "encoded on the boundary" **[t Hooft, 1993; Susskind, 1995]**—see also **[Bousso, 2002]** for a review
- ▶ The large- N limit of the $\mathcal{N} = 4$ SYM theory exhibits a phase transition which can be related to the thermodynamics of AdS black holes **[Witten, 1998]**



The AdS/CFT correspondence

- ▶ Maldacena conjectured that the large- N limit of the maximally supersymmetric $\mathcal{N} = 4$ supersymmetric YM (SYM) theory in four dimensions is dual to type IIB string theory in a $AdS_5 \times S^5$ space **[Maldacena, 1997]**
- ▶ The conjecture arises from the observation that the low-energy dynamics of open strings ending on a stack of N D3 branes in $AdS_5 \times S^5$ can be described in terms of $\mathcal{N} = 4$ SYM



- ▶ Geometric interpretation: There exists a correspondence of symmetries in the two theories
- ▶ A highly non-trivial correspondence, linking the strongly coupled regime of field theory to the weak-coupling limit of a gravity model
- ▶ Identification of the generating functional of connected Green's functions of the supergravity action with the minimum of the supergravity action with given boundary conditions

The AdS/CFT correspondence

- ▶ Maldacena conjectured that the large- N limit of the maximally supersymmetric $\mathcal{N} = 4$ supersymmetric YM (SYM) theory in four dimensions is dual to type IIB string theory in a $AdS_5 \times S^5$ space **[Maldacena, 1997]**
- ▶ The conjecture arises from the observation that the low-energy dynamics of open strings ending on a stack of N D3 branes in $AdS_5 \times S^5$ can be described in terms of $\mathcal{N} = 4$ SYM
- ▶ Geometric interpretation: There exists a correspondence of symmetries in the two theories
 - ▶ \mathcal{R} -symmetry in the gauge theory is $SU(4) \sim SO(6)$ symmetry of S^5
 - ▶ The conformal invariance group in the gauge theory is isomorphic to $SO(2, 4)$, the symmetry group of AdS_5
- ▶ A highly non-trivial correspondence, linking the strongly coupled regime of field theory to the weak-coupling limit of a gravity model
- ▶ Identification of the generating functional of connected Green's functions in the gauge theory with the minimum of the supergravity action with given boundary conditions: correlation functions of gauge theory operators from perturbative calculations in the gravity theory **[Gubser, Klebanov and Polyakov, 1998]**
- ▶ A stringy realization of the holographic principle: the description of dynamics within a volume of space is "encoded on the boundary" [**'t Hooft, 1993; Susskind, 1995**]**—see also [Bousso, 2002]** for a review
- ▶ The large- N limit of the $\mathcal{N} = 4$ SYM theory exhibits a phase transition which can be related to the thermodynamics of AdS black holes **[Witten, 1998]**



The AdS/CFT correspondence

- ▶ Maldacena conjectured that the large- N limit of the maximally supersymmetric $\mathcal{N} = 4$ supersymmetric YM (SYM) theory in four dimensions is dual to type IIB string theory in a $AdS_5 \times S^5$ space **[Maldacena, 1997]**
- ▶ The conjecture arises from the observation that the low-energy dynamics of open strings ending on a stack of N D3 branes in $AdS_5 \times S^5$ can be described in terms of $\mathcal{N} = 4$ SYM
- ▶ Geometric interpretation: There exists a correspondence of symmetries in the two theories
- ▶ A highly non-trivial correspondence, linking the strongly coupled regime of field theory to the weak-coupling limit of a gravity model

$$g^2 = 4\pi g_s$$
$$g^2 N = \frac{R^4}{l_s^4}$$

- ▶ Identification of the generating functional of connected Green's functions in the gauge theory with the minimum of the supergravity action with given boundary conditions: correlation functions of gauge theory operators from perturbative calculations in the gravity theory **[Gubser, Klebanov and Polyakov, 1998]**
- ▶ A stringy realization of the holographic principle: the description of dynamics within a volume of space is "encoded on the boundary" **[t Hooft, 1993; Susskind, 1995]**—see also **[Bousso, 2002]** for a review
- ▶ The large- N limit of the $\mathcal{N} = 4$ SYM theory exhibits a phase transition which can be related to the thermodynamics of AdS black holes **[Witten, 1998]**

The AdS/CFT correspondence

- ▶ Maldacena conjectured that the large- N limit of the maximally supersymmetric $\mathcal{N} = 4$ supersymmetric YM (SYM) theory in four dimensions is dual to type IIB string theory in a $AdS_5 \times S^5$ space **[Maldacena, 1997]**
- ▶ The conjecture arises from the observation that the low-energy dynamics of open strings ending on a stack of N D3 branes in $AdS_5 \times S^5$ can be described in terms of $\mathcal{N} = 4$ SYM
- ▶ Geometric interpretation: There exists a correspondence of symmetries in the two theories
- ▶ A highly non-trivial correspondence, linking the strongly coupled regime of field theory to the weak-coupling limit of a gravity model
- ▶ Identification of the generating functional of connected Green's functions in the gauge theory with the minimum of the supergravity action with given boundary conditions: correlation functions of gauge theory operators from perturbative calculations in the gravity theory **[Gubser, Klebanov and Polyakov, 1998]**
- ▶ A stringy realization of the holographic principle: the description of dynamics within a volume of space is “encoded on the boundary” [**'t Hooft, 1993; Susskind, 1995**—see also **[Bousso, 2002]** for a review
- ▶ The large- N limit of the $\mathcal{N} = 4$ SYM theory exhibits a phase transition which can be related to the thermodynamics of AdS black holes **[Witten, 1998]**

The AdS/CFT correspondence

- ▶ Maldacena conjectured that the large- N limit of the maximally supersymmetric $\mathcal{N} = 4$ supersymmetric YM (SYM) theory in four dimensions is dual to type IIB string theory in a $AdS_5 \times S^5$ space **[Maldacena, 1997]**
- ▶ The conjecture arises from the observation that the low-energy dynamics of open strings ending on a stack of N D3 branes in $AdS_5 \times S^5$ can be described in terms of $\mathcal{N} = 4$ SYM
- ▶ Geometric interpretation: There exists a correspondence of symmetries in the two theories
- ▶ A highly non-trivial correspondence, linking the strongly coupled regime of field theory to the weak-coupling limit of a gravity model
- ▶ Identification of the generating functional of connected Green's functions in the gauge theory with the minimum of the supergravity action with given boundary conditions: correlation functions of gauge theory operators from perturbative calculations in the gravity theory **[Gubser, Klebanov and Polyakov, 1998]**
- ▶ A stringy realization of the holographic principle: the description of dynamics within a volume of space is “encoded on the boundary” **[’t Hooft, 1993; Susskind, 1995]**—see also **[Bousso, 2002]** for a review
- ▶ The large- N limit of the $\mathcal{N} = 4$ SYM theory exhibits a phase transition which can be related to the thermodynamics of AdS black holes **[Witten, 1998]**

The AdS/CFT correspondence

- ▶ Maldacena conjectured that the large- N limit of the maximally supersymmetric $\mathcal{N} = 4$ supersymmetric YM (SYM) theory in four dimensions is dual to type IIB string theory in a $AdS_5 \times S^5$ space **[Maldacena, 1997]**
- ▶ The conjecture arises from the observation that the low-energy dynamics of open strings ending on a stack of N D3 branes in $AdS_5 \times S^5$ can be described in terms of $\mathcal{N} = 4$ SYM
- ▶ Geometric interpretation: There exists a correspondence of symmetries in the two theories
- ▶ A highly non-trivial correspondence, linking the strongly coupled regime of field theory to the weak-coupling limit of a gravity model
- ▶ Identification of the generating functional of connected Green's functions in the gauge theory with the minimum of the supergravity action with given boundary conditions: correlation functions of gauge theory operators from perturbative calculations in the gravity theory **[Gubser, Klebanov and Polyakov, 1998]**
- ▶ A stringy realization of the holographic principle: the description of dynamics within a volume of space is “encoded on the boundary” **[’t Hooft, 1993; Susskind, 1995]**—see also **[Bousso, 2002]** for a review
- ▶ The large- N limit of the $\mathcal{N} = 4$ SYM theory exhibits a phase transition which can be related to the thermodynamics of AdS black holes **[Witten, 1998]**

Non-perturbative predictions for QCD-like theories from holographic models

- ▶ ‘Top-down’ approach: break some symmetries of the $\mathcal{N} = 4$ theory explicitly, add fundamental matter fields to the gauge theory by including new branes in the string theory [**Bertolini, Di Vecchia, Frau, Lerda, and Marotta, 2001**; **Graña and Polchinski, 2001**; **Karch and Katz, 2002**] to get a non-trivial hadron sector with ‘mesons’ and χ SB [**Erdmenger, Evans, Kirsch and Threlfall, 2007**]
- ▶ Description of hydrodynamic and thermodynamic properties for a strongly interacting system, like the QCD plasma, from gauge/gravity duality—see [**Son and Starinets, 2007**; **Mateos, 2007**; **Gubser and Karch, 2009**] and references therein
- ▶ ‘Bottom-up’ approach: construct a 5D gravitational background reproducing the main features of QCD [**Polchinski and Strassler, 2001**; **Erlich, Katz, Son and Stephanov, 2005**; **Da Rold and Pomarol, 2005**; **Karch, Katz, Son and Stephanov, 2006**]
- ▶ Hard-wall *versus* soft-wall AdS/QCD, and related thermodynamic features [**Herzog, 2007**]

Improved holographic QCD model – I

- ▶ Kiritsis and collaborators [**Gürsoy, Kiritsis, Mazzanti and Nitti, 2008**] proposed an AdS/QCD model based on a 5D Einstein-dilaton gravity theory, with the fifth direction dual to the energy scale of the $SU(N)$ gauge theory
- ▶ Field content on the gravity side: metric (dual to the $SU(N)$ energy-momentum tensor), the dilaton (dual to the trace of F^2) and the axion (dual to the trace of $F\tilde{F}$)
- ▶ Gravity action:

$$S_{IHQCD} = -M_P^3 N^2 \int d^5x \sqrt{g} \left[R - \frac{4}{3} (\partial\Phi)^2 + V(\lambda) \right] + 2M_P^3 N^2 \int_{\partial M} d^4x \sqrt{h} K$$

- ▶ Φ is the dilaton field, $\lambda = \exp(\Phi)$ is identified with the running 't Hooft coupling of the dual $SU(N)$ YM theory
- ▶ The effective five-dimensional Newton constant $G_5 = 1 / (16\pi M_P^3 N^2)$ becomes small in the large- N limit

Improved holographic QCD model – II

- ▶ Dilaton potential $V(\lambda)$ defined by requiring asymptotic freedom with a logarithmically running coupling in the UV and linear confinement in the IR of the gauge theory; a possible *Ansatz* is:

$$V(\lambda) = \frac{12}{\ell^2} \left[1 + V_0\lambda + V_1\lambda^{4/3} \sqrt{\log(1 + V_2\lambda^{4/3} + V_3\lambda^2)} \right],$$

where ℓ is the AdS scale (overall normalization), and two free parameters are fixed by imposing that the dual model reproduces the first two coefficients of the $SU(N)$ β -function

- ▶ Gauge/gravity duality expected to hold in the large- N limit only, because calculations in the gravity model neglect string interactions which can become important above a cut-off scale $M_P N^{2/3} \simeq 2.5$ GeV in $SU(3)$
- ▶ First-order transition from a thermal-graviton- to a black-hole-dominated regime in the 5D gravity theory dual to the $SU(N)$ deconfinement transition
- ▶ The model successfully reproduces the main non-perturbative spectral and thermodynamical features of the $SU(3)$ YM theory
- ▶ Can also be used to derive predictions for observables such as the plasma bulk viscosity, drag force and jet quenching parameter [**Gürsoy, Kiritsis, Michalogiorgakis and Nitti, 2009**]

Outline

Physical motivation

The large- N limit

Lattice QCD

Results


Conclusions and outlook

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a




- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables
- ▶ Lattice gauge action [Wilson, 1974]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \quad \text{with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit
- ▶ During the last decade, lattice QCD has entered an era of precision calculations.

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables




- ▶ Lattice gauge action [Wilson, 1974]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \quad \text{with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit
- ▶ During the last decade, lattice QCD has entered an era of precision calculations with large-scale simulations including light dynamical quarks [Davies et al., 2003]

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables



- ▶ Lattice gauge action [Wilson, 1974]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \quad \text{with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit
- ▶ During the last decade, lattice QCD has entered an era of precision calculations with large-scale simulations including light dynamical quark [Davies et al., 2003]

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables
- ▶ Lattice gauge action [**Wilson, 1974**]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \quad \text{with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit $a \rightarrow 0$
- ▶ During the last decade, lattice QCD has entered an era of precision calculations, with large-scale simulations including light dynamical quarks [**Davies *et al.*, 2003**; **Aubin *et al.*, 2004**]
- ▶ Results show striking agreement with experimental data (Wilczek: 'A stunning achievement'; Veltman: 'Wow, that's impressive!')

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables
- ▶ Lattice gauge action [**Wilson, 1974**]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \quad \text{with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit $a \rightarrow 0$
- ▶ During the last decade, lattice QCD has entered an era of precision calculations, with large-scale simulations including light dynamical quarks [**Davies *et al.*, 2003**; **Aubin *et al.*, 2004**]
- ▶ Results show striking agreement with experimental data (Wilczek: 'A stunning achievement'; Veltman: 'Wow, that's impressive!')

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables
- ▶ Lattice gauge action [**Wilson, 1974**]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \operatorname{Re} \operatorname{Tr} U_{\square} \right), \quad \text{with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit $a \rightarrow 0$
- ▶ During the last decade, lattice QCD has entered an era of precision calculations, with large-scale simulations including light dynamical quarks [**Davies *et al.*, 2003**; **Aubin *et al.*, 2004**]
- ▶ Results show striking agreement with experimental data (Wilczek: 'A stunning achievement'; Veltman: 'Wow, that's impressive!')

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables
- ▶ Lattice gauge action [**Wilson, 1974**]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \text{ with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit $a \rightarrow 0$
- ▶ During the last decade, lattice QCD has entered an era of precision calculations, with large-scale simulations including light dynamical quarks [**Davies *et al.*, 2003**; **Aubin *et al.*, 2004**]
- ▶ Results show striking agreement with experimental data (Wilczek: 'A stunning achievement'; Veltman: 'Wow, that's impressive!')

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables
- ▶ Lattice gauge action [**Wilson, 1974**]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \text{ with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit $a \rightarrow 0$
- ▶ During the last decade, lattice QCD has entered an era of precision calculations, with large-scale simulations including light dynamical quarks [**Davies et al., 2003**; **Aubin et al., 2004**]
- ▶ Results show striking agreement with experimental data (Wilczek: 'A stunning achievement'; Veltman: 'Wow, that's impressive!')

Lattice QCD: The basics

- ▶ Discretize a finite hypervolume in Euclidean spacetime by a regular grid with finite spacing a
- ▶ Transcribe gauge and fermion d.o.f. to lattice elements, build lattice observables
- ▶ Lattice gauge action [**Wilson, 1974**]:

$$S = \beta \sum_{\square} \left(1 - \frac{1}{N} \text{Re Tr } U_{\square} \right), \text{ with: } \beta = \frac{2N}{g_0^2}$$

- ▶ A gauge-invariant, non-perturbative regularization
- ▶ Amenable to numerical simulation: Sample configuration space according to a statistical weight proportional to $\exp(-S)$
- ▶ Physical results recovered by extrapolation to the continuum limit $a \rightarrow 0$
- ▶ During the last decade, lattice QCD has entered an era of precision calculations, with large-scale simulations including light dynamical quarks [**Davies et al., 2003**; **Aubin et al., 2004**]
- ▶ Results show striking agreement with experimental data (Wilczek: 'A stunning achievement'; Veltman: 'Wow, that's impressive!')

Thermodynamics on the lattice

- ▶ Thermal averages from simulations on a lattice with compactified Euclidean time direction, with $T = 1/(aN_\tau)$
- ▶ Pressure $p(T)$ via the 'integral method' [Engels *et al.*, 1990]:

$$\begin{aligned} p &= T \frac{\partial}{\partial V} \log \mathcal{Z} \simeq \frac{T}{V} \log \mathcal{Z} = \frac{1}{a^4 N_s^3 N_\tau} \int_{\beta_0}^{\beta} d\beta' \frac{\partial \log \mathcal{Z}}{\partial \beta'} \\ &= \frac{6}{a^4} \int_{\beta_0}^{\beta} d\beta' (\langle U_{\square} \rangle_T - \langle U_{\square} \rangle_0) \end{aligned}$$

Thermodynamics on the lattice

- ▶ Other thermodynamic observables obtained from indirect measurements

- ▶ Trace of the stress tensor $\Delta = \epsilon - 3p$:

$$\Delta = T^5 \frac{\partial}{\partial T} \frac{p}{T^4} = \frac{6}{a^4} \frac{\partial \beta}{\partial \log a} (\langle U_{\square} \rangle_0 - \langle U_{\square} \rangle_T)$$

- ▶ Energy density:

$$\epsilon = \frac{T^2}{V} \frac{\partial}{\partial T} \log \mathcal{Z} = \Delta + 3p$$

- ▶ Entropy density:

$$s = \frac{S}{V} = \frac{\epsilon - f}{T} = \frac{\Delta + 4p}{T}$$


Simulation details

- ▶ Lattice sizes $N_s^3 \times N_\tau$, with $N_s = 20$ or 16 , and $N_\tau = 5$
- ▶ Simulation algorithm: heat-bath [**Kennedy and Pendleton, 1985**] for $SU(2)$ subgroups [**Cabibbo and Marinari, 1982**] and full- $SU(N)$ overrelaxation [**Kiskis, Narayanan and Neuberger, 2003; Dürr, 2004; de Forcrand and Jahn, 2005**]
- ▶ Cross-check with $T = 0$ simulations run using the Chroma suite [**Edwards and Joó, 2004**]
- ▶ Physical scale for $SU(3)$ set using r_0 [**Necco and Sommer, 2001**]
- ▶ Physical scale for $SU(N > 3)$ set using known values for the string tension σ [**Lucini, Teper and Wenger, 2004; Lucini and Teper, 2001**] in combination with the 3-loop lattice β -function [**Allés, Feo and Panagopoulos, 1997; Allton, Teper and Trivini, 2008**] in the mean-field improved lattice scheme [**Parisi, 1980; Lepage and Mackenzie, 1993**]

Measurements of the plaquette

- High precision determination of $(\langle U_{\square} \rangle_T - \langle U_{\square} \rangle_0)$ required


$SU(3)$, $N_s = 20$, $N_\tau = 5$



Measurements of the plaquette

- High precision determination of $(\langle U_{\square} \rangle_T - \langle U_{\square} \rangle_0)$ required


$$\text{SU}(4), N_s = 16, N_\tau = 5$$



Measurements of the plaquette

- ▶ High precision determination of $(\langle U_{\square} \rangle_T - \langle U_{\square} \rangle_0)$ required
- ▶ Data reveal a strong first order bulk transition for $SU(N \geq 4)$

$$SU(6), N_s = 16, N_t = 5$$



Outline

Physical motivation

The large- N limit

Lattice QCD


Results

Conclusions and outlook


ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Improved holographic QCD model vs. lattice data




Improved holographic QCD model vs. lattice data




Improved holographic QCD model vs. lattice data

Trace of the energy-momentum tensor




Improved holographic QCD model vs. lattice data



Improved holographic QCD model vs. lattice data


Entropy density



AdS/CFT vs. lattice data in a 'quasi-conformal' regime

For $T \simeq 3T_C$, the lattice results reveal that the deconfined plasma, while still strongly interacting and far from the Stefan-Boltzmann limit, approaches a scale-invariant regime ...

$p(\epsilon)$ equation of state and approach to conformality




AdS/CFT vs. lattice data in a 'quasi-conformal' regime

... in which the entropy density is comparable with the supergravity prediction for $\mathcal{N} = 4$ SYM [Gubser, Klebanov and Tseytlin, 1998]

$$\frac{s}{s_0} = \frac{3}{4} + \frac{45}{32} \zeta(3) (2\lambda)^{-3/2} + \dots$$

Entropy density vs. 't Hooft coupling




AdS/CFT vs. lattice data in a 'quasi-conformal' regime

... in which the entropy density is comparable with the supergravity prediction for $\mathcal{N} = 4$ SYM [Gubser, Klebanov and Tseytlin, 1998]

$$\frac{s}{s_0} = \frac{3}{4} + \frac{45}{32} \zeta(3) (2\lambda)^{-3/2} + \dots$$

Entropy density vs. 't Hooft coupling




Note that a comparison of $\mathcal{N} = 4$ SYM and full-QCD lattice results for the drag force on heavy quarks also yields $\lambda \simeq 5.5$ [Gubser, 2006]

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

T^2 contributions to the trace anomaly?

The trace anomaly reveals a characteristic T^2 -behavior, possibly of non-perturbative origin [Megías, Ruiz Arriola and Salcedo, 2003; Pisarski, 2006; Andreev, 2007]




Extrapolation to $N \rightarrow \infty$

Based on the parametrization [Bazavov *et al.*, 2009]:

$$\frac{\Delta}{T^4} = \frac{\pi^2}{45} (N^2 - 1) \cdot \left(1 - \left\{ 1 + \exp \left[\frac{(T/T_c) - f_1}{f_2} \right] \right\}^{-2} \right) \left(f_3 \frac{T_c^2}{T^2} + f_4 \frac{T_c^4}{T^4} \right)$$

Extrapolation to the large- N limit



Outline

Physical motivation

The large- N limit

Lattice QCD

Results

Conclusions and outlook

ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Conclusions

- ▶ Equilibrium thermodynamic observables in $SU(N)$ YM theories at temperatures $0.8T_c \leq T \leq 3.4T_c$ show a mild dependence on N
- ▶ Successful comparison with the IHQCD model
- ▶ Quasi-conformal regime of YM and $\mathcal{N} = 4$ SYM predictions—Can lattice data help to pin down realistic parameters for AdS/CFT models of the sQGP?
[Noronha, Gyulassy and Torrieri, 2009]
- ▶ Δ seems to have a T^2 dependence also at large N
- ▶ Extrapolation to the $N \rightarrow \infty$ limit

Projects for the future



ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Projects for the future - I

(in case 'plan A' fails ...)


- ▶ $SU(N)$ screening masses and spatial string tensions, comparisons with AdS/CFT [**Bak, Karch and Yaffe, 2007**] and with IHQCD [**Alanen, Kajantie and Suur-Uski, 2009**]
- ▶ $\text{Tr}F^2$ correlators and dilaton potential [**Noronha, 2009**]
- ▶ Observables related to thermodynamic fluctuations: specific heat, speed of sound *et c.* [**Gavai, Gupta and Mukherjee, 2005**]*—*relevant for the elliptic flow [**Ollitrault, 1992; Teaney, Lauret and Shuryak, 2001**]
- ▶ Renormalized Polyakov loops in various representations [**Damgaard, 1987; Damgaard and Hasenbusch, 1994; Dumitru, Hatta, Lenaghan, Orginos and Pisarski, 2004; Gupta, Hübner and Kaczmarek, 2008**]
- ▶ Transport coefficients [**Meyer, 2007**]

Projects for the future - II


(in case 'plan A' fails ...)

- ▶ High-precision thermodynamics for $SU(N)$ theories in 3D (work in progress with Caselle, Castagnini, Feo and Gliozzi; see also [Bialas, Daniel, Morel and Petersson, 2008])

D=2+1 $SU(N)$ trace anomaly in the confined phase
(preliminary)



D=2+1 $SU(N)$ trace anomaly in the deconfined phase
(preliminary)



ETH

Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich