

## Experimental Study of the New 3.1 GeV Particle by $e^+e^-$ Collision at ADONE.

W. W. ASH (\*), D. C. CHENG (\*\*), B. ESPOSITO, F. FELICETTI, H. OGREN (\*\*\*),  
I. PERUZZI, M. PICCOLO, F. RONGA and G. SCHINA

*Laboratori Nazionali di Frascati del CNEN - Frascati (Roma)*

G. T. ZORN (\*,\*)

*Department of Physics, University of Maryland - College Park, Md.*

B. BARTOLI, B. COLUZZI, E. CUOMO, A. NIGRO, V. SILVESTRINI and F. VANOLI

*Istituto di Fisica dell'Università - Napoli*

*Istituto Nazionale di Fisica Nucleare - Sezione di Napoli*

D. BISELLO, A. MULACHIÉ, M. NIGRO, L. PESCARA and E. SCHIAVUTA

*Istituto di Fisica dell'Università - Padova*

*Istituto Nazionale di Fisica Nucleare - Sezione di Padova*

A. MARINI, P. MONACELLI, L. PAOLUZI, G. PIANO MORTARI,  
F. SEBASTIANI and L. TRASATTI

*Istituto di Fisica dell'Università - Roma*

*Istituto Nazionale di Fisica Nucleare - Sezione di Roma*

(ricevuto il 10 Dicembre 1974)

We present preliminary results of an investigation of electron-positron annihilation in the total c.m. energy range  $\sqrt{s} = (3.09 \div 3.112)$  GeV.

The experiment was performed at ADONE, the Frascati  $e^+e^-$  storage ring.

Evidence for the production of the new particle of mass 3.1 GeV, first observed by the MIT group at Brookhaven (1), and by the Berkeley-SLAC group at SPEAR (2) has already been reported by the ADONE groups (3).

---

(\*) Present address: SLAC, Stanford, Ca.

(\*\*) Present address: University of California, Santa Cruz, Ca.

(\*\*\*) Present address: CERN, Genève.

(\*,\*\*) Supported by U.S.A.E.C.

(1) J. J. AUBERT, U. BECKER, J. P. BIGGS, J. BURGER, M. CHEN, G. EVERHART, P. GOLDHAGEN, J. LEONG, T. MC CORRISTON, T. G. RHOADES, M. ROHDE, S. C. C. TING, SAU LAN WU and Y. Y. LEE: *Phys. Rev. Lett.*, **33**, 1404 (1974).

(2) J. E. AUGUSTIN, A. M. BOYARSKI, M. BREIDENBACH, F. BULOS, J. T. DAKIN, G. J. FELDMAN, G. E.

The present letter concerns the study of the channels

- (1)  $e^+e^- \rightarrow e^+e^-$ ,
- (2)  $e^+e^- \rightarrow \mu^+\mu^-$ ,
- (3)  $e^+e^- \rightarrow \text{many hadrons}$ .

A very narrow resonant behaviour of the cross-section for these processes is confirmed. The total hadronic cross-section integrated over energy is  $\sigma_I = (4.7 \pm 1.6) \cdot 10^{-33} \text{ cm}^2 \text{ GeV}$ .




Fig. 1. - Vertical section of the experimental apparatus.  $C_1, C_1'$  are narrow-gap spark chambers;  $C_2, C_2'$  are wide-gap cylindrical spark chambers for momentum analysis;  $C_3, C_3'$  are thick plate spark chambers for particle identification. MWPC are multiwire proportional chambers;  $S_1, \dots, S_4$  are scintillation counters

FISCHER, D. FRYBERGER, G. HANSON, B. JEAN-MARIE, R. R. LARSEN, V. LÜTH, H. L. LYNCH, D. LYON, C. C. MOREHOUSE, J. M. PATERSON, M. L. PERL, B. RICHTER, P. RAPIDIS, R. F. SCHWITERS, W. M. TANENBAUM, F. VANNUCCI, G. S. ABRAMS, D. BRIGGS, W. CHINOWSKY, C. E. FRIEDBERG, G. GOLDBABER, R. J. HOLLEBEEK, J. A. KADYK, B. LULU, F. PIERRE, G. H. TRILLING, J. S. WHITAKER, J. WISS and J. E. ZIFSE: *Phys. Rev. Lett.*, **33**, 1406 (1974).

(<sup>2</sup>) C. BACCI, R. BALDINI-CELIO, M. BERNARDINI, G. CAPON, R. DEL FABBRO, M. GRILLI, E. IAROCCI, L. H. JONES, M. LOCCI, C. MENCUCCHINI, G. P. MURTAS, G. PENSO, G. SALVINI, M. SPINETTI, M. SPANO, B. STELLA, V. VALENTE, B. BARTOLI, D. BISELLO, B. ESPOSITO, F. FELICETTI, P. MONACELLI, M. NIGRO, L. PAOLUZI, I. PERUZZI, G. PIANO MORTARI, M. PICCOLO, F. RONGA, F. SEBASTIANI, L. TRASATTI, F. VANOLI, G. BARBARINO, G. BARBIELLINI, C. BEMPORAD, R. BIANCASTELLI, M. CASTELLANO, F. CEVENINI, M. CALVETTI, F. COSTANTINI, P. LARICCIA, P. PARASCANDOLO, S. PATRICELLI, E. SASSI, C. SPENCER, L. TORTORA, U. TROYA and S. VITALE: *Phys. Rev. Lett.*, **33**, 1408 (1974).

The MEA <sup>(4)</sup> experimental apparatus as shown in Fig. 1 consists of the following parts:

a) A large (2 m diameter  $\times$  2 m length) solenoid with aluminum coil mounted with its axis perpendicular to the direction of the  $e^+e^-$  beams. Two compensator magnets, covering about 50% of the beam path inside the solenoid, which reduce the magnetic field integrated along the beam path practically to zero.

b) Two 2-plane telescopes of multiwire proportional chambers (MWPC) above and below the beam path at the crossing region. These chambers have the sense wires parallel to the direction of the beams and are used both in the trigger and in the off-line geometrical reconstruction.

c) A system of optical spark chambers used for momentum analysis which comprises two narrow-gap and two cylindrical wide-gap spark chambers coaxial with the solenoid, shown in Fig. 1 as  $C_1$  and  $C_2$ , respectively. The electrodes of  $C_2$  are made of wires in order to reduce multiple scattering and to allow viewing the sparks through the electrodes themselves. This allows to achieve complete stereo view of the tracks. As a matter of fact we use two cameras aligned with the axis of the solenoid and viewing the apparatus from opposite sides.

d) A system of 12 cylindrical thick plate narrow-gap spark chambers  $C_3$  for the detection of showers and nuclear interactions, mounted outside the magnet coil. Their total thickness is  $\approx 6$  radiation lengths and  $\approx 1$  collision length. The solid angle covered by the apparatus for pointlike source is  $\Delta\Omega_c \approx 0.4 \cdot 4\pi (40^\circ \leq \theta \leq 140^\circ)$  for momentum analysis, and  $\Delta\Omega_y \approx 0.27 \cdot 4\pi$  for particle identification (as defined by chambers  $C_3$ ,  $C'_3$ ). The effect of the extended source is to lower  $\Delta\Omega_c$  to about  $0.09 \cdot 4\pi$  at  $\sqrt{s} = 3.1$  GeV.

e) A scintillation counter system ( $S_1$  to  $S_4$ ) used to trigger the apparatus. In the trigger, we require at least one particle with a minimum kinetic energy of about 130 MeV (if pion) going in the upper part of the apparatus ( $S_1S_2S_3S_4$ ) and one particle with about 110 MeV going in the lower part ( $S'_1S'_2S'_3$ ). For two-prong events in order to reduce machine background some configurations (tracks with small dip angle) are rejected.

Cosmic rays are rejected by means of time-of-flight selection over a path length of one meter ( $S_1$  to  $S_3$ ) by requiring a suitable phase relative to the accelerator radio-frequency ( $\Delta t_{RF}$ , see Fig. 2), and by requesting that the source point lies within  $\pm 5$  cm from the beam line. This last rejection is achieved by making use of a fast logic signal from MWPC. The residual counting rate from cosmic rays is 1/15 min.

For each trigger an on-line computer records on magnetic tape all relevant information. The selection of good events is made by using both pictures and computer-recorded information.

Events from the reaction  $e^+e^- \rightarrow \mu^+\mu^-$  are identified by using the following criteria: i) proper position of the source point; ii) correct timing with the beam-beam interaction; iii) correct time of flight in both sides of the apparatus between counters  $S_1S_2$  and  $S_3$  ( $S'_1S'_2$  and  $S'_3$ ); iv) momentum measurement; v) absence of interactions in the thick plate chambers  $C_3$  and  $C'_3$ . Cosmic-ray contamination is negligible. A study of nuclear interactions in  $C_3$  or  $C'_3$  of collinear events sets an upper limit of 3% on the contamination of hadronic collinear events on  $\mu$ -pair production events.

Events from the reaction  $e^+e^- \rightarrow e^+e^-$  must satisfy criteria from i) to iii) and in addition must show electromagnetic showers in  $C_3$  and  $C'_3$ .

<sup>(4)</sup> W. W. ASH, D. GROSSMAN, G. MATTHIAE, G. P. MURTAS, M. NIGRO, G. K. O'NEILL, G. SACERDOTI, R. SANTANGELO, D. SCANNICCHIO and E. SCHIAVUTA: LNF-69/2 (January 1969).

Multihadron events are required to present two noncollinear tracks or more than two tracks in the wide-gap chambers ( $C_2, C'_2$ ), at least one track in  $C_3$  or  $C'_3$ , and to satisfy criteria i) and ii).  $\gamma$ -rays with energy  $E \geq 100$  MeV are also seen as showers in  $C_3, C'_3$ .




Fig. 2. - Time separation  $\Delta t_{RF}$  between the occurrence of the event and a timing signal from a beam pick-up electrode

The experimental results are summarized in Table I. The integrated luminosity  $\mathcal{L}$  has been obtained from the measured rate of small-angle Bhabha scattering observed by the ADONE machine group in a different interaction region with a monitor whose calibration constant was obtained by comparing its rate outside the resonance with the rate of wide-angle Bhabha scattering measured in our apparatus.

In columns 3 to 9 we present the total number of collected multihadron events according to their detected configurations (*e.g.*,  $2C$  = two charged particles detected,  $2C + \geq 1\gamma$  = two charged particles +  $\geq 1$  gamma ray, ...). The corresponding numbers of collinear  $e^+e^-$  and  $\mu^+\mu^-$  events are given in columns 11 and 12.

The rates for processes (1) and (2) integrated over the acceptance of the experimental apparatus are shown in Fig. 3 and 4 as functions of  $\sqrt{s}$ .

In order to evaluate the total multihadron cross-section, a detailed knowledge of the produced final states is needed. At present we use only the information available on the detected charged multiplicity as determined by charge recognition, and on the

TABLE I. - Total number of collected events.

$\sqrt{s}$ (GeV)	Integrated luminosity ( $10^{33} \text{ cm}^{-2}$ )	Multihadron events		
		$N_{2C}$	$N_{2C+\geq 1\gamma}$	$N_{3C}$
(1)	(2)	(3)	(4)	(5)
3.09 ÷ 3.112	$11.8 \pm 1.2$	133	95	125

detected-photon multiplicity. The complete momentum analysis is in progress. The average detected charged multiplicity turns out to be  $\langle N_c \rangle = 3.3 \pm 0.2$ . Furthermore we can set lower limits on the production of final states with at least six and eight charged pions. The ratio of the number of events with a least six (eight) charged pions to the total number of detected multihadron events is found to be  $N_{\geq 6\text{ch}}/N_{\text{tot}} \geq 0.13 \pm 0.02$  ( $N_{\geq 8\text{ch}}/N_{\text{tot}} \geq 0.02 \pm 0.01$ ).




Fig. 3.

Fig. 3. -  $e^+e^- \rightarrow \mu^+\mu^-$  rate integrated over the acceptance of the experimental apparatus vs.  $\sqrt{s}$ .




Fig. 4.

Fig. 4. -  $e^+e^- \rightarrow e^+e^-$  rate integrated over the acceptance of the experimental apparatus vs.  $\sqrt{s}$ .

We have evaluated with a Monte Carlo calculation the detection efficiency for a number of initial pion states (with different charged and neutral multiplicities), leading to various observed configurations (*i.e.*  $2C$ ,  $2C + \geq 1\gamma$ ,  $3C$ ,  $3C + \geq 1\gamma$ , ...). Phase space momentum distribution (PS) has been assumed. The trigger efficiencies are found to be in the range  $0.06 \div 0.12$  for channels like  $e^+e^- \rightarrow 2\pi^+2\pi^-$ ,  $2\pi^+2\pi^-\pi^0$ ,  $2\pi^+2\pi^-2\pi^0$ ,  $2\pi^+2\pi^-3\pi^0$ ,  $3\pi^+3\pi^-\pi^0$ ,  $4\pi^+4\pi^-$ .

The experimental ratio  $R_\gamma$  of the number of events with at least one photon detected to the total number of multihadron events is found to be  $R_\gamma = 0.33 \pm 0.02$ . This value of  $R_\gamma$  and the predictions of our Monte Carlo calculation indicate that the production of final states with one  $\pi^0$  is favoured with respect to final states in which two or more  $\pi^0$ 's are produced.

The observed angular distribution for configurations in which three and four charged particles are detected is somewhat different from the one calculated with the Monte Carlo program. This suggests the existence of correlation effects among the produced particles. In spite of this moderate inconsistency, we have deduced from our PS Monte Carlo calculation an average detection efficiency  $\varepsilon = 0.09 \pm 0.03$  for multihadron events. In Fig. 5 the  $e^+e^- \rightarrow$  multihadron rate integrated over the acceptance of the experimental apparatus is plotted vs.  $\sqrt{s}$ .

$\sqrt{s_{30+ \geq 1\gamma}}$	$N_{4C}$	$N_{40+ \geq 1\gamma}$	$N_{5C}$	Total number of multihadrons	Collinear events	
					$e^+e^-$	$\mu^+\mu^-$
8)	(7)	(8)	(9)	(10)	(11)	(12)
9	73	32	27	554	145	38

The mass resolution is determined by the energy spread of the beams. In ADONE, at  $\sqrt{s} \approx 3.1$  GeV, this spread is  $\Gamma_{2E} = 3$  MeV (FWHM) <sup>(5)</sup>.

The total multihadron cross-section  $\sigma_h$  (Fig. 6) integrated over the energy interval  $\sqrt{s} = (3.099 \div 3.107)$  GeV is found to be  $\sigma_I = (4.7 \pm 1.6) \cdot 10^{-33} \text{ cm}^2 \text{ GeV}$ . The error quoted includes a 30% estimated uncertainty due to the detection efficiency calculation.




Fig. 5.

Fig. 5. —  $e^+e^- \rightarrow$  many hadron rate integrated over the acceptance of the experimental apparatus vs.  $\sqrt{s}$ .




Fig. 6.

Fig. 6. — Total cross-section  $\sigma_h$  for the reaction  $e^+e^- \rightarrow$  many hadrons vs.  $\sqrt{s}$ . The errors are statistical only.

As far as the nature of this particle is concerned, it is difficult at present to make any definite statement. In particular, a detailed study of the  $e^+e^- \rightarrow \mu^+\mu^-$  <sup>(6)</sup> channel may be relevant for the understanding of its properties.

\* \* \*

We wish to thank the ADONE machine group for the excellent performance of the accelerator, which made this experimental possible. The help of both the OPTA group of G. DI STEFANO and his co-workers and of the Mechanical Engineering group was invaluable in the design and setting up of the apparatus. We also thank the Film Developing Service, Mrs. R. MORANI from the Film Scanning Service, and all people from inside and outside the Laboratory who contributed to the success of this experiment.

We are grateful to G. ALTARELLI, C. BERNARDINI, N. CABIBBO, M. GRECO, L. MAIANI, G. PARISI and Y. SRIVASTAVA for many stimulating discussions and suggestions.

<sup>(5)</sup> M. BASSETTI: ADONE Internal Report, E-15 (November 1974).

<sup>(6)</sup> G. ALTARELLI, N. CABIBBO, L. MAIANI, G. PARISI and R. PETRONZIO: *Lett. Nuovo Cimento*, **11**, 609 (1974).