

# Quest for precision in hadronic cross sections at low energy : Monte Carlo tools vs. experimental data

## Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies

S. Actis<sup>36</sup>, A. Arbuzov<sup>8,41</sup>, G. Balossini<sup>30,31</sup>, P. Beltrame<sup>12</sup>, C. Bignamini<sup>30,31</sup>, R. Bonciani<sup>14</sup>, C. M. Carloni Calame<sup>33</sup>, V. Cherepanov<sup>25,26</sup>, M. Czakon<sup>1</sup>, H. Czyż<sup>18,42,45,46</sup>, A. Denig<sup>21</sup>, S. Eidelman<sup>25,26,44</sup>, G. V. Fedotovitch<sup>25,26,41</sup>, A. Ferroglia<sup>22</sup>, J. Gluza<sup>18</sup>, A. Grzelińska<sup>7</sup>, M. Gunia<sup>18</sup>, A. Hafner<sup>21</sup>, S. Jadach<sup>7</sup>, A. Kalinowski<sup>27</sup>, W. Kluge<sup>16</sup>, A. Korchin<sup>19</sup>, J. H. Kühn<sup>17</sup>, P. Mastrolia<sup>13</sup>, G. Montagna<sup>30,31,40,45</sup>, S. E. Müller<sup>21,42</sup>, F. Nguyen<sup>32,40</sup>, O. Nicrosini<sup>31</sup>, D. Nomura<sup>34,43</sup>, G. Pakhlova<sup>23</sup>, G. Pancheri<sup>10</sup>, A. Penin<sup>9</sup>, F. Piccinini<sup>31</sup>, W. Placzek<sup>7</sup>, T. Przedzinski<sup>5</sup>, E. Remiddi<sup>3,4</sup>, T. Riemann<sup>39</sup>, G. Rodrigo<sup>35</sup>, P. Roig<sup>24,35,10</sup>, O. Shekhovtsova<sup>10</sup>, C. P. Shen<sup>15</sup>, A. L. Sibidanov<sup>25</sup>, T. Teubner<sup>20,43</sup>, L. Trentadue<sup>28,29</sup>, G. Venanzoni<sup>10,45,46</sup>, J. J. van der Bij<sup>11</sup>, P. Wang<sup>2</sup>, B. F. L. Ward<sup>37</sup>, Z. Was<sup>7,44</sup>, M. Worek<sup>38,18</sup>, and C. Z. Yuan<sup>2</sup>

- <sup>1</sup> Institut für Theoretische Physik E, RWTH Aachen Universität, D-52056 Aachen, Germany
- <sup>2</sup> Institute of High Energy Physics, Beijing 10049, China
- <sup>3</sup> Dipartimento di Fisica dell'Università di Bologna, I-40126 Bologna, Italy
- <sup>4</sup> INFN, Sezione di Bologna, I-40126 Bologna, Italy
- <sup>5</sup> The Faculty of Physics, Astronomy and Applied Computer Science, Jagellonian University, Reymonta 4, 30-059 Cracow, Poland
- <sup>6</sup> Marian Smoluchowski Institute of Physics, Jagiellonian University, 30-059 Cracow, Poland
- <sup>7</sup> Institute of Nuclear Physics Polish Academy of Sciences, PL-31342 Cracow, Poland
- <sup>8</sup> Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Russia
- <sup>9</sup> Department of Physics, University of Alberta, Edmonton, AB T6G 2J1, Canada
- <sup>10</sup> Laboratori Nazionali di Frascati dell'INFN, Frascati, Italy
- <sup>11</sup> Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, D-79104 Freiburg, Germany
- <sup>12</sup> Physics Department, CERN, Geneva, Switzerland
- <sup>13</sup> CERN, Theory Department, CH-1211 Genève, Switzerland
- <sup>14</sup> Laboratoire de Physique Subatomique et de Cosmologie, Université Joseph Fourier/CNRS-IN2P3/INPG, F-38026 Grenoble, France
- <sup>15</sup> Univ. of Hawaii, USA
- <sup>16</sup> Institut für Experimentelle Kernphysik, Universität Karlsruhe, Karlsruhe, Germany
- <sup>17</sup> Institut für Theoretische Teilchenphysik, Universität Karlsruhe, D-76128 Karlsruhe, Germany.
- <sup>18</sup> Institute of Physics, University of Silesia, PL-40007 Katowice, Poland
- <sup>19</sup> Kharkov Inst. of Physics and Tech. (KFTI), National Science Center; 1 Akademicheskaya Str.; UA-61108, Kharkov, Ukraine
- <sup>20</sup> Department of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, U.K.
- <sup>21</sup> Institut für Kernphysik, Johannes Gutenberg - Universität Mainz, Mainz, Germany
- <sup>22</sup> Institut für Physik (THEP), Johannes Gutenberg-Universität, D-55099 Mainz, Germany
- <sup>23</sup> Institute of Theoretical and Experimental Physics, Moscow, Russia
- <sup>24</sup> Techn. Univ. Muenchen, Physik Dept. T30; James Franck Strasse; D-85747 Garching, Germany
- <sup>25</sup> Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia
- <sup>26</sup> Novosibirsk State University, 630090 Novosibirsk, Russia
- <sup>27</sup> LLR-Ecole Polytechnique, 91128 Palaiseau, France
- <sup>28</sup> Dipartimento di Fisica, Università di Parma, I-43100 Parma, Italy
- <sup>29</sup> INFN, Gruppo Collegato di Parma, I-43100 Parma, Italy
- <sup>30</sup> Dipartimento di Fisica Nucleare e Teorica, Università di Pavia
- <sup>31</sup> INFN, Sezione di Pavia, I-27100 Pavia, Italy
- <sup>32</sup> Dipartimento di Fisica dell'Università "Roma Tre" and INFN Sezione di Roma Tre, Roma, Italy
- <sup>33</sup> School of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, UK
- <sup>34</sup> Theory Group, KEK, Tsukuba, Ibaraki 305-0801, Japan
- <sup>35</sup> Instituto de Fisica Corpuscular (IFIC), Centro mixto UVEG/CSIC; Edificio Institutos de Investigacion, Apartado de Correos 22085; E-46071 Valencia, Espanya
- <sup>36</sup> Paul Scherrer Institut, Wuerenlingen and Villigen, CH-5232 Villigen PSI, Switzerland
- <sup>37</sup> Department of Physics, Baylor University, Waco, Texas 76798-7316, USA

<sup>38</sup> Fachbereich C, Bergische Universität Wuppertal, D-42097 Wuppertal, Germany

<sup>39</sup> Deutsches Elektronen-Synchrotron, DESY, D-15738 Zeuthen, Germany

<sup>40</sup> Section 2 conveners

<sup>41</sup> Section 3 conveners

<sup>42</sup> Section 4 conveners

<sup>43</sup> Section 6 conveners

<sup>44</sup> Section 5 conveners

<sup>45</sup> Working group convener

<sup>46</sup> Correspondings authors: henryk.czyz@us.edu.pl, guido.montagna@pv.infn.it, graziano.venanzoni@lnf.infn.it

Received: date / Revised version: date

**Abstract.** We present last years achievements of the experimental and theoretical groups working on hadronic cross section measurements and tau physics and sketch the prospects in this fields for next few years to come. We describe the status of the Monte Carlo event generators used by the experimental collaborations at  $e^+e^-$  machines with energies up to the energies of B-factories and for analyses of tau decays, and give an overview of the experimental results concerning hadronic cross section measurement and tau decays. We cover experimental results and Monte Carlo generators concerning luminosity measurements, measurements of hadronic cross sections using scan and radiative return method, and tau decays. The accuracies of the Monte Carlo generators and their tests are overviewed. The available radiative corrections both included and not included in the codes are discussed. A special section is devoted to the vacuum polarisation contributions.

**PACS.** PACS-key describing text of that key – PACS-key describing text of that key

## Contents

1	Introduction . . . . .	2
2	Luminosity . . . . .	4
3	R measurement from energy scan . . . . .	35
4	Radiative return . . . . .	44
5	Tau decays . . . . .	72
6	Vacuum polarization . . . . .	82
7	Summary . . . . .	88

## 1 Introduction

The systematic comparison of the Standard Model (SM) predictions with precise experimental data served in the last few decades as an invaluable tool to test the theory at the quantum level. It has also provided stringent constraints on many “new physics” scenarios. The (so far) remarkable agreement between the measurements of the electroweak observables and their SM predictions is a striking experimental confirmation of the theory, even if there are a few observables where the agreement is not so satisfactory. On the other hand, the Higgs boson has not yet been observed, and there are strong theoretical arguments hinting at the presence of physics beyond the SM. Future colliders, like the upcoming LHC or an  $e^+e^-$  International Linear Collider (ILC), will hopefully answer many questions, offering at the same time great physics potential and a new challenge to provide even more precise theoretical predictions.

Precision tests of the Standard Model require an appropriate inclusion of higher order effects and the knowledge of very precise input parameters. One of the basic input parameters is the fine-structure constant  $\alpha$  which

depends logarithmically on the energy scale. It has been determined at zero momentum transfer with an impressive accuracy of 0.37 parts per billion (ppb) [1] from the measurement of the anomalous electron magnetic moment [1], relying on the validity of perturbative QED. At the  $M_Z$  scale the shift of the fine-structure constant involves low energy non-perturbative hadronic effects, which cause a dramatic reduction of accuracy, by several orders of magnitude [2, 3, 4, 5]. These low energy strong-interaction effects can be computed from hadronic  $e^+e^-$  annihilation data via the dispersion relation [6]:

$$\Delta\alpha_{\text{had}}^{(5)}(M_Z^2) = - \left( \frac{\alpha M_Z^2}{3\pi} \right) \text{Re} \int_{m_\pi^2}^{\infty} ds \frac{R(s)}{s(s - M_Z^2 - i\epsilon)}, \quad (1)$$

where  $R(s) = \sigma^{(0)}(s)/(4\pi\alpha^2/3s)$  and  $\sigma^{(0)}(s)$  is the total cross section for  $e^+e^-$  annihilation into hadronic state.

The current accuracy of this dispersive integral is of the order of 1%, dominated by the error of the hadronic cross section measurement in the energy region below few GeV [7, 8, 9, 10, 2, 3, 4, 5, 11, 12, 13, 14]. This error represents the largest source to the uncertainty of the effective electromagnetic coupling constant, (recent evaluations give  $\Delta\alpha(M_Z^2) = 22 \times 10^{-5}$  [5, 14]), becoming a limiting factor on the effective electroweak mixing parameter  $\sin^2\theta_f$ , and in turn on the prediction of the Higgs mass [15]. Table 1 (from Ref. [2]) shows that an uncertainty  $\delta\Delta\alpha_{\text{had}}^{(5)} \sim 5 \times 10^{-5}$ , needed for precision physics at a future linear collider, requires the measurement of the hadronic cross section with a precision of  $O(1\%)$  from threshold up to the  $\Upsilon$  peak.

Like the effective fine-structure constant at the scale  $M_Z$ , the SM determination of the anomalous magnetic moment of the muon  $a_\mu$  is presently limited by the evaluation

$\delta\Delta\alpha_{\text{had}}^{(5)} \times 10^5$	$\delta(\sin^2\theta_{\text{eff}}^{\text{lept}}) \times 10^5$	Request on $R$
22	7.9	Present
7	2.5	$\delta R/R \sim 1\%$ up to $J/\psi$
5	1.8	$\delta R/R \sim 1\%$ up to $\Upsilon$

**Table 1.** [2] Values of the uncertainties  $\delta\Delta\alpha_{\text{had}}^{(5)}$  (first column) and the errors induced by these uncertainties on the theoretical SM prediction for  $\sin^2\theta_{\text{eff}}^{\text{lept}}$  (second column). The third column indicates the corresponding requirements for the  $R$  measurement.

of the hadronic vacuum polarisation effects, which cannot be computed perturbatively at low energies. However, using analyticity and unitarity, it was shown long ago that this term can be computed from hadronic  $e^+e^-$  annihilation data via the dispersive integral [16]:

$$\begin{aligned} a_\mu^{\text{HLO}} &= (1/4\pi^3) \int_{m_\pi^2}^{\infty} ds K(s) \sigma^{(0)}(s) \\ &= (\alpha^2/3\pi^2) \int_{m_\pi^2}^{\infty} ds K(s) R(s)/s. \end{aligned} \quad (2)$$

The kernel function  $K(s)$  decreases monotonically with increasing  $s$ . This integral is similar to the one entering the evaluation of the hadronic contribution  $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$  in Eq. (1). Here, however, the weight function in the integrand gives a stronger weight to low-energy data. A recent compilation of  $e^+e^-$  data gives [17]:

$$a_\mu^{\text{had}} = (695.5 \pm 4.1) \times 10^{-10}. \quad (3)$$

Similar values are obtained by other groups [14, 18, 19, 20].

By adding this contribution to the rest of the SM contributions, a recent update of the SM prediction of  $a_\mu$ , which uses the result in [21] gives [17, 22]:

$a_\mu^{\text{SM}} = 116591834(49) \times 10^{-11}$ . The difference with the experimental average [23],  $a_\mu^{\text{exp}} = 116592080(63) \times 10^{-11}$  is then  $\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = +246(80) \times 10^{-11}$ , i.e., 3.1 standard deviations (all errors were added in quadrature). Slightly higher discrepancies are obtained in Refs. [14, 19, 20]. As in the case of  $\alpha(M_Z^2)$ , the uncertainty on the theoretical evaluation of  $a_\mu$  is still dominated by the hadronic contribution at low energies, and a reduction of the uncertainty is necessary in order to match the precision of the proposed experiments on the measurement of  $a_\mu$  at FNAL [24] and J-PARK [25].

The precise determination of the hadronic cross sections (accuracy  $\leq 1\%$ ) requires a deep control of higher order effects like Radiative Corrections (RC) and the non-perturbative hadronic contribution to the running of  $\alpha$  (vacuum polarisation, VP) into Monte Carlo (MC) programs. Particularly in the last years, the increasing precision reached on the experimental side at the  $e^+e^-$  colliders (VEPP-2M, DAΦNE, BEPC, PEP-II and KEKB) led to the development of dedicated high precision theoretical tools: BabaYaga (and its successor BabaYaga@NLO) for the measurement of the luminosity, MCGPJ for the

simulation of the exclusive QED channels, and PHOKHARA for the simulation of the process with Initial State Radiation (ISR)  $e^+e^- \rightarrow \text{hadrons} + \gamma$ , are examples of MC generators which include NLO corrections with per mille accuracy. In parallel to these efforts, well-tested codes such as BHWIDE (developed for LEP/SLC colliders) were adopted.

Theoretical accuracies of these generators were estimated, whenever possible, by evaluating higher order missing contributions (from this point of view, the great progress in the calculation of two-loop corrections to the Bhabha scattering cross section was essential to establish the theoretical accuracy of the existing generators for the luminosity measurement). Usually only analytical or semi-analytical estimates of missing terms exist which don't take into account realistic experimental cuts. In addition MC event generators include different parametrisation for the VP, which affect the prediction (and the precision) of the cross sections, and RC are usually implemented with different formulations.

By these arguments it becomes evident how important are the comparisons of MC generators for a common set of input parameters and experimental cuts. Such *tuned* comparisons, which started in the LEP era, are a key step in the validation of the generators, since it allows to check that the different details entering the complex structure of the generators are under control, and free of possible bugs. This was the main motivation for the “Working Group on Radiative Corrections and Monte Carlo tools”, (Radio Montecarlo), which was formed 3 years ago bringing together experts (theorists and experimentalists) working in the field of low energy  $e^+e^-$  physics and partly also the  $\tau$  community.

In addition to tuned comparisons, the technical details of the MC generators, the recent progress (like new calculations), and the open issues were also discussed in regular meetings.

This report is a summary of all these efforts: it provides a self-contained and up-to-date description of the progress which occurred in the last years towards precision hadronic physics at low energies, together with new original results like comparisons and estimates of high order effects (e.g. pion pair correction to the Bhabha process) in presence of realistic experimental cuts.

The report is divided into five sections: the first three are devoted to the status of the MC tools for Luminosity R-scan, and Initial State Radiation (ISR).

Tau spectral function in hadronic decays are also used to estimate  $a_\mu^{\text{had}}$ , since they can be related to  $e^+e^-$  annihilation cross section via isospin symmetry [26, 27, 28, 29]. The substantial difference between the  $e^+e^-$ - and  $\tau$ -based determinations of  $a_\mu^{\text{had}}$ , even if isospin violation corrections are taken into account, shows that further common theoretical and experimental efforts are necessary to understand this phenomenon. In Section 4 the experimental status and MC tools for tau decays are discussed. The recent improvements of the generators TAUOLA and PHOTOS are discussed and perspectives for further developments are sketched.

Section 5 focuses on the theoretical treatment of non-perturbative effects of light quarks to the Vacuum Polarisation, which is a key ingredient for the high precision determination of the hadronic cross section.

## 2 Luminosity

The present Section addresses the most important experimental and theoretical issues involved in the precision determination of the luminosity at meson factories. The luminosity is the key ingredient underlying all the measurements and studies of the physics processes discussed in the other Sections. Particular emphasis is put on the theoretical accuracy inherent to the event generators used in the experimental analysis, in comparison with the most advanced perturbative calculations and experimental precision requirements. The effort done during the activity of the working group to perform tuned comparisons between the predictions of the most accurate programs is described in detail. New calculations, leading to an update of the theoretical error associated to the prediction of the luminosity cross section, are also presented. The aim of the Section is to provide a self-contained and up-to-date description of the progress occurred during the last few years towards high-precision luminosity monitoring at flavour factories, as well as of the still open issues necessary for future advances.

The structure of the Section is as follows. After an introduction on the motivation for precision luminosity measurements at meson factories (Section 2.1), the leading-order (LO) cross sections of the two QED processes of major interest, i.e. Bhabha scattering and photon pair production, are presented in Section 2.2, together with the formulae for the next-to-leading-order (NLO) photonic corrections to the above processes. The remarkable progress on the calculation of next-to-next-leading-order (NNLO) QED corrections to the Bhabha cross section, as occurred in the last few years, is reviewed in Section 2.3. In particular, this Section presents new exact results on lepton and hadron pair corrections, taking into account realistic event selection criteria. Section 2.4 is devoted to the description of the theoretical methods used in the Monte Carlo (MC) generators for the simulation of multiple photon radiation. The matching of such contributions with NLO corrections is also described in Section 2.4. The main features of the MC programs used by the experimental collaborations are summarized in Section 2.5. Numerical results for the radiative corrections implemented into the MC generators are shown in Section 2.6 for both the Bhabha process and two-photon production. The tuned comparisons between the predictions of the most precise generators are presented and discussed in detail in Section 2.7, considering the Bhabha process at different centre-of-mass (c.m.) energies and with realistic experimental cuts. The theoretical accuracy presently reached by the luminosity tools is addressed in Section 2.8, where the most important sources of uncertainty are discussed quantitatively. The estimate of the total error affecting the calculation of the Bhabha cross section is

given, as main conclusion of the present work, in Section 2.9, updating and making more robust results available in the literature. Some still open issues are drawn in Section 2.9 as well.

### 2.1 Motivation

The luminosity of a collider is the normalization constant between the event rate and the cross section of a given process. For an accurate measurement of the cross section of an electron-positron ( $e^+e^-$ ) annihilation process, the precise knowledge of the collider luminosity is mandatory.

The luminosity depends on three factors: beam-beam crossing frequency, beam currents and the beam overlap area in the crossing region. However, the last quantity is difficult to determine accurately from the collider optics. Thus, experiments prefer to determine the luminosity by the counting rate of well selected events whose cross section is known with good accuracy, using the formula [30]

$$\int \mathcal{L} dt = \frac{N}{\epsilon \sigma}, \quad (4)$$


where  $N$  is the number of events of the chosen reference process,  $\epsilon$  the experimental selection efficiency and  $\sigma$  the theoretical cross section of the reference process. Therefore, the total luminosity error will be given by the sum in quadrature of the fractional experimental and theoretical uncertainties.

Since the advent of low luminosity  $e^+e^-$  colliders, a great effort was devoted to obtain good precision in the cross section of electromagnetic processes, extending the pioneering work of the earlier days [6]. At the  $e^+e^-$  colliders, operating in the c.m. energy range  $1 \text{ GeV} < \sqrt{s} < 3 \text{ GeV}$ , such as ACO at Orsay, VEPP-II at Novosibirsk and Adone at Frascati, the luminosity measurement was based on Bhabha scattering [31,32] with final-state electrons and positrons detected at small angles, or single and double bremsstrahlung processes [33], thanks to their high statistics. The electromagnetic cross sections scale as  $1/s$ , while elastic  $e^+e^-$  scattering has a steep dependence on the polar angle,  $\sim 1/\theta^3$ , thus providing high rate for small values of  $\theta$ .

Also at high-energy accelerators running in the '90s around the  $Z$  pole to perform precision tests of the Standard Model (SM), as LEP at CERN and SLC at Stanford, the experiments used small-angle Bhabha scattering events as a luminosity monitoring process. Indeed, for the very forward angular acceptances considered by LEP/SLC collaborations, the Bhabha process is dominated by the electromagnetic interaction and, therefore, calculable, at least in principle, with very high accuracy. At the end of LEP and SLC operation, a total (experimental plus theoretical) precision of one per mille (or better) was achieved [34,35,36,37,38,39,40], thanks to the work of different theoretical groups and the excellent performance of precision luminometers.

At current low- and intermediate-energy high-luminosity meson factories, the small polar angle region is dif-





**Fig. 1.** Comparison between large-angle Bhabha KLOE data (points) and MC (histogram) distributions for the  $e^\pm$  polar angle  $\theta$  (left) and for the acollinearity,  $\zeta = |\theta_{e^+} + \theta_{e^-} - 180^\circ|$  (right), where the flight direction of the  $e^\pm$  is given by the position of clusters in the calorimeter. In each case, MC and data histograms are normalized to unity. From [30].

difficult to access for the presence of the low-beta insertions close to the beam crossing region, while wide-angle Bhabha scattering produces a large counting rate and can be exploited for a precise measurement of the luminosity.

Therefore, also in this latter case of  $e^\pm$  scattered at large angles, e.g. larger than  $55^\circ$  for the KLOE experiment [30] running at DAΦNE, in Frascati and larger than  $40^\circ$  for the CLEO-c experiment [41] running at CESR, Cornell, the main advantages of Bhabha scattering are preserved:

1. large statistics. For example at DAΦNE, a statistical error  $\delta\mathcal{L}/\mathcal{L} \sim 0.3\%$  is reached in about two hours of data taking, even at the lowest luminosities;
2. high accuracy for the calculated cross section;
3. clean event topology of the signal and small amount of background.


In Eq. (4) the cross section is usually evaluated by inserting event generators, which include radiative corrections at a high level of precision, into the MC code simulating the detector response. The code has to be developed to reproduce the detector performance (geometrical acceptance, reconstruction efficiency and resolution of the measured quantities) to a high level of confidence.

In most cases the major sources of the systematic errors of the luminosity measurement are differences of efficiencies and resolutions between data and MC.

In the case of KLOE the largest experimental error of the luminosity measurement is due to a different polar angle resolution between data and MC which is observed at the edges of the accepted interval for Bhabha scattering events. Fig. 1 shows a comparison between large angle Bhabha KLOE data and MC, at left for the polar angle and at right for the acollinearity  $\zeta = |\theta_{e^+} + \theta_{e^-} - 180^\circ|$ . One observes a very good agreement between data and

MC, but also the differences (of about 0.3 %) at the sharp interval edges. The analysis cut,  $\zeta < 9^\circ$ , applied to the acollinearity distribution is very far from the bulk of the distribution and does not introduce noteworthy systematic errors. Also in the CLEO-c luminosity measurement with Bhabha scattering events, the detector modeling is the main source of experimental error. In particular, uncertainties include those due to finding and reconstruction of the electron shower, in part due to the nature of the electron shower, as well as the steep  $e^\pm$  polar angle distribution.

The luminosity measured with Bhabha scattering events is often checked by using other QED processes, such as  $e^+e^- \rightarrow \mu^+\mu^-$  or  $e^+e^- \rightarrow \gamma\gamma$ . In KLOE, the luminosity measured with  $e^+e^- \rightarrow \gamma\gamma$  events differs by 0.3% with respect to Bhabha events. In CLEO-c,  $e^+e^- \rightarrow \mu^+\mu^-$  events are also used and the luminosity determined from  $\gamma\gamma$  ( $\mu^+\mu^-$ ) is found to be 2.1% (0.6%) larger than that from Bhabha events. Fig. 2 shows the CLEO-c data for the polar angle distributions of all three processes, compared with the corresponding MC predictions. The three QED processes are also used by the BaBar experiment at the PEP-II collider, Stanford, yielding a luminosity determination with an error of about 1% [42]. Large-angle Bhabha scattering is the normalization process adopted by the CMD-2 and SND collaborations at VEPP-2M, Novosibirsk, while both BES at BEPC, Beijing and Belle at KEKB, Tsukuba measure luminosity using the processes  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow \gamma\gamma$  with the final-state particles detected at wide polar angles and an experimental accuracy of a few per cent. However, the BES-III aims at reaching a few per mille error in the luminosity measurement in the near future [43].




**Fig. 2.** Distributions of CLEO-c  $\sqrt{s} = 3.774$  GeV data (circles) and MC simulations (histogram) for the polar angle of the positive lepton (upper two plots) in  $e^+e^-$  and  $\mu^+\mu^-$  events, and the mean value of  $|\cos\theta_\gamma|$  of the two photons in  $\gamma\gamma$  events. MC histograms are normalized to the number of data events. From [41].

The need of precision, namely better than 1%, and possibly redundant measurements of the collider luminosity is of utmost importance to perform accurate measurements of the  $e^+e^- \rightarrow \text{hadrons}$  cross sections, which are the key ingredient for evaluating the hadronic contribution to the running of the electromagnetic coupling constant  $\alpha$  and the muon anomaly  $g - 2$ .

## 2.2 LO cross sections and NLO corrections

As remarked in Section 2.1, the processes of interest for luminosity measurement at meson factories are Bhabha scattering and electron-positron annihilation into two photons and muon pairs. Here we present the LO formulae of the cross section of the processes  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow \gamma\gamma$ , as well as the QED corrections to their cross sections in the NLO approximation of the perturbation theory. The reaction  $e^+e^- \rightarrow \mu^+\mu^-$  is discussed in Section 3.



**Fig. 3.** LO Feynman diagrams for the Bhabha process in QED, corresponding to  $s$ -channel annihilation and  $t$ -channel scattering.

### 2.2.1 LO cross sections

For the Bhabha scattering process

$$e^-(p_-) + e^+(p_+) \rightarrow e^-(p'_-) + e^+(p'_+) \quad (5)$$

at the Born level with simple one-photon exchange (see Fig. 3) the differential cross section reads

$$\frac{d\sigma_0^{\text{Bhabha}}}{d\Omega_-} = \frac{\alpha^2}{4s} \left( \frac{3+c^2}{1-c} \right)^2 + \mathcal{O}\left(\frac{m_e^2}{s}\right), \quad (6)$$

where

$$s = (p_- + p_+)^2, \quad c = \cos\theta_-. \quad (7)$$

The angle  $\theta_-$  is defined between the initial and final electron three-momenta,  $d\Omega_- = d\phi_- d\cos\theta_-$ , and  $\phi_-$  is the azimuthal angle of the outgoing electron. The small mass correction terms suppressed by the ratio  $m_e^2/s$  are negligible for the energy range and the angular acceptances which are of interest here.

At meson factories the Bhabha scattering cross section is largely dominated by  $t$ -channel photon exchange, followed by  $s$ - $t$  interference and  $s$ -channel annihilation. Furthermore,  $Z$ -boson exchange contributions and other electroweak effects are suppressed at least by a factor  $s/M_Z^2$ . In particular, for large-angle Bhabha scattering with a c.m. energy  $\sqrt{s} = 1$  GeV the  $Z$  boson contribution amounts to about  $-1 \times 10^{-5}$ . For  $\sqrt{s} = 3$  GeV it amounts to  $-1 \times 10^{-4}$  and  $-1 \times 10^{-3}$  for  $\sqrt{s} = 10$  GeV. So only at  $B$ -factories the electroweak effects should be taken into account at the tree level, when aiming at a per mille precision level.

The LO differential cross section of the two-photon annihilation channel (see Fig. 4)

$$e^+(p_+) + e^-(p_-) \rightarrow \gamma(q_1) + \gamma(q_2)$$

can be obtained by a crossing relation from the Compton scattering cross section computed by Brown and Feynman [44]. It reads as follows

$$\frac{d\sigma_0^{\gamma\gamma}}{d\Omega_1} = \frac{\alpha^2}{s} \left( \frac{1+c_1^2}{1-c_1^2} \right) + \mathcal{O}\left(\frac{m_e^2}{s}\right), \quad (8)$$

where  $d\Omega_1$  denotes the differential solid angle of the first photon. It is assumed that both final photons are registered in a detector, and that their polar angles with respect to the initial beam directions are not small ( $\theta_{1,2} \gg m_e/E$ , where  $E$  is the beam energy).




Fig. 4. LO Feynman diagrams for the process  $e^+e^- \rightarrow \gamma\gamma$ .

### 2.2.2 NLO corrections

The complete set of NLO radiative corrections, emerging at  $\mathcal{O}(\alpha)$  of the perturbation theory, to Bhabha scattering and two-photon annihilation can be split into gauge-invariant subsets: QED corrections, due to emission of real photons off the charged leptons and exchange of virtual photons between them, and purely weak contributions arising from the electroweak sector of the SM.

The complete  $\mathcal{O}(\alpha)$  QED corrections to Bhabha scattering are known since a long time [45, 46]. The first complete NLO prediction in the electroweak SM was performed in [47], followed by [48] and several others. At the NNLO, the leading virtual weak corrections from the top quark were derived first in [49] and are available in the fitting programs ZFITTER [50, 51] and TOPAZ0 [52, 53, 54] extensively used by the experimentalists for the extraction of the electroweak parameters at LEP/SLC. The weak NNLO corrections in the Standard Model are also known for the  $\rho$ -parameter [55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71] and the weak mixing angle [72, 73, 74, 75, 76, 77], as well as corrections from Sudakov logarithms [78, 79, 80, 81, 82, 83, 84, 85]. Both NLO and NNLO weak effects are negligible at low energies and are not implemented yet in numerical packages for Bhabha scattering at meson factories. In pure QED, the situation is considerably different due to a remarkable progress made on NNLO corrections in recent years, as emphasized and discussed in detail in Section 2.3.

As usual, the photonic corrections can be split into two parts according to their kinematics. The first part preserves the Born-like kinematics and contains the effects due to one-loop amplitudes (virtual corrections) and single soft-photon emission. Examples of Feynman diagrams giving rise to such corrections are represented in Fig. 5. The energy of a soft photon is assumed not to exceed an energy  $\Delta E$ , where  $E$  is the beam energy and the auxiliary parameter  $\Delta \ll 1$  should be chosen in such a way that the validity of soft-photon approximation is guaranteed. The second contribution is due to hard photon emission, i.e. to single bremsstrahlung with photon energy above  $\Delta E$  and corresponds to the radiative process  $e^+e^- \rightarrow e^+e^-\gamma$ .

Following [86, 87], the soft plus virtual (SV) correction can be cast into the form

$$\frac{d\sigma_{B+S+V}^{\text{Bhabha}}}{d\Omega_-} = \frac{d\sigma_0^{\text{Bhabha}}}{d\Omega_-} \left\{ 1 + \frac{2\alpha}{\pi}(L-1) \left[ 2 \ln \Delta + \frac{3}{2} \right] - \frac{8\alpha}{\pi} \ln(\text{ctg} \frac{\theta}{2}) \ln \Delta + \frac{\alpha}{\pi} K_{SV}^{\text{Bhabha}} \right\}, \quad (9)$$

where the factor  $K_{SV}^{\text{Bhabha}}$  is given by

$$K_{SV}^{\text{Bhabha}} = -1 - 2\text{Li}_2(\sin^2 \frac{\theta}{2}) + 2\text{Li}_2(\cos^2 \frac{\theta}{2}) + \frac{1}{(3+c^2)^2} \left[ \frac{\pi^2}{3}(2c^4 - 3c^3 - 15c) + 2(2c^4 - 3c^3 + 9c^2 + 3c + 21) \ln^2(\sin \frac{\theta}{2}) - 4(c^4 + c^2 - 2c) \ln^2(\cos \frac{\theta}{2}) - 4(c^3 + 4c^2 + 5c + 6) \ln^2(\text{tg} \frac{\theta}{2}) + 2(c^3 - 3c^2 + 7c - 5) \ln(\cos \frac{\theta}{2}) + 2(3c^3 + 9c^2 + 5c + 31) \ln(\sin \frac{\theta}{2}) \right], \quad (10)$$

and depends on the scattering angle, because of the contribution due to initial-final-state interference and box diagrams (see Fig. 6). It is worth noticing that the SV correction contains a leading logarithmic (LL) part enhanced by the collinear logarithm  $L = \ln(s/m_e^2)$ . Among the virtual corrections there is also a numerically important effect due to vacuum polarization in the photon propagator. Its contribution is omitted in Eq. (10) but can taken into account in the standard way by insertion of the resummed vacuum polarization operators in the photon propagators of the Born-level Bhabha amplitudes.

The differential cross section of the single hard bremsstrahlung process

$$e^+(p_+) + e^-(p_-) \rightarrow e^+(p'_+) + e^-(p'_-) + \gamma(k)$$

for scattering angles being large compared with  $m_e/E$  reads

$$d\sigma_{\text{hard}}^{\text{Bhabha}} = \frac{\alpha^3}{2\pi^2 s} R_{e\bar{e}\gamma} d\Gamma_{e\bar{e}\gamma}, \quad (11)$$

$$d\Gamma_{e\bar{e}\gamma} = \frac{d^3 p'_+ d^3 p'_- d^3 k}{\varepsilon'_+ \varepsilon'_- k^0} \delta^{(4)}(p_+ + p_- - p'_+ - p'_- - k),$$

$$R_{e\bar{e}\gamma} = \frac{WT}{4} - \frac{m_e^2}{(\chi'_+)^2} \left( \frac{s}{t} + \frac{t}{s} + 1 \right)^2 - \frac{m_e^2}{(\chi'_-)^2} \left( \frac{s}{t_1} + \frac{t_1}{s} + 1 \right)^2 - \frac{m_e^2}{\chi_+^2} \left( \frac{s_1}{t} + \frac{t}{s_1} + 1 \right)^2 - \frac{m_e^2}{\chi_-^2} \left( \frac{s_1}{t_1} + \frac{t_1}{s_1} + 1 \right)^2,$$

where

$$W = \frac{s}{\chi_+ \chi_-} + \frac{s_1}{\chi'_+ \chi'_-} - \frac{t_1}{\chi'_+ \chi_+} - \frac{t}{\chi'_- \chi_-} + \frac{u}{\chi'_+ \chi_-} + \frac{u_1}{\chi'_- \chi_+},$$

$$T = \frac{ss_1(s^2 + s_1^2) + tt_1(t^2 + t_1^2) + uu_1(u^2 + u_1^2)}{ss_1 tt_1},$$


and the invariants are defined as

$$s_1 = 2p'_- p'_+, \quad t = -2p_- p'_-, \quad t_1 = -2p_+ p'_+,$$


$$u = -2p_- p'_+, \quad u_1 = -2p_+ p'_-, \quad \chi_{\pm} = kp_{\pm}, \quad \chi'_{\pm} = kp'_{\pm}.$$

NLO QED radiative corrections to the two-photon annihilation channel were obtained in [88, 89, 90, 91], while weak corrections were computed in [92].





**Fig. 5.** Examples of Feynman diagrams for real and virtual NLO QED initial-state corrections to the  $s$ -channel contribution to the Bhabha process.



**Fig. 6.** Feynman diagrams for NLO QED box corrections to the  $s$ -channel contribution to the Bhabha process.

In the one-loop approximation the part of the differential cross section with the Born-like kinematics reads

$$\begin{aligned} \frac{d\sigma_{B+S+V}^{\gamma\gamma}}{d\Omega_1} &= \frac{d\sigma_0^{\gamma\gamma}}{d\Omega_1} \left\{ 1 + \frac{\alpha}{\pi} \left[ (L-1) \left( 2 \ln \Delta + \frac{3}{2} \right) \right. \right. \\ &\quad \left. \left. + K_{SV}^{\gamma\gamma} \right] \right\}, \\ K_{SV}^{\gamma\gamma} &= \frac{\pi^2}{3} + \frac{1-c_1^2}{2(1+c_1^2)} \left[ \left( 1 + \frac{3}{2} \frac{1+c_1}{1-c_1} \right) \ln \frac{1-c_1}{2} \right. \\ &\quad \left. + \left( 1 + \frac{1-c_1}{1+c_1} + \frac{1}{2} \frac{1+c_1}{1-c_1} \right) \ln^2 \frac{1-c_1}{2} + (c_1 \rightarrow -c_1) \right], \\ c_1 &= \cos \theta_1, \quad \theta_1 = \widehat{\mathbf{q}_1 \mathbf{p}_-}. \end{aligned} \quad (12)$$

In addition, the three-photon production process

$$e^+(p_+) + e^-(p_-) \rightarrow \gamma(q_1) + \gamma(q_2) + \gamma(q_3)$$

must be included. Its cross section is given by


$$\begin{aligned} d\sigma^{e^+e^- \rightarrow 3\gamma} &= \frac{\alpha^3}{8\pi^2 s} R_{3\gamma} d\Gamma_{3\gamma}, \\ R_{3\gamma} &= s \frac{\chi_3^2 + (\chi_3')^2}{\chi_1 \chi_2 \chi_1' \chi_2'} - 2m_e^2 \left[ \frac{\chi_1^2 + \chi_2^2}{\chi_1 \chi_2 (\chi_3')^2} + \frac{(\chi_1')^2 + (\chi_2')^2}{\chi_1' \chi_2' \chi_3^2} \right] \\ &\quad + (\text{cyclic permutations}), \\ d\Gamma_{3\gamma} &= \frac{d^3 q_1 d^3 q_2 d^3 q_3}{q_1^0 q_2^0 q_3^0} \delta^{(4)}(p_+ + p_- - q_1 - q_2 - q_3), \end{aligned} \quad (13)$$

where

$$\chi_i = q_i p_-, \quad \chi_i' = q_i p_+, \quad i = 1, 2, 3.$$

The process has to be treated as a radiative correction to the two-photon production. The energy of the third photon should exceed the soft-photon energy threshold  $\Delta E$ . In practice, the tree photon contribution, as well as the radiative Bhabha process  $e^+e^- \rightarrow e^+e^-\gamma$ , should be simulated with the help of a MC event generator in order to take into account the proper experimental criteria of a given event selection.

In addition to the corrections discussed above, also the effect of vacuum polarization, due to the insertion of fermion loops inside the photon propagators, must be included in the precise calculation of the Bhabha scattering cross section. Its theoretical treatment, which faces the non-trivial problem of the non-perturbative contribution due to light quarks, is addressed in detail in Section 6. However, numerical results for such a correction are presented in Section 2.6 and Section 2.8.



**Fig. 7.** Cross sections of the processes  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow \gamma\gamma$  in the LO and NLO approximations, as a function of the c.m. energy at meson factories (upper panel). In the lower panel, the relative contribution due to the NLO QED corrections (in per cent) to the two processes is shown.

In Fig. 7 the cross sections of the Bhabha and two-photon production processes in the LO and NLO approximations are shown as a function of the c.m. energy between  $\sqrt{s} \simeq 2m_\pi$  and  $\sqrt{s} \simeq 10$  GeV (upper panel). The results were obtained imposing the following cuts for the Bhabha process:

$$\begin{aligned} \theta_{\pm}^{\min} &= 45^\circ & \theta_{\pm}^{\max} &= 135^\circ \\ E_{\pm}^{\min} &= 0.3\sqrt{s} & \xi_{\max} &= 10^\circ \end{aligned} \quad (14)$$

where  $\theta_{\pm}^{\min, \max}$  are the angular acceptance cuts,  $E_{\pm}^{\min}$  are the minimum energy thresholds for the detection of the final-state electron/positron and  $\xi_{\max}$  is the maximum  $e^+e^-$  acollinearity. For the photon pair production processes we used correspondingly:

$$\begin{aligned} \theta_{\gamma}^{\min} &= 45^\circ & \theta_{\gamma}^{\max} &= 135^\circ \\ E_{\gamma}^{\min} &= 0.3\sqrt{s} & \xi_{\max} &= 10^\circ \end{aligned} \quad (15)$$

where, as in Eq. (14),  $\theta_{\gamma}^{\min, \max}$  are the angular acceptance cuts,  $E_{\gamma}^{\min}$  is the minimum energy threshold for the detection of at least two photons and  $\xi_{\max}$  is the maximum

acollinearity between the most energetic and next-to-most energetic photon.

The cross sections display the typical  $1/s$  QED behaviour. The relative effect of NLO corrections is shown in the lower panel. It can be seen that the NLO corrections are largely negative and increase with increasing c.m. energies, because of the growing importance of the collinear logarithm  $L = \ln(s/m_e^2)$ . The corrections to  $e^+e^- \rightarrow \gamma\gamma$  are about one half of those to Bhabha scattering, because of the absence of final-state radiation effects in photon pair production.

## 2.3 NNLO Corrections to the Bhabha Scattering Cross Section

Beyond the NLO corrections discussed in the previous Section, in recent years a significant effort was devoted to the calculation of the perturbative corrections to the Bhabha process at the NNLO in QED.

The calculation of the full NNLO corrections to the Bhabha scattering cross section requires three types of ingredients: *i*) the two-loop matrix elements for the  $e^+e^- \rightarrow e^+e^-$  process; *ii*) the one-loop matrix elements for the  $e^+e^- \rightarrow e^+e^-\gamma$  process, both in the case in which the additional photon is soft or hard; *iii*) the tree-level matrix elements for  $e^+e^- \rightarrow e^+e^-\gamma\gamma$ , with two soft, two hard or one soft and one hard photons. Also the process  $e^+e^- \rightarrow e^+e^-e^+e^-$ , with one of the two  $e^+e^-$  pairs remaining undetected, contributes to the Bhabha signature at NNLO. Depending on the kinematics, other final states like, e.g.,  $e^+e^-\mu^+\mu^-$  or those with hadrons are also possible.

The advent of new calculational techniques and a deeper understanding of the IR structure of unbroken gauge theories, such as QED or QCD, made the calculation of the complete set of two-loop QED corrections possible. The history of this calculation will be presented in Section 2.3.1.

Some remarks on the one-loop matrix elements with three particles in the final state are in order now. The diagrams involving the emission of a soft photon are known and they were included in the calculations of the two-loop matrix elements, in order to remove the IR soft divergencies. However, although the contributions due to a hard collinear photon are taken into account in logarithmic accuracy by the MC generators, a full calculation of the diagrams involving a hard photon in a general phase-space configuration is still missing. In Section 2.3.2, we shall comment on the possible strategies which can be adopted in order to calculate these corrections <sup>1</sup>.

As a general comment, it must be noticed that the fixed-order corrections calculated up to NNLO are taken into account at the LL, and, partially, next-to-leading-log (NLL) level in the most precise MC generators, which include, as will be discussed in Section 2.4 and Section

2.5, the logarithmically enhanced contributions of soft and collinear photons at all orders in perturbation theory.

Concerning the tree level graphs with four particles in the final state, the production of a soft  $e^+e^-$  pair was considered in the literature by the authors of [94] by following the evaluation of pair production [95,96] within the calculation of the  $\mathcal{O}(\alpha^2 L)$  single-logarithmic accurate small-angle Bhabha cross section [35], and it is included in the two-loop calculation (see Section 2.3.1). New results on lepton and hadron pair corrections which are at present approximately included in the available Bhabha codes are presented in Section 2.3.3.

### 2.3.1 Virtual corrections for the $e^+e^- \rightarrow e^+e^-$ process

The calculation of the virtual two-loop QED corrections to the Bhabha scattering differential cross section was carried out in the last 10 years. This calculation was made possible by an improvement of the techniques employed in the evaluation of multiloop Feynman diagrams. An essential tool used to manage the calculation is the Laporta algorithm [97,98,99,100], which allows to reduce a generic combination of dimensionally-regularized scalar integrals to a combination of a small set of independent integrals called the “Master Integrals” (MIs) of the problem under consideration. The calculation of the MIs is then pursued by means of a variety of methods. Particularly important are the differential equations method [101,102,103,104,105,106,107] and the Mellin-Barnes techniques [108,109,110,111,112,113,114,115,116,117]. Both methods proved to be very useful in the evaluation of virtual corrections to Bhabha scattering since they are especially effective in problems with a small number of different kinematic parameters. They both allow one to obtain an analytic expression for the integrals, which must be written in terms of a suitable functional basis. A basis which was extensively employed in the calculation of multiloop Feynman diagrams of the type discussed here is represented by the Harmonic Polylogarithms [118,119,120,121,122,123,124,125,126] and their generalizations. Another fundamental achievement which allowed to complete the calculation of the QED two-loop corrections was an improved understanding of the IR structure of QED. In particular, the relation between the collinear logarithms in which the electron mass  $m_e$  plays the role of a natural cut-off and the corresponding poles in the dimensionally regularized massless theory was extensively investigated in [127,128,129,130].

The first complete diagrammatic calculation of the two-loop QED virtual corrections to Bhabha scattering can be found in [131]. However, this result was obtained in the fully massless approximation ( $m_e = 0$ ), by employing dimensional regularization (DR) to regulate both soft and collinear divergencies. Today, the complete set of two-loop corrections to Bhabha scattering in pure QED have been evaluated using  $m_e$  as a collinear regulator, as required in order to include these fixed-order calculations in available Monte Carlo event generators. The Feynman diagrams involved in the calculation can be divided in three

<sup>1</sup> As it will be emphasized in Section 2.8 and Section 2.9 the complete calculation of this class of corrections became available [93] during the completion of the present work.

gauge-independent sets: *i*) diagrams without fermion loops (“photonic” diagrams), *ii*) diagrams involving a closed electron loop, and *iii*) diagrams involving a closed loop of hadrons or a fermion heavier than the electron. Some of the diagrams belonging to the aforementioned sets are shown in Figs. 8–11. These three sets are discussed in more detail below.

### Photonic Corrections


A large part of the NNLO photonic corrections can be evaluated in a closed analytic form, retaining the full dependence on  $m_e$  [132], by using the Laporta algorithm for the reduction of the Feynman diagrams to a combination of MIs, and then the differential equations method for their analytic evaluation. With this technique it is possible to calculate, for instance, the NNLO corrections to the form factors [133, 134, 135, 136]. However, a calculation of the two-loop photonic boxes retaining the full dependence on  $m_e$  seems to be beyond the reach of this method. This is due to the fact that the number of MIs belonging to the same topology is, in some cases, large. Therefore, one must solve analytically large systems of first-order ordinary linear differential equations; this is not possible in general. Alternatively, in order to calculate the different MIs involved, one could use the Mellin-Barnes techniques, as shown in [114, 115, 136, 137, 138, 139], or a combination of both methods. The calculation is very complicated and a full result is not available yet<sup>2</sup>. However, the full dependence on  $m_e$  is not phenomenologically relevant. In fact, the physical problem exhibits a well defined mass hierarchy. The mass of the electron is always very small compared to the other kinematic invariants and can be safely neglected everywhere, with the exception of the terms in which it acts as a collinear regulator. The ratio of the photonic NNLO corrections to the Born cross section is the following

$$\frac{d\sigma^{(2,\text{PH})}}{d\sigma^{(\text{Born})}} = \left(\frac{\alpha}{\pi}\right)^2 \sum_{i=0}^2 \delta^{(\text{PH},i)} (L_e)^i + \mathcal{O}\left(\frac{m_e^2}{s}, \frac{m_e^2}{t}\right), \quad (16)$$

where  $L_e = \ln(s/m_e^2)$  and the coefficients  $\delta^{(\text{PH},i)}$  contain infrared logarithms and are functions of the scattering angle  $\theta$ . The approximation given by Eq. (16) is sufficient for a phenomenological description of the process<sup>3</sup>. The coefficients of the double and single collinear logarithm in Eq. (16),  $\delta^{(\text{PH},2)}$  and  $\delta^{(\text{PH},1)}$ , were obtained in [140, 141]. However, the precision required for luminosity measurements at  $e^+e^-$  colliders demands the calculation of the non-logarithmic coefficient,  $\delta^{(\text{PH},0)}$ . The latter was obtained in [127, 128] by reconstructing the differential cross

<sup>2</sup> For the planar double box diagrams, all the MIs are known [137] for small  $m_e$ , while the MIs for the non-planar double box diagrams are not completed.


<sup>3</sup> It can be shown that the terms suppressed by a positive power of  $m_e^2/s$  do not play any phenomenological role already at very low c.m. energies,  $\sqrt{s} \sim 10$  MeV. Moreover, the terms  $m_e^2/t$  (or  $m_e^2/u$ ) become important in the extremely forward (backward) region, unreachable for the experimental setup.



**Fig. 8.** Some of the diagrams belonging to the class of the “photonic” NNLO corrections to the Bhabha scattering differential cross section. The additional photons in the final state are soft.

section in the  $s \gg m_e^2 \neq 0$  limit from the dimensionally regularized massless approximation [131]. The main idea of the method developed in [127, 128] is outlined below. As far as the leading term in the small electron mass expansion is considered, the difference between the massive and the dimensionally regularized massless Bhabha scattering can be viewed as a difference between two regularization schemes for the infrared divergences. With the known massless two-loop result at hand, the calculation of the massive one is reduced to constructing the *infrared matching* term which relates the two above mentioned regularization schemes. To perform the matching an auxiliary amplitude is constructed, which has the same structure of the infrared singularities but it is sufficiently simple to be evaluated at least at the leading order in the small mass expansion. A particular form of the auxiliary amplitude is dictated by the general theory of infrared singularities in QED and involves the exponent of the one-loop correction as well as the two-loop corrections to the logarithm of the electron form factor. The difference between the full and the auxiliary amplitudes is infrared finite. It can be evaluated by using dimensional regularization for each amplitude and then taking the limit of four space-time dimensions. The infrared divergences, which induce the asymptotic dependence of the virtual corrections on the electron and photon masses, are absorbed into the auxiliary amplitude while the technically most nontrivial calculation of the full amplitude is performed in the massless approximation. The matching of the massive and massless results is then necessary only for the auxiliary amplitude and it is straightforward. Thus the two-loop massless result for the scattering amplitude along with the two-loop massive electron form factor [142] are sufficient to obtain the two-loop photonic correction to the differential cross section in the small electron mass limit.

A method based on a similar principle was subsequently developed in [129, 130]; the authors of [130] confirmed the result of [127, 128] for the NNLO photonic corrections to the Bhabha scattering differential cross section.



**Fig. 9.** Some of the diagrams belonging to the class of the “electron loop” NNLO corrections. The additional photons or electron-positron pair in the final state are soft.

### Electron Loop Corrections

The NNLO electron loop corrections arise from the interference of two-loop Feynman diagrams with the tree-level amplitude as well as from the interference of one-loop diagrams, as long as one of the diagrams contributing to each term involves a closed electron loop. This set of corrections presents a single two-loop box topology, and it is therefore technically less challenging to evaluate with respect to the photonic correction set. The calculation of the electron loop corrections was completed a few years ago [143, 144, 145, 146]; the final result retains the full dependence of the differential cross section on the electron mass  $m_e$ . The MIs involved in the calculation were identified by means of the Laporta algorithm and evaluated with the differential equation method. As expected, after UV renormalization the differential cross section presented only residual IR poles which were removed by adding the contribution of the soft photon emission diagrams. The resulting NNLO differential cross section could be conveniently written in terms of 1- and 2-dimensional Harmonic Polylogarithms (HPLs) of maximum weight three. Expanding the cross section in the limit  $s, |t| \gg m_e^2$ , the ratio of the NNLO corrections to the Born cross section can be written as in Eq. (16):

$$\frac{d\sigma^{(2, \text{EL})}}{d\sigma^{(\text{Born})}} = \left(\frac{\alpha}{\pi}\right)^2 \sum_{i=0}^3 \delta^{(\text{EL}, i)}(L_e)^i + \mathcal{O}\left(\frac{m_e^2}{s}, \frac{m_e^2}{t}\right). \quad (17)$$

Note that the series now contains a cubic collinear logarithm. This logarithm appears, with an opposite sign, in the corrections due to the production of an electron-positron pair (the soft-pair production was considered in [94]). When the two contributions are considered together in the full NNLO, the cubic collinear logarithms cancel. Therefore, the physical cross section includes at most a double logarithm, as in Eq. (16).

The explicit expression of all the coefficients  $\delta^{(\text{EL}, i)}$ , obtained by expanding the results of [143, 144, 145] was confirmed by two different groups [130, 146]. In [130] the small electron mass expansion was performed within the soft-collinear effective theory (SCET) framework, while the analysis in [146] employed the asymptotic expansion

of the MIs.

### Heavy-Flavor and Hadronic Corrections

Finally, we consider the corrections originating from two-loop Feynman diagrams involving a heavy flavor fermion loop<sup>4</sup>. Since this set of corrections involves one more mass scale with respect to the corrections analyzed in the previous sections, a direct diagrammatic calculation is in principle a more challenging task. Recently, in [130] the authors applied their technique based on SCET to Bhabha scattering and obtained the heavy flavor NNLO corrections in the limit in which  $s, |t|, |u| \gg m_f^2 \gg m_e^2$ , where  $m_f$  is the mass of the heavy fermion running in the loop. Their result was very soon confirmed in [146] by means of a method based on the asymptotic expansion of Mellin-Barnes representation of the MIs involved in the calculation. However, the results obtained in the approximation  $s, |t|, |u| \gg m_f^2 \gg m_e^2$  cannot be applied to the case in which  $\sqrt{s} < m_f$  (as in the case of a tau loop at  $\sqrt{s} \sim 1$  GeV), and they apply only to a relatively narrow angular region perpendicular to the beam direction when  $\sqrt{s}$  is not very much larger than  $m_f$  (as in the case of top-quark loops at ILC). It was therefore necessary to calculate the heavy flavor corrections to Bhabha scattering assuming only that the electron mass is much smaller than the other scales in the process, but retaining the full dependence on the heavy mass,  $s, |t|, |u|, m_f^2 \gg m_e^2$ .

The calculation was carried out in two different ways: in [147, 148] it was done analytically, while in [149, 150] it was done numerically with dispersion relations.


The technical problem of the diagrammatic calculation of Feynman integrals with four scales can be simplified by considering carefully, once more, the structure of the collinear singularities of the heavy-flavor corrections. The ratio of the NNLO heavy flavor corrections to the Born cross section is given by

$$\frac{d\sigma^{(2, \text{HF})}}{d\sigma^{(\text{Born})}} = \left(\frac{\alpha}{\pi}\right)^2 \sum_{i=0}^1 \delta^{(\text{HF}, i)}(L_e)^i + \mathcal{O}\left(\frac{m_e^2}{s}, \frac{m_e^2}{t}\right), \quad (18)$$

where now the coefficients  $\delta^{(i)}$  are functions of the scattering angle  $\theta$  and, in general, of the mass of the heavy fermions involved in the virtual corrections. It is possible to prove that, in a physical gauge, all the collinear singularities factorize and can be absorbed in the external field renormalization [151]. This observation has two consequences in the case at hand. The first one is that box diagrams are free of collinear divergencies in a physical gauge; since the sum of all boxes forms a gauge independent block, it can be concluded that the sum of all box diagrams is free of collinear divergencies in any gauge. The second consequence is that the single collinear logarithm in Eq. (18) arises from vertex corrections only. Moreover, if one chooses on-shell UV renormalization conditions, the irreducible two-loop vertex graphs are free

<sup>4</sup> Here by “heavy flavor” we mean a muon or a  $\tau$ -lepton, as well as a heavy quark, like the top, the  $b$ - or the  $c$ -quark, depending on the c.m. energy range that we are considering.





**Fig. 10.** Some of the diagrams belonging to the class of the “heavy fermion” NNLO corrections. The additional photons in the final state are soft.

of collinear singularities. Therefore, among all the two-loop diagrams contributing to the NNLO heavy flavor corrections to Bhabha scattering, only the reducible vertex corrections are logarithmically divergent in the  $m_e \rightarrow 0$  limit<sup>5</sup>. The latter are easily evaluated even if they depend on two different masses. By exploiting these two facts, one can obtain the NNLO heavy-flavor corrections to the Bhabha scattering differential cross section assuming only that  $s, |t|, |u|, m_f^2 \gg m_e^2$ . In particular, one can set  $m_e = 0$  from the beginning in all the two-loop diagrams with the exception of the reducible ones. This procedure allows one to effectively eliminate one mass scale from the two-loop boxes, so that these graphs can be evaluated with the techniques already employed in the diagrammatic calculation of the electron loop corrections<sup>6</sup>. In the case in which the heavy flavor fermion is a quark, it is straightforward to modify the calculation of the two-loop self-energy diagrams to obtain the mixed QED-QCD corrections to Bhabha scattering [148].

An alternative approach to the calculation of the heavy flavor corrections to Bhabha scattering, is based on dispersion relations. This method also applies to hadronic corrections. The hadronic and heavy-fermion corrections to the Bhabha-scattering cross section can be obtained by appropriately inserting the renormalized irreducible photon vacuum-polarization function  $\Pi$  in the photon propagator:

$$\frac{g_{\mu\nu}}{q^2 + i\delta} \rightarrow \frac{g_{\mu\alpha}}{q^2 + i\delta} (q^2 g^{\alpha\beta} - q^\alpha q^\beta) \Pi(q^2) \frac{g_{\beta\nu}}{q^2 + i\delta}. \quad (19)$$

The vacuum polarization  $\Pi$  can be represented by a once-subtracted dispersion integral [6],

$$\Pi(q^2) = -\frac{q^2}{\pi} \int_{4M^2}^{\infty} dz \frac{\text{Im } \Pi(z)}{z} \frac{1}{q^2 - z + i\delta}. \quad (20)$$

<sup>5</sup> Additional collinear logarithms arise also from the interference of one-loop diagrams in which at least one vertex is present.

<sup>6</sup> The necessary MIs can be found in [148, 152, 153, 154].

The contributions to  $\Pi$  may then be determined from a (properly normalized) production cross section by the optical theorem [155],

$$\text{Im } \Pi_{\text{had}}(z) = -\frac{\alpha}{3} R_{\text{had}}(z). \quad (21)$$

In this way, the hadronic vacuum polarization may be obtained from the experimental data for  $R_{\text{had}}$ :

$$R_{\text{had}}(z) = \frac{\sigma(\{e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons}\}; z)}{(4\pi\alpha^2)/(3z)}. \quad (22)$$

In the low-energy region the inclusive experimental data may be used [27, 156]. Around a narrow hadronic resonance with mass  $M_{\text{res}}$  and width  $\Gamma_{\text{res}}^{e^+e^-}$  one may use the relation

$$R_{\text{res}}(z) = \frac{9\pi}{\alpha^2} M_{\text{res}} \Gamma_{\text{res}}^{e^+e^-} \delta(z - M_{\text{res}}^2), \quad (23)$$

and in the remaining regions the perturbative QCD prediction [157]. Contributions to  $\Pi$  arising from leptons and heavy quarks with mass  $m_f$ , charge  $Q_f$  and color  $C_f$  can be computed directly in perturbation theory:

$$R_f(z; m_f) = Q_f^2 C_f \left( 1 + 2 \frac{m_f^2}{z} \right) \sqrt{1 - 4 \frac{m_f^2}{z}}. \quad (24)$$

As a result of the above formulas, the massless photon propagator gets replaced by a massive propagator, whose effective mass  $z$  is subsequently integrated over:

$$\frac{g_{\mu\nu}}{q^2 + i\delta} \rightarrow \frac{\alpha}{3\pi} \int_{4M^2}^{\infty} \frac{dz R(z)}{z(q^2 - z + i\delta)} \left( g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2 + i\delta} \right). \quad (25)$$


For self-energy corrections to Bhabha scattering at one-loop order, the dispersion relation approach was first employed in [158]. Two-loop applications of this technique, prior to Bhabha scattering, are the evaluation of the hadronic vertex correction [159] and of two-loop hadronic corrections to the lifetime of the muon [160]. The approach was also applied to the evaluation of the two-loop form factors in QED in [161, 162, 163].

The fermionic and hadronic corrections to Bhabha scattering at one-loop accuracy come only from the self-energy diagram; see for details Section 6. At two-loop level there are reducible and irreducible self-energy contributions, vertices and boxes. The reducible corrections are easily treated. For the evaluation of the irreducible two-loop diagrams, it is advantageous that they are one-loop diagrams with self-energy insertions, because the application of the dispersion technique as described here is possible.

The kernel function for the irreducible two-loop vertex was derived in [159] and verified in e.g. [150] and the three kernel functions for the two-loop box functions were first obtained in [164, 149, 150] and verified in [165]. A complete collection of all the relevant formulae may be found in [150], and the corresponding Fortran code bhhbnnloh is publicly available at the web page [166]

[www-zeuthen.desy.de/theory/research/bhabha/](http://www-zeuthen.desy.de/theory/research/bhabha/).





**Fig. 11.** Some of the diagrams belonging to the class of the “hadronic” corrections. The additional photons in the final state are soft.

In [150], the dependence of the various heavy-fermion NNLO corrections on  $\ln(s/m_f^2)$  for  $s, |t|, |u| \gg m_f^2$  was studied. The irreducible vertex behaves (before a combination with real pair emission terms) like  $\ln^3(s/m_f^2)$  [159], while the sum of the various infrared divergent diagrams as a whole behaves like  $\ln(s/m_f^2) \ln(s/m_e^2)$ . This is in accordance with Eq. (18), but the limit plays no effective role at the energies studied here.


As a result of efforts of recent years we have now for all the non-photon virtual two-loop contributions at least two completely independent calculations. The net result as a ratio of the NNLO corrections to the Born cross section in per mille, is shown in Fig. 12 for KLOE and in Fig. 13 for BaBar/Belle<sup>7</sup>. While the non-photon corrections stay at 1 per mille or less for KLOE, they reach a few per mille at the BaBar/Belle energy range. The NNLO photonic corrections are the dominant contributions and amount to some per mille, both at  $\phi$  and  $B$  factories. However, as already emphasized, the bulk of both photonic and non-photon corrections is incorporated into the generators used by the experimental collaborations. Hence, the consistent comparison between the results of NNLO calculations and the MC predictions at the same perturbative level allows to assess the theoretical accuracy of the luminosity tools, as will be discussed quantitatively in Section 2.8.

### 2.3.2 Fixed-Order Calculation of the Hard Photon Emission at One Loop


The one-loop matrix element for the process  $e^+e^- \rightarrow e^+e^-\gamma$  is one of the contributions to the complete set of NNLO corrections to Bhabha scattering. Its evaluation requires the non-trivial computation of one-loop tensor integrals associated to pentagon-diagrams.

According to the standard Passarino-Veltman (PV) approach [168], one-loop tensor integrals can be expressed in terms of MIs with trivial numerators that are independent of the loop variable, each multiplied by a Lorentz

<sup>7</sup> The pure self-energy corrections deserve a special discussion and are thus omitted in the plots.



**Fig. 12.** Two-loop photonic and non-photon corrections to Bhabha scattering at  $\sqrt{s} = 1.02$  GeV, normalized to the QED tree-level cross section, as a function of the electron polar angle. No cuts. The parameterizations of  $R_{had}$  due to [167] and [27, 156, 157] are very close to each other.



**Fig. 13.** Two-loop photonic and non-photon corrections to Bhabha scattering at  $\sqrt{s} = 10.56$  GeV, normalized to the QED tree-level cross section, as a function of the electron polar angle. No cuts. The parameterizations of  $R_{had}$  is due to [167].

structure depending only on combinations of the external momenta and the metric tensor. The achievement of the complete PV-reduction amounts to solve a non-trivial system of equations. Due to its size, it is reasonable replacing the analytic techniques by numerical tools. It is difficult to implement the PV-reduction numerically, since it gives rise to Gram determinants. The latter naturally arise in the procedure of inverting a system and they can vanish at special phase space points. This fact requires a proper modification of the reduction algorithm [169, 170, 171, 172,

[173, 174, 175]. A viable solution for the complete algebraic reduction of tensor-pentagon (and tensor-hexagon) integrals was formulated in [176, 177, 178], by exploiting the algebra of signed minors [179]. In this approach the cancellation of powers of inverse Gram determinants was performed recently in [180, 181].

The computation of the one-loop five-point amplitude  $e^+e^- \rightarrow e^+e^-\gamma$  can be alternatively performed by using generalized-unitarity cutting rules (see [182] for a detailed compilation of references). In the following we propose two ways to achieve the result, *via* an analytical and *via* a semi-numerical method. The application of generalized cutting-rules as an on-shell method of calculation is based on two fundamental properties of scattering amplitudes: *i*) analyticity, according to which any amplitude is determined by its singularity structure [183, 184, 185, 155, 186]; *ii*) and unitarity, according to which the residues at the singularities are determined by products of simpler amplitudes. Turning these properties into a tool for computing scattering amplitudes is possible because of the underlying representation of the amplitude in terms of Feynman integrals and their PV-reduction, which grants the existence of a representation of any one-loop amplitudes as linear combination of MIs, each multiplied by a rational coefficient. In the case of  $e^+e^- \rightarrow e^+e^-\gamma$ , pentagon-integrals may be expressed, through PV-reduction, by a linear combination of 17 MIs (including 3 boxes, 8 triangles, 5 bubbles, and 1 tadpole). Since the required MIs are analytically known [187, 188, 189, 177, 171, 190, 191], the determination of their coefficients is needed for reconstructing the amplitude as a whole. To this aim, one may use the Mathematica program hexagon [180, 181]. Also matching the generalized cuts of the amplitude with the cuts of the MIs provides an efficient way to extract their (rational) coefficients out of the amplitude itself. In general the fulfillment of multiple-cut conditions requires loop momenta with complex components. The effect of the cut conditions is to freeze some of its components, when not all, according to the number of the cuts. With the quadruple-cut [192] the loop momentum is completely frozen, yielding the algebraic determination of the coefficients of  $n$ -point functions with  $n \geq 4$ . In cases where fewer than four denominators are cut, like triple-cut [193, 194, 195], double-cut [196, 197, 198, 199, 200, 194], and single-cut [201], the loop momentum is not frozen: the free-components are left over as phase-space integration variables.

For each multiple-cut, the evaluation of the phase-space integral would generate, in general, logarithms and a non-logarithmic term. The coefficient of a given  $n$ -point MI finally appears in the non-logarithmic term of the corresponding  $n$ -particle cut, where all the internal lines are on-shell (while the logarithms correspond to the cuts of higher-point MIs which share that same cut). Therefore all the coefficients of MIs can be determined in a *top-down* algorithm, starting from the quadruple-cuts for the extraction of the 4-point coefficients, and following with the triple-, double-, and single-cuts, for the coefficients of 3-, 2- and 1-point, respectively. The coefficient of an  $n$ -point MI ( $n \geq 2$ ) can be also obtained by specializing to

the case at hands the generating formulas given in [202] for general one-loop amplitudes.

Instead of the analytic evaluation of the multiple-cut phase-space integrals, it is worth considering the feasibility of computing the process  $e^+e^- \rightarrow e^+e^-\gamma$  with a seminumerical technique by now known as OPP-reduction [203, 204], based on the decomposition of the numerator of any one-loop integrand in terms of its denominators [205, 206, 207, 208]. Within this approach the coefficients of the MIs can be found simply by solving a system of numerical equations, avoiding any explicit integration. The OPP-reduction algorithm exploits the polynomial structures of the integrand when evaluated at values of the loop-momentum fulfilling multiple cut-conditions: *i*) for each  $n$ -point MI, one considers the  $n$ -particle cut obtained by setting all the propagating lines on-shell; *ii*) such a cut is associated to a polynomial in terms of the free components of the loop-momentum, which corresponds to the numerator of the integrand evaluated at the solution of the on-shell conditions; *iii*) the constant-term of that polynomial is the coefficient of the MI.

Hence the difficult task of evaluating one-loop Feynman integrals is reduced to the much simpler problem of polynomial fitting, recently optimized by using a projection technique based on the Discrete Fourier Transform [209].

In general the result of a dimensional-regulated amplitude in the 4-dimensional limit, being  $D (= 4 - 2\epsilon)$  the regulating parameter, is expected to contain (poly)logarithms, often referred to as the cut-constructible term, and a pure rational term. In a later paper [210], which completed the OPP-method, the rising of the rational term was attributed to two potential sources (of UV-divergent integrals): one, defined  $R_1$ , due to the  $D$ -dimensional completion of the 4-dimensional contribution of the numerator; a second one, called  $R_2$ , due to the  $(-2\epsilon)$ -dimensional algebra of Dirac-matrices. Therefore in the OPP-approach the calculation of the one-loop amplitude  $e^+e^- \rightarrow e^+e^-\gamma$  can proceed through two computational stages:

1. the coefficients of the MIs that are responsible both for the cut-constructible and for the  $R_1$ -rational terms can be determined by applying the OPP-reduction discussed above [203, 204, 209];
2. the  $R_2$ -rational term can be computed by using additional tree-level-like diagrammatic rules, very much resembling the computation of the counter terms needed for the renormalization of UV-divergencies [210].

The numerical influence of the radiative loop diagrams, including the pentagon diagrams, is expected not to be particularly large. However, the calculation of such corrections would greatly help to assess the physical precision of existing luminosity programs.

### 2.3.3 Pair Corrections

As was mentioned in the paragraph on virtual heavy flavor and hadronic corrections of Section 2.3.1, these virtual corrections have to be combined with real corrections in order to get physically sensible results. The virtual NNLO

**Table 2.** The NNLO lepton and pion pair corrections to the Bhabha scattering Born cross section  $\sigma_B$ : virtual corrections  $\sigma_v$ , soft and hard real photon emissions  $\sigma_s, \sigma_h$ , and pair emission contributions  $\sigma_{pairs}$ . Total pair correction cross sections obtainable from the sum  $\sigma_{s+v+h} + \sigma_{pairs}$ . All cross sections in nanobarns according to the cuts given in the text.

Electron pair corrections					
	$\sigma_B$	$\sigma_h$	$\sigma_{v+s}$	$\sigma_{v+s+h}$	$\sigma_{pairs}$
KLOE	529.469	9.502	-11.567	-2.065	0.271
BaBar	6.744	0.246	-0.271	-0.025	0.017
Muon pair corrections					
	$\sigma_B$	$\sigma_h$	$\sigma_{v+s}$	$\sigma_{v+s+h}$	$\sigma_{pairs}$
KLOE	529.469	1.494	-1.736	-0.241	—
BaBar	6.744	0.091	-0.095	-0.004	0.0005
Tau pair corrections					
	$\sigma_B$	$\sigma_h$	$\sigma_{v+s}$	$\sigma_{v+s+h}$	$\sigma_{pairs}$
KLOE	529.469	0.020	-0.023	-0.003	—
BaBar	6.744	0.016	-0.017	-0.0007	$< 10^{-7}$
Pion pair corrections					
	$\sigma_B$	$\sigma_h$	$\sigma_{v+s}$	$\sigma_{v+s+h}$	$\sigma_{pairs}$
KLOE	529.469	1.174	-1.360	-0.186	—
BaBar	6.744	0.062	-0.065	-0.003	0.00003

electron, muon, tau and pion corrections have to be combined with the emission of real electron, muon, tau and pion pairs, respectively. The real pair production cross sections are finite, but cut dependent. We consider here the pion pair production as it is the dominant part of the hadronic corrections and can serve as an estimate of the role of the whole set of hadronic corrections. The description of all relevant hadronic contributions is much more involved task and will not be covered in this review. As was first explicitly shown for Bhabha scattering in [94] for electron pairs, and also discussed in [150], there appear exact cancellations of terms of the order  $\ln^3(s/m_e^2)$  or  $\ln^3(s/m_f^2)$ , so that the leading terms are at most of order  $\ln^2(s/m_e^2), \ln^2(s/m_f^2)$ .

In Table 2 we show NNLO lepton and pion pair contributions with typical kinematical cuts for the KLOE and BaBar experiments. Besides contributions from unresolved pair emissions  $\sigma_{pairs}$ , we also add unresolved real hard photon emission contributions  $\sigma_h$ . The corrections  $\sigma_{pairs}$  from fermions have been calculated with the Fortran package HELAC-PHEGAS [211, 212, 213, 214], the real pion corrections with EKHARA [215, 216], the NNLO hard photonic corrections  $\sigma_h$  with a program [217] based on the generator BHAGEN-1PH [218]. The latter depend, technically, on the soft photon cut-off  $E_\gamma^{\min} = \omega$ . After adding up with  $\sigma_{v+s}$ , the sum of the two  $\sigma_{v+s+h}$  is independent of that; in fact we use here  $\omega/E_{\text{beam}} = 10^{-4}$ . The  $\sigma_{v+s}$  is determined with an updated version of the Fortran package bhhnnlohf [150, 166] in order to cover also pion pair corrections. The cuts applied in Table 2 for the KLOE experiment are

- $\sqrt{s} = 1.02$  GeV
- $E_{\min} = 0.4$  GeV
- $55^\circ < \theta_\pm < 125^\circ$

- $\xi_{\max} = 9^\circ$ ,

and for the BaBar experiment

- $\sqrt{s} = 10.56$  GeV
- $|\cos(\theta_\pm)| < 0.7$  and  $|\cos(\theta_-)| < 0.65$
- $|\mathbf{p}_+|/E_{\text{beam}} > 0.75$  and  $|\mathbf{p}_-|/E_{\text{beam}} > 0.5$  or  $|\mathbf{p}_-|/E_{\text{beam}} > 0.75$  and  $|\mathbf{p}_+|/E_{\text{beam}} > 0.5$
- $\xi_{\max}^{3d} = 30^\circ$ .

Here  $E_{\min}$  is the energy threshold for the final-state electron/positron,  $\theta_\pm$  are the electron/positron polar angles and  $\xi_{\max}$  is the maximum allowed polar angle acollinearity:

$$\xi = |\theta_+ + \theta_- - 180^\circ|, \quad (26)$$

and  $\xi_{\max}^{3d}$  is the maximum allowed three dimensional acollinearity:

$$\xi^{3d} = \left| \arccos \left( \frac{\mathbf{p}_+ \cdot \mathbf{p}_-}{(|\mathbf{p}_-||\mathbf{p}_+|)} \right) \times \frac{180^\circ}{\pi} - 180^\circ \right|. \quad (27)$$

For  $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ , cuts are applied only to the  $e^+e^-$  pair. In the case of  $e^+e^- \rightarrow e^+e^-e^+e^-$ , all possible  $e^\pm e^\mp$  combinations are checked and if at least one pair fulfils the cuts the event is accepted.

At KLOE the electron pair corrections contribute about  $3 \times 10^{-3}$  and at BaBar about  $1 \times 10^{-3}$ , while all the other contributions of pair production are even smaller. Like in small-angle Bhabha scattering at LEP/SLC the pair corrections [219] are largely dominated by the electron pair contribution.

## 2.4 Multiple photon effects and matching with NLO corrections

### 2.4.1 Universal methods for leading logarithmic corrections

From inspection of Eq. (9) and Eq.(12) for the SV NLO QED corrections to the cross section of the Bhabha scattering and  $e^+e^- \rightarrow \gamma\gamma$  process, it can be seen that large logarithms  $L = \ln(s/m_e^2)$ , due to collinear photon emission, are present. Similar large logarithmic terms arise after integration of the hard photon contributions from the kinematical domains of photon emission at small angles with respect to charged particles. For the energy range of meson factories the logarithm is large numerically, i.e.  $L \sim 15$  at the  $\phi$ -factories and  $L \sim 20$  at the  $B$ -factories, and the corresponding terms give the bulk of the total radiative correction. These contributions represent also the dominant part of the NNLO effects discussed in Section 2.3. Therefore the logarithmically enhanced contributions due to emission of soft and collinear photons must be taken into account at all orders in perturbation theory, to achieve the required theoretical accuracy.

The methods for the calculation of higher-order (HO) QED corrections on the basis of the generators employed nowadays at flavour factories were already widely and successfully used in the 90s at LEP/SLC for electroweak tests

of the SM. They were adopted for the calculation of both the small-angle Bhabha scattering cross section (necessary for the high-precision luminosity measurement) and  $Z$ -boson observables. Hence, the theory accounting for the control of HO QED corrections at meson factories can be considered particularly robust, having passed the very stringent tests of LEP/SLC era.

The most popular and standard methods to keep under control multiple photon effects are the QED Structure Function (SF) approach [220, 221, 222, 223] and Yennie-Frautschi-Suura (YFS) exponentiation [224]. The former is used in all the versions of the generator BabaYaga [225, 226, 227] and MCGPJ [228] (albeit according to different realizations), while the latter is the theoretical recipe adopted in BHWIDE [229]. Actually, analytical QED SFs  $D(x, Q^2)$ , valid in the strictly collinear approximation, are implemented in MCGPJ, whereas BabaYaga is based on a MC Parton Shower (PS) algorithm to reconstruct  $D(x, Q^2)$  numerically.

### *The Structure Function approach*

Let us consider the annihilation process  $e^-e^+ \rightarrow X$ , where  $X$  is some given final state and  $\sigma_0(s)$  its LO cross section. Initial-state (IS) QED radiative corrections can be described according to the following picture. Before arriving at the annihilation point, the incoming electron (positron) of four-momentum  $p_{-(+)}$  radiates real and virtual photons. These photons, due to the dynamical features of QED, are mainly radiated along the direction of motion of the radiating particles, and their effect is mainly to reduce the original four-momentum of the incoming electron (positron) to  $x_{1(2)}p_{-(+)}$ . After this pre-emission, the hard scattering process  $e^-(x_1p_-)e^+(x_2p_+) \rightarrow X$  takes place, at a reduced squared c.m. energy  $\hat{s} = x_1x_2s$ . The resulting cross section, corrected for IS QED radiation, can be represented as follows [220, 221, 222]

$$\sigma(s) = \int_0^1 dx_1 dx_2 D(x_1, s) D(x_2, s) \sigma_0(x_1 x_2 s) \Theta(\text{cuts}), \quad (28)$$

where  $D(x, s)$  is the electron SF, representing the probability that an incoming electron (positron) radiates a collinear photon, retaining a fraction  $x$  of its original momentum at the energy scale  $Q^2 = s$ , and  $\Theta(\text{cuts})$  stands for a rejection algorithm taking care of experimental cuts. When considering photonic radiation only the non-singlet part of the SF is of interest. If the running of the QED coupling constant is neglected, the non-singlet part of the SF is the solution of the following Renormalization Group (RG) equation, analogous to the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) equation of QCD [230, 231, 232]:

$$s \frac{\partial}{\partial s} D(x, s) = \frac{\alpha}{2\pi} \int_x^1 \frac{dz}{z} P_+(z) D\left(\frac{x}{z}, s\right), \quad (29)$$

where  $P_+(z)$  is the regularized Altarelli-Parisi (AP) splitting function electron  $\rightarrow$  electron + photon, given by

$$P_+(z) = P(z) - \delta(1-z) \int_0^1 dx P(x),$$

$$P(z) = \frac{1+z^2}{1-z}. \quad (30)$$

Equation (29) can be also transformed into an integral equation, subject to the boundary condition  $D(x, m_e^2) = \delta(1-x)$ :

$$D(x, s) = \delta(1-x) + \frac{\alpha}{2\pi} \int_{m_e^2}^s \frac{dQ^2}{Q^2} \int_x^1 \frac{dz}{z} P_+(z) D\left(\frac{x}{z}, Q^2\right). \quad (31)$$

Equation (31) can be solved exactly by means of numerical methods, such as the inverse Mellin transform method. However, this derivation of  $D(x, s)$  turns out to be problematic in view of phenomenological applications. Therefore, approximate (but very accurate) analytical representations of the solution of the evolution equation are of major interest for practical purposes. This type of solution was the one typically adopted in the context of LEP/SLC phenomenology. A first analytical solution can be obtained in the soft photon approximation, i.e. in the limit  $x \simeq 1$ . This solution, also known as Gribov-Lipatov (GL) approximation, exponentiates at all perturbative orders the large logarithmic contributions of infrared and collinear origin, but it does not take into account hard-photon (collinear) effects. This drawback can be overcome by solving the evolution equation iteratively. At the  $n$ -th step of the iteration, one obtains the  $\mathcal{O}(\alpha^n)$  contribution to the SF for any value of  $x$ . By combining the GL solution with the iterative one, in which the soft-photon part has been eliminated in order to avoid double counting, one can build a hybrid solution of the evolution equation. It exploits all the positive features of the two kinds of solutions and is not affected by the limitations intrinsic to each of them. Two classes of hybrid solutions, namely the additive and factorized ones, are known in the literature and both were adopted for applications to LEP/SLC precision physics. A typical additive solution, where the GL approximation  $D_{GL}(x, s)$  is supplemented by finite-order terms present



in the iterative solution, is given by [233]

$$\begin{aligned}
D_A(x, s) &= \sum_{i=0}^3 d_A^{(i)}(x, s), \\
d_A^{(0)}(x, s) &= \frac{\exp\left[\frac{1}{2}\beta\left(\frac{3}{4} - \gamma_E\right)\right]}{\Gamma\left(1 + \frac{1}{2}\beta\right)} \frac{1}{2}\beta(1-x)^{\frac{1}{2}\beta-1}, \\
d_A^{(1)}(x, s) &= -\frac{1}{4}\beta(1+x), \\
d_A^{(2)}(x, s) &= \frac{1}{32}\beta^2[(1+x)(-4\ln(1-x) + 3\ln x) \\
&\quad - 4\frac{\ln x}{1-x} - 5 - x], \\
d_A^{(3)}(x, s) &= \frac{1}{384}\beta^3\left\{(1+x)[18\zeta(2) - 6\text{Li}_2(x) \right. \\
&\quad - 12\ln^2(1-x)] + \frac{1}{1-x}\left[-\frac{3}{2}(1+8x+3x^2)\ln x \right. \\
&\quad + \frac{1}{2}(1+7x^2)\ln^2 x - 12(1+x^2)\ln x \ln(1-x) \\
&\quad - 6(x+5)(1-x)\ln(1-x) \\
&\quad \left. - \frac{1}{4}(39-24x-15x^2)\right]\Big\}, \tag{32}
\end{aligned}$$

where  $\Gamma$  is the Euler gamma-function,  $\gamma_E \approx 0.5772$  the Euler-Mascheroni constant,  $\zeta$  the Riemann  $\zeta$ -function and  $\beta$  is the large collinear factor

$$\beta = \frac{2\alpha}{\pi} \left[ \ln\left(\frac{s}{m_e^2}\right) - 1 \right]. \tag{33}$$

Explicit examples of factorized solutions, which are obtained by multiplying the GL solution by finite-order terms, in such a way that, order by order, the iterative contributions are exactly recovered, can be found in [234]. For the calculation of HO corrections with a per mille accuracy analytical SFs in additive and factorized form containing up to  $\mathcal{O}(\alpha^3)$  finite-order terms are sufficient and in excellent agreement. They also agree with the accuracy much better than 0.1 with the exact numerical solution of the QED evolution equation. Explicit solutions up to the fifth order in  $\alpha$  were calculated in [235, 236].

The RG method described above was applied in [237] for the treatment of LL QED radiative corrections to various processes of interest for physics at meson factories. Such a formulation was later implemented in the generator MCGPJ. For example, according to [237], the Bhabha scattering cross section, accounting for LL terms in all orders,  $\mathcal{O}(\alpha^n L^n)$ ,  $n = 1, 2, \dots$ , of perturbation theory, is given by

$$\begin{aligned}
d\sigma_{\text{LLA}}^{\text{Bhabha}} &= \sum_{a,b,c,d=e^\pm, \gamma} \int_{\bar{z}_1}^1 dz_1 \int_{\bar{z}_2}^1 dz_2 D_{ae^-}^{\text{str}}(z_1) D_{be^+}^{\text{str}}(z_2) \\
&\quad \times d\sigma_0^{ab \rightarrow cd}(z_1, z_2) \int_{\bar{y}_1}^1 \frac{dy_1}{Y_1} D_{e^-c}^{\text{frg}}\left(\frac{y_1}{Y_1}\right) \int_{\bar{y}_2}^1 \frac{dy_2}{Y_2} D_{e^+d}^{\text{frg}}\left(\frac{y_2}{Y_2}\right) \\
&\quad + \mathcal{O}\left(\alpha^2 L, \alpha \frac{m_e^2}{s}\right), \tag{34}
\end{aligned}$$

where  $d\sigma_0^{ab \rightarrow cd}(z_1, z_2)$  is the differential LO cross section of the process  $ab \rightarrow cd$  with energy fractions of the incoming particles being scaled by factors  $z_1$  and  $z_2$  with respect to the initial electron and positron, respectively. In the notation of [237], the electron SF  $D_{ab}^{\text{str}}(z)$  is distinguished from the electron fragmentation function  $D_{ab}^{\text{frg}}(z)$  to point out the role played by IS radiation (described by  $D_{ab}^{\text{str}}(z)$ ) with respect to the one due to final-state radiation (described by  $D_{ab}^{\text{frg}}(z)$ ). However, because of their probabilistic meaning, the electron structure and fragmentation functions coincide. In Eq. (34) the quantities  $Y_{1,2}$  are the energy fractions of particles  $c$  and  $d$  with respect to the beam energy. Explicit expressions for  $Y_{1,2} = Y_{1,2}(z_1, z_2, \cos\theta)$  and other details on the kinematics can be found in [237]. The lower limits of the integrals,  $\bar{z}_{1,2}$  and  $\bar{y}_{1,2}$ , should be defined according to the experimental conditions of particle detection and kinematical constraints. For the case of the  $e^+e^- \rightarrow \gamma\gamma$  process, one has to change the master formula (34) by picking up the two-photon final state. Formally it can be done just by choosing the proper fragmentation functions,  $D_{\gamma c}^{\text{frg}}$  and  $D_{\gamma d}^{\text{frg}}$ .

The photonic part of the non-singlet electron structure (fragmentation) function in  $\mathcal{O}(\alpha^n L^n)$  considered in [237] reads as follows

$$\begin{aligned}
D_{ee}^{NS, \gamma}(z) &= \delta(1-z) + \sum_{i=1}^n \left(\frac{\alpha}{2\pi}(L-1)\right)^i \frac{1}{i!} \left[P_{ee}^{(0)}(z)\right]^{\otimes i}, \\
D_{\gamma e}(z) &= \frac{\alpha}{2\pi}(L-1)P_{\gamma e}(z) + \mathcal{O}(\alpha^2 L^2), \\
D_{e\gamma}(z) &= \frac{\alpha}{2\pi}LP_{\gamma e}(z) + \mathcal{O}(\alpha^2 L^2), \\
P_{ee}^{(0)}(z) &= \left[\frac{1+z^2}{1-z}\right]_+ \\
&= \lim_{\Delta \rightarrow 0} \left\{ \delta(1-z)(2\ln \Delta + \frac{3}{2}) + \Theta(1-z-\Delta) \frac{1+z^2}{1-z} \right\}, \\
\left[P_{ee}^{(0)}(z)\right]^{\otimes i} &= \int_z^1 \frac{dt}{t} P_{ee}^{(i-1)}(t) P_{ee}^{(0)}\left(\frac{z}{t}\right), \tag{35} \\
P_{e\gamma}(z) &= z^2 + (1-z)^2, \quad P_{\gamma e}(z) = \frac{1+(1-z)^2}{z}.
\end{aligned}$$

Starting from the second order in  $\alpha$  there appear also non-singlet and singlet  $e^+e^-$  pair contributions to the structure function:

$$\begin{aligned}
D_{ee}^{NS, e^+e^-}(z) &= \frac{1}{3} \left(\frac{\alpha}{2\pi}L\right)^2 P_{ee}^{(1)}(z) + \mathcal{O}(\alpha^3 L^3), \\
D_{ee}^{S, e^+e^-}(z) &= \frac{1}{2!} \left(\frac{\alpha}{2\pi}L\right)^2 R(z) + \mathcal{O}(\alpha^3 L^3), \\
R(z) &= P_{e\gamma} \otimes P_{\gamma e}(z) = \frac{1-z}{3z}(4+7z+4z^2) \\
&\quad + 2(1+z)\ln z. \tag{36}
\end{aligned}$$

Note that radiation of a real pair, *i.e.* appearance of additional electrons and positrons in the final state, require the application of nontrivial conditions of experimental particle registration. Unambiguously, that can be done only



within a MC event generator based on four-particle matrix elements, as already discussed in Section 2.3.

In the same way as in QCD, the LL cross sections depend on the choice of the factorization scale  $Q^2$  in the argument of the large logarithm  $L = \ln(Q^2/m_e^2)$ , which is not fixed a priori by the theory. However, the scale should be taken of the order of the characteristic energy transfer in the process under consideration. Typical choices are  $Q^2 = s$ ,  $Q^2 = -t$ , and  $Q^2 = st/u$ . The first one is good for annihilation channels, like  $e^+e^- \rightarrow \mu^+\mu^-$ , the second one is optimal for small-angle Bhabha scattering, where the  $t$ -channel exchange dominates, see [238]. The last choice allows to exponentiate the leading contribution due to initial-final state interference [239] and is particularly suited for large-angle Bhabha scattering in QED. The option  $Q^2 = st/u$  is adopted in all the versions of the generator BabaYaga. Reduction of the scale dependence can be achieved by taking into account next-to-leading corrections in  $\mathcal{O}(\alpha^n L^{n-1})$ , next-to-next-to-leading ones in  $\mathcal{O}(\alpha^n L^{n-2})$  etc.

#### The Parton Shower algorithm

The PS algorithm is a method for providing a MC iterative solution of the evolution equation, at the same time generating the four-momenta of the electron and photon at a given step of the iteration. It was developed within the context of QCD and later applied in QED too.

In order to implement the algorithm, it is first necessary to assume the existence of an upper limit for the energy fraction  $x$ , in such a way that the AP splitting function is regularized as follows

$$P_+(z) = \theta(x_+ - z)P(z) - \delta(1 - z) \int_0^{x_+} dx P(x). \quad (37)$$

Of course, in the limit  $x_+ \rightarrow 1$  Eq. (37) recovers the usual definition of the AP splitting function given in Eq. (30). By inserting the modified AP vertex into Eq. (29), one obtains

$$\begin{aligned} s \frac{\partial}{\partial s} D(x, s) &= \frac{\alpha}{2\pi} \int_x^{x_+} \frac{dz}{z} P(z) D\left(\frac{x}{z}, s\right) \\ &\quad - \frac{\alpha}{2\pi} D(x, s) \int_x^{x_+} dz P(z). \end{aligned} \quad (38)$$

Separating the variables and introducing the Sudakov form factor

$$\Pi(s_1, s_2) = \exp \left[ -\frac{\alpha}{2\pi} \int_{s_2}^{s_1} \frac{ds'}{s'} \int_0^{x_+} dz P(z) \right], \quad (39)$$

which is the probability that the electron evolves from virtuality  $-s_2$  to  $-s_1$  without emitting photons of energy fraction larger than  $1 - x_+ \equiv \epsilon$  ( $\epsilon \ll 1$ ), Eq. (38) can be recast into integral form as follows

$$\begin{aligned} D(x, s) &= \Pi(s, m_e^2) D(x, m_e^2) \\ &\quad + \frac{\alpha}{2\pi} \int_{m_e^2}^s \frac{ds'}{s'} \Pi(s, s') \int_x^{x_+} \frac{dz}{z} P(z) D\left(\frac{x}{z}, s'\right). \end{aligned} \quad (40)$$

The formal iterative solution of Eq. (40) can be represented by the following infinite series

$$\begin{aligned} D(x, s) &= \sum_{n=0}^{\infty} \prod_{i=1}^n \left\{ \int_{m_e^2}^{s_{i-1}} \frac{ds_i}{s_i} \Pi(s_{i-1}, s_i) \right. \\ &\quad \times \left. \frac{\alpha}{2\pi} \int_{x/(z_1 \dots z_{i-1})}^{x_+} \frac{dz_i}{z_i} P(z_i) \right\} \Pi(s_n, m_e^2) D\left(\frac{x}{z_1 \dots z_n}, m_e^2\right). \end{aligned} \quad (41)$$

The particular form of Eq. (41) allows to exploit a MC method for building the solution iteratively. The steps of the algorithm are as follows:

- 1 – set  $Q^2 = m_e^2$ , and fix  $x = 1$  according to the boundary condition  $D(x, m_e^2) = \delta(1 - x)$ ;
- 2 – generate a random number  $\xi$  in the interval  $[0, 1]$ ;
- 3 – if  $\xi < \Pi(s, Q^2)$  stop the evolution; otherwise
- 4 – compute  $Q'^2$  as solution of the equation  $\xi = \Pi(Q'^2, Q^2)$ ;
- 5 – generate a random number  $z$  according to the probability density  $P(z)$  in the interval  $[0, x_+]$ ;
- 6 – substitute  $x \rightarrow xz$  and  $Q^2 \rightarrow Q'^2$ ; go to 2.

The  $x$  distribution of the electron SF as obtained by means of the PS algorithm and a numerical solution (based on the inverse Mellin transform method) of the QED evolution equation is shown in Fig. 14. Perfect agreement is seen. Once  $D(x, s)$  has been reconstructed by the algorithm, the master formula of Eq. (28) can be used for the calculation of LL corrections to the cross section of interest. This cross section must be independent of the soft-hard photon separator  $\epsilon$ , in the limit of small  $\epsilon$  values. This can be clearly seen in Fig. 15, where the QED corrected Bhabha cross section as a function of the fictitious parameter  $\epsilon$  is shown for DAΦNE energies with the cuts of Eq. (14), but for an angular acceptance  $\theta_{\pm}$  of  $55^\circ \div 125^\circ$ . The cross section reaches a plateau for  $\epsilon$  smaller than  $10^{-4}$ .

The main advantage of the PS algorithm with respect to the analytical solutions of the electron evolution is the possibility of going beyond the strictly collinear approximation and generating transverse momentum  $p_{\perp}$  of electrons and photons at each branching. In fact, the kinematics of the branching process  $e(p) \rightarrow e'(p') + \gamma(q)$  can be written as


$$\begin{aligned} p &= (E, \mathbf{0}, p_z) \\ p' &= (zE, \mathbf{p}_{\perp}, p'_z) \\ q &= ((1 - z)E, -\mathbf{p}_{\perp}, q_z). \end{aligned} \quad (42)$$

Once the variables  $p^2$ ,  $p'^2$  and  $z$  are generated by the PS algorithm, the on-shell condition  $q^2 = 0$ , together with the longitudinal momentum conservation, allows to obtain an expression for the  $p_{\perp}$  variable:


$$p_{\perp}^2 = (1 - z)(zp^2 - p'^2), \quad (43)$$

at first order in  $p^2/E^2 \ll 1$ ,  $p_{\perp}^2/E^2 \ll 1$ .

However, some not correct behaviour of the exclusive photon kinematics reconstruction is related to this PS picture, due to the approximations inherent to Eq. (43). First



**Fig. 14.** Comparison for the  $x$  distribution of the electron SF as obtained by means of a numerical solution of the QED evolution equation (solid line) and the PS algorithm (histogram). From [225].



**Fig. 15.** QED corrected Bhabha cross section at DAΦNE as a function of the infrared regulator  $\epsilon$  of the PS approach, according to the setup of Eq. (14). The error bars correspond to  $1\sigma$  MC errors. From [227].

of all, since within the PS algorithm the generation of  $p'^2$  and  $z$  are independent, it can happen that in some branching the  $p_\perp^2$  as given by Eq. (43) is negative. In order to avoid this problem, the introduction of any kinematical cut on  $p^2$  or  $z$  generation (or the regeneration of the whole event) would mean a not correct reconstruction of the SF  $x$  distribution, which is important for a precise cross section calculation. Furthermore, in the PS scheme, each fermion produces its photon cascade independently of the other ones, missing the effects due to the interference of radiation coming from different charged particles. As far as inclusive cross sections (i.e., no cuts are imposed on the generated photons) are considered, these effects are largely integrated out but they become important when more ex-

clusive variables distributions are looked at, as shown in [240]

Concerning the first problem, it can be overcome choosing a generation of  $p_\perp$  of the photons different from Eq. (43). For example, one can choose to extract the photon  $\cos\vartheta_\gamma$  according to the universal leading poles  $1/p \cdot k$  present in the matrix element for photon emission. Namely, one can generate  $\cos\vartheta_\gamma$  as

$$\cos\vartheta_\gamma \propto \frac{1}{1 - \beta \cos\vartheta_\gamma}, \quad (44)$$

where  $\beta$  is the speed of the emitting particle. In this way, photon energy and angle are generated independently, different from Eq. (43). The nice feature of this prescription is that  $p_\perp^2 = E_\gamma^2 \sin^2\vartheta_\gamma$  is always well defined and the  $x$  distribution reproduces exactly the SF, because any further kinematical cuts must be imposed to avoid unphysical events. At this stage, the PS is used only to generate the energies and multiplicity of the photons. The problem of including the radiation interference is still unsolved, because the variables of photons emitted by a fermion are still uncorrelated with those of the other charged particles. The issue of including photon interference can be successfully worked out looking at the YFS formula [224]:

$$d\sigma_n \approx d\sigma_0 \frac{e^{2n}}{n!} \prod_{l=1}^n \frac{d^3\mathbf{k}_l}{(2\pi)^3 2k_l^0} \sum_{i,j=1}^N \eta_i \eta_j \frac{-p_i \cdot p_j}{(p_i \cdot k_l)(p_j \cdot k_l)}. \quad (45)$$

It gives the differential cross section  $d\sigma_n$  for the emission of  $n$  photons, whose momenta are  $k_1, \dots, k_n$ , from a kernel process described by  $d\sigma_0$  and involving  $N$  fermions, whose momenta are  $p_1, \dots, p_N$ . In Eq. (45)  $\eta_i$  is a charge factor, which is  $+1$  for incoming  $e^-$  or outgoing  $e^+$  and  $-1$  for incoming  $e^+$  or outgoing  $e^-$ . Note that Eq. (45) is valid in the soft limit ( $k_i \rightarrow 0$ ). The important point is that it also accounts for coherence effects. From the YFS formula it is straightforward to read out the angular spectrum of the  $l^{th}$  photon:

$$\cos\vartheta_l \propto - \sum_{i,j=1}^N \eta_i \eta_j \frac{1 - \beta_i \beta_j \cos\vartheta_{ij}}{(1 - \beta_i \cos\vartheta_{il})(1 - \beta_j \cos\vartheta_{jl})}. \quad (46)$$

It is worth noticing that in the LL prescription, the same quantity writes as

$$\cos\vartheta_l \propto \sum_{i=1}^N \frac{1}{1 - \beta_i \cos\vartheta_{il}}, \quad (47)$$

whose terms are of course contained in Eq. (46).

In order to consider also coherence effects in the angular distribution of the photons, one can generate  $\cos\vartheta_\gamma$  according to Eq. (46), rather than to Eq. (47). This recipe [240] is adopted in BabaYaga v3.5 and BabaYaga@NLO.

#### Yennie-Frautschi-Suura exponentiation

The YFS exponentiation procedure, implemented in the code BHWIDE, is a technique for summing up all the

infrared (IR) singularities present in any process accompanied by photonic radiation [224]. It is inherently exclusive, i.e. all the summations of the IR singular contributions are done before any phase-space integration over the virtual or real photon four-momenta are performed. The method was mainly developed by S. Jadach, B.F.L. Ward and collaborators to realize precision MC tools. In the following, the general ideas underlying the procedure are summarized.

Let us consider the scattering process  $e^+(p_1)e^-(p_2) \rightarrow f_1(q_1) \cdots f_n(q_n)$ , where  $f_1(q_1) \cdots f_n(q_n)$  represents a given arbitrary final state, and let  $\mathcal{M}_0$  be its tree-level matrix element. By using standard Feynman-diagram techniques, it is possible to show that the same process, when accompanied by  $l$  additional real photons radiated by the IS particles, and under the assumption that the  $l$  additional photons are soft, i.e. their energy is much smaller than any energy scale involved in the process, can be described by the factorized matrix element built up by the LO one,  $\mathcal{M}_0$ , times the product of  $l$  eikonal currents, namely

$$\mathcal{M} \simeq \mathcal{M}_0 \prod_{i=1}^l \left[ e \left( \frac{\varepsilon_i(k_i) \cdot p_2}{k_i \cdot p_2} - \frac{\varepsilon_i(k_i) \cdot p_1}{k_i \cdot p_1} \right) \right], \quad (48)$$

where  $e$  is the electron charge,  $k_i$  are the momenta of the photons and  $\varepsilon_i(k_i)$  their polarization vectors. Taking the square of the matrix element in Eq. (48) and multiplying by the proper flux factor and the Lorentz-invariant phase space volume, the cross section for the process  $e^+(p_1)e^-(p_2) \rightarrow f_1(q_1) \cdots f_n(q_n) + l$  real photons can be written as

$$d\sigma_r^{(l)} = d\sigma_0 \frac{1}{l!} \prod_{i=1}^l \left[ k_i dk_i d\cos\vartheta_i d\varphi_i \frac{1}{2(2\pi)^3} \sum_{\varepsilon_i} e^2 \left( \frac{\varepsilon_i(k_i) \cdot p_2}{k_i \cdot p_2} - \frac{\varepsilon_i(k_i) \cdot p_1}{k_i \cdot p_1} \right)^2 \right]. \quad (49)$$

By summing over the number of final-state photons, one obtains the cross section for the original process accompanied by an arbitrary number of real photons, namely

$$\begin{aligned} d\sigma_r^{(\infty)} &= \sum_{l=0}^{\infty} d\sigma_r^{(l)} \\ &= d\sigma_0 \exp \left[ k dk d\cos\vartheta d\varphi \frac{1}{2(2\pi)^3} \sum_{\varepsilon} e^2 \left( \frac{\varepsilon(k) \cdot p_2}{k \cdot p_2} - \frac{\varepsilon(k) \cdot p_1}{k \cdot p_1} \right)^2 \right]. \end{aligned} \quad (50)$$

Equation (50), being limited to real radiation only, is IR divergent once the phase space integrations are performed down to zero photon energy. This problem, as well known, finds its solution in the matching between real and virtual photonic radiation. Equation (50) already shows the key feature of exclusive exponentiation, i.e. summing up all the perturbative contributions before performing any phase space integration.

In order to get meaningful radiative corrections, besides IS real photon corrections it is necessary to consider also IS virtual photon corrections, i.e. the corrections due to additional internal photon lines connecting the IS electron and positron. For a vertex-type amplitude, the result can be written as

$$\mathcal{M}_{V_1} = -i \frac{e^2}{(2\pi)^4} \int d^4k \frac{1}{k^2 + i\varepsilon} \bar{v}(p_1) \gamma^\mu \frac{-(\not{p}_1 + \not{k}) + m}{2p_1 \cdot k + k^2 + i\varepsilon} \times \Gamma \frac{(\not{p}_2 + \not{k}) + m}{2p_2 \cdot k + k^2 + i\varepsilon} \gamma_\mu u(p_2), \quad (51)$$

where  $\Gamma$  stands for the Dirac structure of the LO process, in such a way that  $\mathcal{M}_0 = \bar{v}(p_1) \Gamma u(p_2)$ . The soft-photon part of the amplitude can be extracted by taking  $k^\mu \simeq 0$  in all the numerators. In this approximation, the amplitude of Eq. (51) becomes

$$\begin{aligned} \mathcal{M}_{V_1} &= \mathcal{M}_0 \times V, \\ V &= \frac{2i\alpha}{(2\pi)^3} \int d^4k \frac{4p_1 \cdot p_2}{(2p_1 \cdot k + k^2 + i\varepsilon)(2p_2 \cdot k + k^2 + i\varepsilon)} \times \frac{1}{k^2 + i\varepsilon}. \end{aligned} \quad (52)$$

It can be seen that, as in the real case, the IR virtual correction factorizes off the LO matrix element so that it is universal, i.e. independent of the details of the process under consideration, and divergent in the IR portion of the phase space.

The correction given by  $n$  soft virtual photons can be seen to factorize with an additional  $1/n!$  factor, namely

$$\mathcal{M}_{V_n} = \mathcal{M}_0 \times \frac{1}{n!} V^n, \quad (53)$$

so that by summing over all the additional soft virtual photons one obtains

$$\mathcal{M}_V = \mathcal{M}_0 \times \exp[V]. \quad (54)$$

As already noticed both the real and virtual factors are IR divergent. In order to obtain meaningful expressions one has to adopt some regularization procedure. One possibility is to give the photon a (small) mass  $\lambda$  and modify Eqs. (49) and (52) accordingly. Once all the expressions are properly regularized, one can write down the YFS master formula, which takes into account real and virtual photonic corrections to the LO process. In virtue of the factorization properties discussed above, the master formula can be obtained from Eq. (50) with the substitution  $d\sigma_0 \rightarrow d\sigma_0 |\exp(V)|^2$ , i.e.

$$\begin{aligned} d\sigma &= d\sigma_0 |\exp(V)|^2 \exp \left[ k dk d\cos\vartheta d\varphi \frac{1}{2(2\pi)^3} \sum_{\varepsilon} e^2 \left( \frac{\varepsilon(k) \cdot p_2}{k \cdot p_2} - \frac{\varepsilon(k) \cdot p_1}{k \cdot p_1} \right)^2 \right]. \end{aligned} \quad (55)$$

As a last step it is possible to perform analytically the IR cancellation between virtual and very soft real photons. Actually, since very soft real photons do not affect

the kinematics of the process, the real photon exponent can be split into a contribution coming from photons with energy less than a cutoff  $k_{min}$  plus a contribution from photons with energy above it. The first contribution can be integrated over all its phase space and then combined with the virtual exponent. After this step it is possible to remove the regularizing photon mass by taking the limit  $\lambda \rightarrow 0$ , so that Eq. (55) becomes

$$d\sigma = d\sigma_0 \exp(Y) \exp \left[ k dk d\Theta(k - k_{min}) \cos \vartheta d\varphi \frac{1}{2(2\pi)^3} \sum_{\varepsilon} e^2 \left( \frac{\varepsilon(k) \cdot p_2}{k \cdot p_2} - \frac{\varepsilon(k) \cdot p_1}{k \cdot p_1} \right)^2 \right], \quad (56)$$

where  $Y$  is given by

$$Y = 2V + \int k dk d\Theta(k_{min} - k) \cos \vartheta d\varphi \frac{1}{2(2\pi)^3} \sum_{\varepsilon} e^2 \left( \frac{\varepsilon(k) \cdot p_2}{k \cdot p_2} - \frac{\varepsilon(k) \cdot p_1}{k \cdot p_1} \right)^2. \quad (57)$$

The explicit form of  $Y$  can be derived by performing all the details of the calculation, and reads

$$Y = \beta \ln \frac{k_{min}}{E} + \delta_{YFS}, \quad \delta_{YFS} = \frac{1}{4}\beta + \frac{\alpha}{\pi} \left( \frac{\pi^2}{3} - \frac{1}{2} \right). \quad (58)$$

#### 2.4.2 Matching NLO and higher-order corrections

As it will be shown numerically in Section 2.6, NLO corrections must be combined with multiple photon emission effects to achieve a theoretical accuracy at the per mille level. This combination, technically known as *matching*, is on the grounds of the most precise generators used for luminosity monitoring, i.e. BabaYaga@NLO, BHWIDE and MCGPJ. Although the matching is implemented according to different theoretical details, some general aspects are common to all the recipes and must be emphasized:

1. it is possible to match NLO and HO corrections consistently, avoiding double counting of LL contributions at order  $\alpha$  and preserving the advantages of resummation of soft and collinear effects beyond  $\mathcal{O}(\alpha)$ ;
2. the convolution of NLO corrections with HO terms allows to include the dominant part of NNLO corrections, given by infrared-enhanced  $\alpha^2 L$  sub-leading contributions. This was argued and demonstrated analytically and numerically in [36] through comparison with the available  $\mathcal{O}(\alpha^2)$  corrections to  $s$ -channel processes and  $t$ -channel Bhabha scattering. Such an aspect of the matching procedure is crucial to settle the theoretical accuracy of the generators by means of explicit comparisons with the exact NNLO perturbative corrections discussed in Section 2.3, and will be addressed in Section 2.8
3. BabaYaga@NLO and BHWIDE implement a fully factorized matching recipe, while MCGPJ includes some terms in additive form, as will be visible in the formulae to be reported.

In the following we summarize the basic features of the matching procedure as implemented in the codes MCGPJ, BabaYaga@NLO and BHWIDE.

The matching approach realized in the MC event generator MCGPJ was developed in [228]. In particular Bhabha scattering with complete  $\mathcal{O}(\alpha)$  and HO LL photonic corrections is represented in the following way:

$$\begin{aligned} \frac{d\sigma^{e^+e^- \rightarrow e^+e^- (\gamma)}}{d\Omega_-} &= \int_{\bar{z}_1}^1 dz_1 \int_{\bar{z}_2}^1 dz_2 D_{ee}^{NS,\gamma}(z_1) D_{ee}^{NS,\gamma}(z_2) \\ &\times \frac{d\hat{\sigma}_0^{\text{Bhabha}}(z_1, z_2)}{d\Omega_-} \left( 1 + \frac{\alpha}{\pi} K_{SV} \right) \Theta(\text{cuts}) \\ &\times \int_{y_{th}}^{Y_1} \frac{dy_1}{Y_1} \int_{y_{th}}^{Y_2} \frac{dy_2}{Y_2} D_{ee}^{NS,\gamma}\left(\frac{y_1}{Y_1}\right) D_{ee}^{NS,\gamma}\left(\frac{y_2}{Y_2}\right) \\ &+ \frac{\alpha}{\pi} \int_{\Delta}^1 \frac{dx}{x} \left\{ \left[ \left( 1 - x + \frac{x^2}{2} \right) \ln \frac{\theta_0^2(1-x)^2}{4} + \frac{x^2}{2} \right] \right. \\ &\times 2 \frac{d\hat{\sigma}_0^{\text{Bhabha}}}{d\Omega_-} + \left[ \left( 1 - x + \frac{x^2}{2} \right) \ln \frac{\theta_0^2}{4} + \frac{x^2}{2} \right] \\ &\times \left[ \frac{d\hat{\sigma}_0^{\text{Bhabha}}(1-x, 1)}{d\Omega_-} + \frac{d\hat{\sigma}_0^{\text{Bhabha}}(1, 1-x)}{d\Omega_-} \right] \left. \right\} \Theta(\text{cuts}) \\ &- \frac{\alpha^2}{4s} \left( \frac{3+c^2}{1-c} \right)^2 \frac{8\alpha}{\pi} \ln(\text{ctg} \frac{\theta}{2}) \ln \frac{\Delta\varepsilon}{\varepsilon} \\ &+ \frac{\alpha^3}{2\pi^2 s} \int_{\substack{k^0 > \Delta\varepsilon \\ \theta_i > \theta_0}} \frac{WT}{4} \Theta(\text{cuts}) \frac{d\Gamma_{e\bar{e}\gamma}}{d\Omega_-}. \end{aligned} \quad (59)$$

Here the step functions  $\Theta(\text{cuts})$  stand for the particular cuts applied. The auxiliary parameter  $\theta_0$  defines cones around the directions of the motion of the charged particles in which the emission of hard photons is approximated by the factorized form by convolution of collinear radiation factors [241] with the Born cross section. The dependence on the parameters  $\Delta$  and  $\theta_0$  cancels out in the sum with the last term of Eq. (59), where the photon energy and emission angles with respect to all charged particles are limited from below ( $k^0 > \Delta\varepsilon, \theta_i > \theta_0$ ). Taking into account vacuum polarization the Born level Bhabha cross section with reduced energies of the incoming elec-

tron and positron can be cast in the following form:

$$\begin{aligned} \frac{d\hat{\sigma}_0^{\text{Bhabha}}(z_1, z_2)}{d\Omega_-} &= \frac{4\alpha^2}{s^2} \left\{ \frac{1}{|1 - \Pi(\hat{t})|^2} \frac{a^2 + z_2^2(1+c)^2}{2z_1^2(1-c)^2} \right. \\ &+ \frac{1}{|1 - \Pi(\hat{s})|^2} \frac{z_1^2(1-c)^2 + z_2^2(1+c)^2}{2a^2} \\ &\left. - \text{Re} \frac{1}{(1 - \Pi(\hat{t}))(1 - \Pi(\hat{s}))^*} \frac{z_2^2(1+c)^2}{az_1(1-c)} \right\} d\Omega_-, \\ \hat{s} &= z_1 z_2 s, \quad \hat{t} = -\frac{s z_1^2 z_2(1-c)}{z_1 + z_2 - (z_1 - z_2)c}. \end{aligned} \quad (60)$$

where  $\Pi(Q^2)$  is the photon self-energy correction. Note that in the cross section above the cosine of the scattering angle,  $c$ , is given for the original c.m. reference frame of the colliding beams.

For the two-photon production channel, a similar representation is used in MCGPJ:

$$\begin{aligned} d\sigma^{e^+e^- \rightarrow \gamma\gamma(\gamma)} &= \int_{\bar{z}_1}^1 dz_1 D_{ee}^{NS,\gamma}(z_1) \int_{\bar{z}_2}^1 dz_2 D_{ee}^{NS,\gamma}(z_2) \\ &\times d\hat{\sigma}_0^{\gamma\gamma}(z_1, z_2) \left( 1 + \frac{\alpha}{\pi} K_{SV}^{\gamma\gamma} \right) + \frac{\alpha}{\pi} \int_{\Delta}^1 \frac{dx}{x} \\ &\times \left[ \left( 1 - x + \frac{x^2}{2} \right) \ln \frac{\theta_0^2}{4} + \frac{x^2}{2} \right] \left[ d\hat{\sigma}_0(1-x, 1) \right. \\ &\left. + d\hat{\sigma}_0(1, 1-x) \right] + \frac{1}{3} \frac{4\alpha^3}{\pi^2 s^2} \int_{\substack{z_i \geq \Delta \\ \pi - \theta_0 \geq \theta_i \geq \theta_0}} d\Gamma_{3\gamma} \\ &\times \left[ \frac{z_3^2(1+c_3^2)}{z_1^2 z_2^2 (1-c_1^2)(1-c_2^2)} + \text{two cyclic permutations} \right], \\ z_i &= \frac{q_i^0}{\varepsilon}, \quad c_i = \cos \theta_i, \quad \theta_i = \widehat{\mathbf{p} - \mathbf{q}_i}, \end{aligned} \quad (61)$$

where the cross section with reduced energies has the form

$$\frac{d\hat{\sigma}_0^{\gamma\gamma}(z_1, z_2)}{d\Omega_1} = \frac{2\alpha^2}{s} \frac{z_1^2(1-c_1)^2 + z_2^2(1+c_1)^2}{(1-c_1^2)(z_1 + z_2 + (z_2 - z_1)c_1)^2},$$

and the factor  $1/3$  in the last term of Eq. (61) takes into account the identity of the final-state photons. The sum of the last two terms does not depend on  $\Delta$  and  $\theta_0$ .

Concerning BabaYaga@NLO the matching starts from the observation that Eq. (28) for the QED corrected all-order cross section can be rewritten in terms of the PS ingredients as

$$d\sigma_{LL}^\infty = \Pi(Q^2, \varepsilon) \sum_{n=0}^{\infty} \frac{1}{n!} |\mathcal{M}_{n,LL}|^2 d\Phi_n, \quad (62)$$

The expansion at  $\mathcal{O}(\alpha)$  of Eq. (62) does not coincide, by construction, with an exact  $\mathcal{O}(\alpha)$  result. In fact

$$\begin{aligned} d\sigma_{LL}^\alpha &= \left[ 1 - \frac{\alpha}{2\pi} I_+ \log \frac{Q^2}{m^2} \right] |\mathcal{M}_0|^2 d\Phi_0 + |\mathcal{M}_{1,LL}|^2 d\Phi_1 \\ &\equiv [1 + C_{\alpha,LL}] |\mathcal{M}_0|^2 d\Phi_0 + |\mathcal{M}_{1,LL}|^2 d\Phi_1, \end{aligned} \quad (63)$$

where  $I_+ \equiv \int_0^{1-\varepsilon} P(z) dz$ , whereas an exact NLO cross section can be always cast in the form

$$d\sigma^\alpha = [1 + C_\alpha] |\mathcal{M}_0|^2 d\Phi_0 + |\mathcal{M}_1|^2 d\Phi_1. \quad (64)$$

The coefficients  $C_\alpha$  contain the complete  $\mathcal{O}(\alpha)$  virtual and soft-bremsstrahlung corrections in units of the Born squared amplitude and  $|\mathcal{M}_1|^2$  is the exact squared matrix element with the emission of one hard photon. We remark that  $C_{\alpha,LL}$  has the same logarithmic structure as  $C_\alpha$  and that  $|\mathcal{M}_{1,LL}|^2$  has the same singular behaviour like  $|\mathcal{M}_1|^2$ .

In order to match the LL and NLO calculations the following correction factors, which are by construction infrared safe and free of collinear logarithms, are introduced

$$F_{SV} = 1 + (C_\alpha - C_{\alpha,LL}), \quad F_H = 1 + \frac{|\mathcal{M}_1|^2 - |\mathcal{M}_{1,LL}|^2}{|\mathcal{M}_{1,LL}|^2}, \quad (65)$$

so that the exact  $\mathcal{O}(\alpha)$  cross section can be expressed, up to terms of  $\mathcal{O}(\alpha^2)$ , in terms of its LL approximation as

$$d\sigma^\alpha = F_{SV}(1 + C_{\alpha,LL}) |\mathcal{M}_0|^2 d\Phi_0 + F_H |\mathcal{M}_{1,LL}|^2 d\Phi_1. \quad (66)$$

Driven by Eq. (66), Eq. (62) can be improved by writing the resummed matched cross section as

$$\begin{aligned} d\sigma_{\text{matched}}^\infty &= F_{SV} \Pi(Q^2, \varepsilon) \\ &\sum_{n=0}^{\infty} \frac{1}{n!} \left( \prod_{i=0}^n F_{H,i} \right) |\mathcal{M}_{n,LL}|^2 d\Phi_n. \end{aligned} \quad (67)$$

The correction factors  $F_{H,i}$  follow from the definition (65) for each photon emission. The  $\mathcal{O}(\alpha)$  expansion of Eq. (67) coincides now with the exact NLO cross section of Eq. (64) and all HO LL contributions are the same as in Eq. (62). This formulation is implemented in BabaYaga@NLO for both Bhabha scattering and photon pair production, using, of course, the appropriate SV and hard bremsstrahlung formulae. This matching formulation has been also applied to the study of Drell-Yan-like processes, by combining the complete  $\mathcal{O}(\alpha)$  electroweak corrections with QED shower evolution in the generator HORACE [242, 243, 244, 245].

As far as BHWIDE is concerned, this MC event generator realizes the process

$$e^+(p_1) + e^-(q_1) \longrightarrow e^+(p_2) + e^-(q_2) + \gamma_1(k_1) + \dots + \gamma_n(k_n) \quad (68)$$

via the YFS exponentiated cross section formula

$$\begin{aligned} d\sigma &= e^{2\alpha\Re B + 2\alpha\tilde{B}} \sum_{n=0}^{\infty} \frac{1}{n!} \int \prod_{j=1}^n \frac{d^3 k_j}{k_j^0} \int \frac{d^4 y}{(2\pi)^4} \\ &e^{iy(p_1 + q_1 - p_2 - q_2 - \sum_j k_j) + D} \tilde{\beta}_n(k_1, \dots, k_n) \frac{d^3 p_2 d^3 q_2}{p_2^0 q_2^0}, \end{aligned} \quad (69)$$

where the real infrared function  $\tilde{B}$  and the virtual infrared function  $B$  are given presently [229], and here we note the usual connections

$$\begin{aligned} 2\alpha\tilde{B} &= \int^{k \leq K_{\text{max}}} \frac{d^3 k}{k_0} \tilde{S}(k), \\ D &= \int d^3 k \frac{\tilde{S}(k)}{k^0} (e^{-iy \cdot k} - \theta(K_{\text{max}} - k)), \end{aligned} \quad (70)$$



for the standard YFS infrared real emission factor

$$\tilde{S}(k) = \frac{\alpha}{4\pi^2} \left[ Q_f Q_{f'} \left( \frac{p_1}{p_1 \cdot k} - \frac{q_1}{q_1 \cdot k} \right)^2 + \dots \right], \quad (71)$$

if  $Q_f$  is the electric charge of  $f$  in units of the positron charge. Here, the “...” represent the remaining terms in  $\tilde{S}(k)$  obtained from the one given by respective substitutions of  $Q_f, p_1, Q_{f'}, q_1$  with corresponding values for the other pairs of the respective external charged legs according to the YFS prescription in Ref. [224, 246] (wherein due attention is taken to obtain the correct relative sign of each of the terms in  $\tilde{S}(k)$  according to this latter prescription). We have explicitly the representations

$$\begin{aligned} 2\alpha \Re B(p_1, q_1, p_2, q_2) + 2\alpha \tilde{B}(p_1, q_1, p_2, q_2; k_m) = \\ R_1(p_1, q_1; k_m) + R_1(p_2, q_2; k_m) + R_2(p_1, p_2; k_m) + \\ R_2(q_1, q_2; k_m) - R_2(p_1, q_2; k_m) - R_2(q_1, p_2; k_m), \end{aligned} \quad (72)$$

with

$$R_1(p, q; k_m) = R_2(p, q; k_m) + \left( \frac{\alpha}{\pi} \right) \frac{\pi^2}{2}, \quad (73)$$

$$\begin{aligned} R_2(p, q; k_m) = & \frac{\alpha}{\pi} \left\{ \left( \ln \frac{2pq}{m_e^2} - 1 \right) \ln \frac{k_m^2}{p^0 q^0} + \frac{1}{2} \ln \frac{2pq}{m_e^2} \right. \\ & - \frac{1}{2} \ln^2 \frac{p^0}{q^0} - \frac{1}{4} \ln^2 \frac{(\Delta + \delta)^2}{4p^0 q^0} - \frac{1}{4} \ln^2 \frac{(\Delta - \delta)^2}{4p^0 q^0} \\ & - \Re \text{Li}_2 \left( \frac{\Delta + \omega}{\Delta + \delta} \right) - \Re \text{Li}_2 \left( \frac{\Delta + \omega}{\Delta - \delta} \right) \\ & - \Re \text{Li}_2 \left( \frac{\Delta - \omega}{\Delta + \delta} \right) - \Re \text{Li}_2 \left( \frac{\Delta - \omega}{\Delta - \delta} \right) \\ & \left. + \frac{\pi^2}{3} - 1 \right\}, \end{aligned} \quad (74)$$

where  $\Delta = \sqrt{2pq + (p^0 - q^0)^2}$ ,  $\omega = p^0 + q^0$ ,  $\delta = p^0 - q^0$ , and  $k_m$  is a soft photon cut-off in the CMS ( $E_{\gamma}^{\text{soft}} < k_m \ll E_{\text{beam}}$ ).

The YFS hard photon residuals  $\tilde{\beta}_i$  in (69),  $i = 0, 1$ , are given exactly through  $\mathcal{O}(\alpha)$  in Ref. [229] for BHWIDE, so that this latter event generator calculates the YFS exponentiated exact  $\mathcal{O}(\alpha)$  cross section for  $e^+e^- \rightarrow e^+e^- + n(\gamma)$  with multiple initial, initial-final, and final state radiation using a corresponding MC realization of Eq. (69) in the wide angle regime. The  $\mathcal{O}(\alpha)$  electroweak corrections library, relevant for higher energies, is taken from Refs. [247, 87].

The result (69) is an exact re-arrangement of the loop expansion for the respective cross section and it is independent of the dummy parameter  $K_{\text{max}}$ . To see how to derive it, one may proceed as follows. Let the amplitude for the emission of  $n$  real photons in the Bhabha process be

$$\mathcal{M}^{(n)} = \sum_{\ell} M_{\ell}^{(n)}, \quad (75)$$

$M_{\ell}^{(n)}$  is the contribution to  $\mathcal{M}^{(n)}$  from Feynman diagrams with  $\ell$  virtual loops. The key result in the YFS theory of

Ref. [224, 246] on virtual corrections is that we may rewrite Eq. (75) as the exact representation

$$\mathcal{M}^{(n)} = e^{\alpha B} \sum_{j=0}^{\infty} \mathbf{m}_j^{(n)}, \quad (76)$$

where we have defined

$$\alpha B = \int \frac{d^4 k}{(k^2 - \lambda^2 + i\epsilon)} S(k), \quad (77)$$

with the virtual infrared emission factor given by

$$\begin{aligned} S(k) = & \frac{-i\alpha}{8\pi^2} \sum_{i' < j} Z_{i'} \theta_{i'} Z_j \theta_j \left( \frac{(2\bar{p}_{i'} \theta_{i'} - k)_{\mu}}{k^2 - 2k\bar{p}_{i'} \theta_{i'} + i\epsilon} \right. \\ & \left. + \frac{(2\bar{p}_j \theta_j + k)_{\mu}}{k^2 + 2k\bar{p}_j \theta_j + i\epsilon} \right)^2. \end{aligned} \quad (78)$$

Here,  $\lambda$  is an infrared regulator mass, and we follow Ref. [224, 246] and identify the sign of the  $j$ -th external line charge here as  $Z_j = Q_j$  and  $\theta_j = +(-)$  for outgoing (incoming) 4-momentum  $\bar{p}_j$ , so that here  $\bar{p}_1 = p_1$ ,  $\bar{p}_2 = q_1$ ,  $\bar{p}_3 = p_2$ ,  $\bar{p}_4 = q_2$ ,  $Z_1 = +1$ ,  $\theta_1 = -$ ,  $Z_2 = -1$ ,  $\theta_2 = -$ ,  $Z_3 = +1$ ,  $\theta_3 = +$ ,  $Z_4 = -1$ ,  $\theta_4 = +$ . The amplitudes  $\{\mathbf{m}_j^{(n)}\}$  are free of all virtual infrared divergences.

Using the result (76) for  $\mathcal{M}^{(n)}$ , we get the attendant differential cross section by the standard methods as

$$\begin{aligned} d\hat{\sigma}^n = & \frac{e^{2\alpha \Re B}}{n!} \int \prod_{l=1}^n \frac{d^3 k_l}{(k_l^2 + \lambda^2)^{1/2}} \\ & \times \bar{\rho}^{(n)}(p_1, q_1, p_2, q_2, k_1, \dots, k_n) \frac{d^3 p_2 d^3 q_2}{p_2^0 q_2^0} \\ & \times \delta^{(4)} \left( p_1 + q_1 - p_2 - q_2 - \sum_{i=1}^n k_i \right), \end{aligned} \quad (79)$$

where we have defined

$$\bar{\rho}^{(n)}(p_1, q_1, p_2, q_2, k_1, \dots, k_n) = \sum_{\text{spin}} \left| \sum_{j=0}^{\infty} \mathbf{m}_j^{(n)} \right|^2, \quad (80)$$

in the incoming  $e^+e^-$  CMS system, and we have absorbed the remaining kinematical factors for the initial state flux and spin averages into the normalization of the amplitudes  $\mathcal{M}^{(n)}$  for pedagogical reasons, so that the  $\bar{\rho}^{(n)}$  are averaged over initial spins and summed over final spins. We then use the key result of Ref. [224, 246] on real corrections to write the exact result

$$\begin{aligned} \bar{\rho}^{(n)}(p_1, q_1, p_2, q_2, k_1, \dots, k_n) = & \prod_{i=1}^n \tilde{S}(k_i) \bar{\beta}_0 + \dots + \\ & \sum_{i=1}^n \tilde{S}(k_i) \bar{\beta}_{n-1}(k_1, \dots, k_{i-1}, k_{i+1}, \dots, k_n) \\ & + \bar{\beta}_n(k_1, \dots, k_n), \end{aligned} \quad (81)$$

**Table 3.** MC generators used for luminosity monitoring at meson factories.

Generator	Theory	Accuracy
BabaYaga v3.5	Parton Shower	$\sim 0.5 \div 1\%$
BabaYaga@NLO	$\mathcal{O}(\alpha) + \text{PS}$	$\sim 0.1\%$
BHAGENF	$\mathcal{O}(\alpha)$	$\sim 1\%$
BHWIDE	$\mathcal{O}(\alpha)$ YFS	$\sim 0.5\%(\text{LEP1})$
BKQED	$\mathcal{O}(\alpha)$	$\sim 1\%$
MCGPJ	$\mathcal{O}(\alpha) + \text{SF}$	$< 0.2\%$

where the hard photon residuals  $\{\bar{\beta}_j\}$  are determined recursively [224, 246] and are free of all virtual and all real infrared singularities to all orders in  $\alpha$ . Introducing the result (81) into Eq. (79) and summing over the number of real photons  $n$  leads directly to master formula (69). We see that it allows for exact exclusive treatment of hard photonic effects on an event-by-event basis.

## 2.5 Monte Carlo generators

To measure the luminosity, event generators, rather than analytical calculations, are mandatory to provide theoretical results of real experimental interest. The software tools used in early measurements of the luminosity at flavour factories (and sometimes still used in recent experimental publications) include generators such as BHAGENF [248], BabaYaga v3.5 [226] and BKQED [249, 250]. These MC programs, however, are based either on a fixed NLO calculation (such as BHAGENF and BKQED) or include corrections to all orders in perturbation theory, but in the LL approximation only (like BabaYaga v3.5). Therefore the precision of these codes can be estimated to lie in the range  $0.5 \div 1\%$ , depending on the adopted experimental cuts.

The increasing precision reached on the experimental side during the last years led to the development of new dedicated theoretical tools, such as BabaYaga@NLO and MCGPJ, and the adoption of already well-tested codes, like BHWIDE, the latter extensively used at high-energy LEP/SLC colliders for the simulation of the large-angle Bhabha process. As already emphasized in Section 2.4.2, all these three codes include NLO corrections in combination with multiple photon contributions and have, therefore, a precision tag of  $\sim 0.1\%$ . As described in the following, the experiments typically use more than one generator, to keep the luminosity theoretical error under control through the comparison of independent predictions.

A list of the MC tools used in the luminosity measurement at meson factories is given in Table 3, which summarizes the main ingredients of their formulation for radiative corrections and the estimate of their theoretical accuracy.

The basic theoretical and phenomenological features of the different generators are summarized in the following.

1. BabaYaga v3.5 – It is a MC generator developed by the Pavia group at the starting of DAΦNE operation us-

ing a QED PS approach for the treatment of LL QED corrections to luminosity processes and later improved to account for the interference of radiation emitted by different charged legs in the generation of the momenta of the final-state particles. The main drawback of BabaYaga v3.5 is the absence of  $\mathcal{O}(\alpha)$  non-log contributions, resulting in a theoretical precision of  $\sim 0.5\%$  for large-angle Bhabha scattering and of about  $1\%$  for  $\gamma\gamma$  and  $\mu^+\mu^-$  final states. It is used by the CLEO-c collaboration for the study of all the three luminosity processes.

2. BabaYaga@NLO – It is the presently released version of BabaYaga, based on the matching of exact  $\mathcal{O}(\alpha)$  corrections with QED PS, as described in Section 2.4.2. The accuracy of the current version is estimated to be at  $0.1\%$  level for large-angle Bhabha scattering, two-photon and  $\mu^+\mu^-$ <sup>8</sup> production. It is presently used by the KLOE and BaBar collaborations, and under consideration by the BES-III experiment. Like BabaYaga v3.5, BabaYaga@NLO is available at the web page of the Pavia phenomenology group [www.pv.infn.it/~hepcomplex/babayaga.html](http://www.pv.infn.it/~hepcomplex/babayaga.html).
3. BHAGENF/BKQED – BKQED is the event generator developed by Berends and Kleiss and based on the classical exact NLO calculations of [249, 250] for all QED processes. It was intensively used at LEP to perform tests of QED through the analysis of the  $e^+e^- \rightarrow \gamma\gamma$  process and is adopted by the BaBar collaboration for the simulation of the same reaction. BHAGENF is a code realized by Drago and Venanzoni at the beginning of DAΦNE operation to simulate Bhabha events, adapting the calculations of [249] to include the contribution of the  $\phi$  resonance. Both generators lack the effect of HO corrections and, as such, have a precision accuracy of about  $1\%$ . The BHAGENF code is available at the web address [www.lnf.infn.it/~graziano/bhagenf/bhabha.html](http://www.lnf.infn.it/~graziano/bhagenf/bhabha.html).
4. BHWIDE – It is a MC code realized in Krakow-Knoxville at the time of LEP/SLC operation and described in [229]. In this generator exact  $\mathcal{O}(\alpha)$  corrections are matched with the resummation of soft and collinear logarithms through the YFS exponentiation approach. According to the authors the precision is estimated to be about  $0.5\%$  for c.m. energies around the  $Z$  resonance. This accuracy estimate was derived through detailed comparisons of the BHWIDE predictions with those of other LEP tools in the presence of the full set of NLO corrections, including purely weak corrections. However since the latter are phenomenologically unimportant at  $e^+e^-$  accelerators of moderately high energies and the QED theoretical ingredients of BHWIDE are very similar to the formulation of both BabaYaga@NLO and MCGPJ, one can argue that BHWIDE accuracy for physics at flavour factories is at the level of  $0.1\%$ . It is adopted by the KLOE, BaBar and

<sup>8</sup> At present, finite mass effects in the virtual corrections to  $e^+e^- \rightarrow \mu^+\mu^-$ , which should be included for precision simulations at the  $\phi$ -factories, are not included in BabaYaga@NLO.

BES collaborations. The code is available at [placzek.home.cern.ch/placzek/bhwide/](http://placzek.home.cern.ch/placzek/bhwide/).

5. MCGPJ – It is the generator developed by the Dubna-Novosibirsk collaboration and used at VEPP-2M collider. This program includes exact  $\mathcal{O}(\alpha)$  corrections supplemented with HO LL contributions related to the emission of collinear photon jets and taken into account through analytical QED collinear SF, as described in Section 2.4.2. The theoretical precision is estimated to be better than 0.2%. The generator is available at the web address [cmd.inp.nsk.su/~sibid/](http://cmd.inp.nsk.su/~sibid/).

It is worth noticing that the theoretical uncertainty of the most accurate generators based on the matching of exact NLO with LL resummation starts at the level of  $\mathcal{O}(\alpha^2)$  NNL contributions, as far as photonic corrections are concerned. Other sources of error affecting their physical precision are discussed in detail in Section 2.8

## 2.6 Numerical results

Before showing the results which enable us to settle the technical and theoretical accuracy of the generators, it is worth discussing the impact of various sources of radiative corrections implemented in the programs used in the experimental analysis. This allows one to understand which corrections are strictly necessary to achieve a precision at the per mille level for both the calculation of integrated cross section and the simulation of more exclusive distributions.

### 2.6.1 Integrated cross sections

The first set of phenomenological results about radiative corrections refer to the Bhabha cross section, as obtained by means of the code BabaYaga@NLO, according to different perturbative and precision levels. In Table 4 we show the values for the Born cross section  $\sigma_0$ , the  $\mathcal{O}(\alpha)$  PS and exact cross section,  $\sigma_\alpha^{\text{PS}}$  and  $\sigma_\alpha^{\text{NLO}}$ , respectively, as well as the LL PS cross section  $\sigma^{\text{PS}}$  and the matched cross section  $\sigma_{\text{matched}}$ . Furthermore, the cross section in the presence of the vacuum polarization correction  $\sigma_0^{\text{VP}}$  is also shown. The results correspond to the c.m. energies  $\sqrt{s} = 1, 4, 10$  GeV, and were obtained with the selection criteria of Eq. (14), but for an angular acceptance of  $55^\circ \leq \theta_\pm \leq 125^\circ$  resembling realistic data taking at meson factories. One should keep in mind that the cuts of Eq. (14) tend to single out quasi-elastic Bhabha events and that the energy of final-state electron/positron corresponds to a so-called “bare” event selection (i.e. without photon recombination), which corresponds to what is done in practice at flavour factories. In particular the rather stringent energy and acollinearity cuts enhance the impact of soft and collinear radiation with respect to a more inclusive setup.

From these cross section values, it is possible to calculate the relative effect of various corrections, namely the contribution of vacuum polarization and exact  $\mathcal{O}(\alpha)$

**Table 4.** Bhabha cross section (in nb) at meson factories according to different precision levels and using the cuts of Eq. (14), but with an angular acceptance of  $55^\circ \leq \theta_\pm \leq 125^\circ$ . The numbers in parentheses are  $1\sigma$  MC errors.

$\sqrt{s}(\text{GeV})$	1.02	4	10
$\sigma_0$	529.4631(2)	44.9619(1)	5.5026(2)
$\sigma_0^{\text{VP}}$	542.657(6)	46.9659(1)	5.85526(3)
$\sigma_{\text{NLO}}$	451.523 (6)	37.1654 (6)	4.4256 (2)
$\sigma_\alpha^{\text{PS}}$	454.503 (6)	37.4186 (6)	4.4565 (1)
$\sigma_{\text{matched}}$	455.858 (5)	37.6731 (4)	4.5046 (3)
$\sigma^{\text{PS}}$	458.437 (4)	37.8862 (4)	4.5301 (2)

QED corrections, of non-logarithmic (NLL) terms entering the  $\mathcal{O}(\alpha)$  cross section, of HO corrections in the  $\mathcal{O}(\alpha)$  matched PS scheme and finally of NNL effects beyond order  $\alpha$  largely dominated by  $\mathcal{O}(\alpha^2 L)$  contributions. The above per cent corrections are shown in Table 5 and can be derived from the cross section results of Table 4 according to the following definitions

$$\begin{aligned}
 \delta_{\text{VP}} &\equiv \frac{\sigma_0^{\text{VP}} - \sigma_0}{\sigma_0}, & \delta_\alpha &\equiv \frac{\sigma_{\text{NLO}} - \sigma_0}{\sigma_0}, \\
 \delta_\alpha^{\text{NLL}} &\equiv \frac{\sigma_{\text{NLO}} - \sigma_\alpha^{\text{PS}}}{\sigma_{\text{NLO}}}, & \delta_{\text{HO}} &\equiv \frac{\sigma_{\text{matched}} - \sigma_{\text{NLO}}}{\sigma_{\text{NLO}}}, \\
 \delta_{\alpha^2 L} &\equiv \frac{\sigma_{\text{matched}} - \sigma_{\text{NLO}} - \sigma_\alpha^{\text{PS}} + \sigma_\alpha^{\text{PS}}}{\sigma_{\text{NLO}}}.
 \end{aligned}$$

From Table 5 it can be seen that  $\mathcal{O}(\alpha)$  corrections decrease the Bhabha cross section by about 15÷17% at the  $\phi$  and  $\tau$ -charm factories, and by about 20% at the  $B$ -factories. Within the full set of  $\mathcal{O}(\alpha)$  corrections, non-log terms are of the order of 0.5%, almost independently of c.m. energy, as expected, and with a mild dependence on the angular acceptance cuts as due to box/interference contributions. The effect of HO corrections due to multiple photon emission is about 1% at the  $\phi$  and  $\tau$ -charm factories and reaches about 2% at the  $B$ -factories. The contribution of (approximate)  $\mathcal{O}(\alpha^2 L)$  corrections is at the 0.1% level, while the vacuum polarization increases the cross section by about 2% around 1 GeV, and of about 5% and 6% at 4 GeV and 10 GeV, respectively. Concerning the latter correction the non-perturbative hadronic contribution to the running of  $\alpha$  was parameterized in terms of the HADR5N routine [251,252,4] included in BabaYaga@NLO both in the LO and NLO diagrams. We have checked that the results obtained for the vacuum polarization correction in terms of the parametrization [156] agree at the  $10^{-4}$  level with those obtained with HADR5N, as shown in detail in Section 2.8. Those routines return a data driven error thus affecting the theoretical precision of the calculation of the Bhabha cross section as will be discussed in Section 2.9.

Analogous results about the size of radiative corrections to the process  $e^+e^- \rightarrow \gamma\gamma$  are given in Table 6 [253]. They were obtained using BabaYaga@NLO, according to

the experimental cuts of Eq. (15) for the c.m. energies  $\sqrt{s} = 1, 3, 10$  GeV.


**Table 5.** Relative size of different sources of correction (in per cent) to the large-angle Bhabha cross section for typical selection cuts at  $\phi$ ,  $\tau$ -charm and  $B$  factories.

$\sqrt{s}(\text{GeV})$	1.02	4.	10.
$\delta_\alpha$	-14.73	-17.32	-19.57
$\delta_\alpha^{\text{NLL}}$	-0.66	-0.68	-0.70
$\delta_{\text{HO}}$	0.97	1.35	1.79
$\delta_{\alpha^2 L}$	0.09	0.09	0.11
$\delta_{\text{VP}}$	2.43	4.46	6.03

**Table 6.** Photon pair production cross sections (in nb) at different accuracy levels and relative corrections (in per cent) for the setup of Eq. (15) and the c.m. energies  $\sqrt{s} = 1, 3, 10$  GeV.

$\sqrt{s}$ (GeV)	1	3	10
$\sigma_0$	137.53	15.281	1.3753
$\sigma_{\text{NLO}}$	129.45	14.211	1.2620
$\sigma_\alpha^{\text{PS}}$	128.55	14.111	1.2529
$\sigma_{\text{matched}}$	129.77	14.263	1.2685
$\sigma^{\text{PS}}$	128.92	14.169	1.2597
$\delta_\alpha$	-5.87	-7.00	-8.24
$\delta_\alpha^{\text{NLL}}$	0.70	0.71	0.73
$\delta_{\text{HO}}$	0.24	0.37	0.51

The numerical errors coming from the MC integration are not shown in Table 5 because they are beyond the quoted digits. From Table 5 it can be seen that the exact  $\mathcal{O}(\alpha)$  corrections lower the Born cross section by about 5.9% ( $\phi$  resonance), 7.0% (at  $\sqrt{s} = 3$  GeV) and 8.2% ( $\Upsilon$  resonance). The effect due to  $\mathcal{O}(\alpha^n L^n)$  (with  $n \geq 2$ ) terms is quantified by the contribution  $\delta_{\text{HO}}$ , which is a positive correction of about 0.2% (at the  $\phi$  resonance), 0.4% ( $\tau$ -charm factories) and 0.5% (at the  $\Upsilon$  resonance), and therefore important in the light of the aimed per mille accuracy. On the other hand also next-to-leading  $\mathcal{O}(\alpha)$  corrections, quantified by the contribution  $\delta_\alpha^{\text{NLL}}$ , are necessary at the precision level of 0.1%, since their contribution is of about 0.7% almost independently of the c.m. energy. To further corroborate the precision reached in the cross section calculation of  $e^+e^- \rightarrow \gamma\gamma$ , we also evaluated the effect due to the most important sub-leading  $\mathcal{O}(\alpha^2)$  photonic corrections and given by order  $\alpha^2 L$  contributions. It turns out that the effect due to  $\mathcal{O}(\alpha^2 L)$  corrections does not exceed the 0.05% level. Obviously, the contribution of vacuum polarization is absent in  $\gamma\gamma$  production and this is an advantage for particularly precise predictions as the uncertainty associated with the hadronic part of vacuum polarization does not affect the cross section calculation.




**Fig. 16.** Invariant mass distribution of the Bhabha process at KLOE, according to BabaYaga v3.5 (OLD), BabaYaga@NLO (NEW) and an exact NLO calculation. The inset shows the relative effect of NLO corrections, given by the difference of BabaYaga v3.5 and BabaYaga@NLO predictions. From [227].

## 2.6.2 Distributions


Besides the integrated cross section, various differential cross sections are used by the experimentalists to monitor the collider luminosity. In Fig. 16 and Fig. 17 we show two distributions which are particularly sensitive to the details of photon radiation, i.e. the  $e^+e^-$  invariant mass and acollinearity distribution, in order to quantify the size of NLO and HO corrections. The distributions are obtained according to the exact  $\mathcal{O}(\alpha)$  calculation and with the two BabaYaga versions, BabaYaga v3.5 and BabaYaga@NLO. From Fig. 16 and Fig. 17 it can be clearly seen that multiple photon corrections introduce significant deviations with respect to an  $\mathcal{O}(\alpha)$  simulation, especially in the hard tails of the distributions where they amount to several per cent. To make clearly visible the contribution of exact  $\mathcal{O}(\alpha)$  non-log terms the inset shows the relative differences between the predictions of BabaYaga v3.5 (denoted as OLD) and BabaYaga@NLO (denoted as NEW). Actually as discussed in Section 2.4.2 these differences mainly come from non-log NLO contributions and to a smaller extent from  $\mathcal{O}(\alpha^2 L)$  terms. Their effect is flat and at the level of 0.5% for the acollinearity distribution while they reach the several per cent level in the hard tail of the invariant mass distribution.

It is also worth noticing that LL radiative corrections beyond  $\alpha^2$  can be quite important for accurate simulations at least when considering differential distributions. This means that even with a complete NNLO calculation at hand it would be desirable to match such corrections with the resummation of all the remaining LL effects. In Fig. 18, the relative effect of HO corrections beyond  $\alpha^2$  dominated by the  $\alpha^3$  contributions (dashed line) is shown in comparison with that of the  $\alpha^2$  corrections (solid line) on the acollinearity distribution for the Bhabha process at DAΦNE. As it can be seen the  $\alpha^3$  effect can be as large as 10% in the phase space region of soft photon emission, corresponding to small acollinearity angles with almost back-to-back final-state fermions.






**Fig. 17.** Acollinearity distribution of the Bhabha process at KLOE, according to BabaYaga v3.5 (OLD) and BabaYaga@NLO (NEW). The inset shows the relative effect of NLO corrections, given by the difference of BabaYaga v3.5 and BabaYaga@NLO predictions. From [227].



**Fig. 18.** Relative effect of HO corrections  $\alpha^2 L^2$  and  $\alpha^n L^n$  ( $n \geq 3$ ) to the acollinearity distribution of the Bhabha process at KLOE. From [227].

Concerning the process  $e^+e^- \rightarrow \gamma\gamma$  we show in Fig. 19 the energy distribution of the most energetic photon, while the acollinearity distribution of the two most energetic photons is represented in Fig. 20. The distributions refer to exact  $\mathcal{O}(\alpha)$  corrections matched with the PS algorithm (solid line), to the exact NLO calculation (dashed line) and to all-order pure PS predictions of BabaYaga v3.5 (dashed-dotted line). In the inset of each plot, the relative effect due to multiple photon contributions ( $\delta_{\text{HO}}$ ) and non-logarithmic terms entering the improved PS algorithm ( $\delta_{\alpha}^{\text{NLL}}$ ) is also shown, according to the definitions given in Eq. (82).

For the energy distribution of the most energetic photon particularly pronounced effects due to exponentiation are present. In the statistically dominant region, HO corrections reduce the  $\mathcal{O}(\alpha)$  distribution by about 20%, while they give rise to a significant hard tail close to the energy threshold of  $0.3\sqrt{s}$  as a consequence of the higher photon multiplicity of the resummed calculation with respect to the fixed-order NLO prediction. Needless to say, the rela-



**Fig. 19.** Energy distribution of the most energetic photon in the process  $e^+e^- \rightarrow \gamma\gamma$ , according to the PS matched with  $\mathcal{O}(\alpha)$  corrections denoted as exp (solid line), the exact  $\mathcal{O}(\alpha)$  calculation (dashed line) and the pure all-order PS as in BabaYaga v3.5 (dashed-dotted line). Inset: relative effect (in per cent) of multiple photon corrections (solid line) and of non-log contributions of the matched PS algorithm (dashed line). From [253].


tive effect of multiple photon corrections below about 0.46 GeV not shown in the inset is finite but huge. This representation was chosen to make also visible in the inset the contribution of  $\mathcal{O}(\alpha)$  non-log terms, that otherwise would be hardly seen in comparison with the multiple photon corrections. Concerning the acollinearity distribution, the contribution of higher-order corrections is positive and of about 10% for quasi-back-to-back photon events, whereas it is negative and decreasing from  $\sim -30\%$  to  $\sim -10\%$  for increasing acollinearity values. As far as the contributions of non-logarithmic effects dominated by next-to-leading  $\mathcal{O}(\alpha)$  corrections are concerned, they contribute at the level of several per mille for the acollinearity distribution, while they lie in the several per cent range for the energy distribution.

As a whole, the results of the present Section emphasize that both exact  $\mathcal{O}(\alpha)$  and HO photonic corrections are necessary, as well as taking into account the running of  $\alpha$ , for a 0.1% theoretical precision in the calculation of both the cross sections and distributions.

## 2.7 Tuned comparisons

The typical procedure followed in the literature to establish the technical precision of the theoretical tools is to perform tuned comparisons between the predictions of independent programs using the same set of input parameters and experimental cuts. This strategy was initiated in the 90s during the CERN workshops for precision physics at LEP and is still in use when considering processes of interest for physics at hadron colliders demanding particularly accurate theoretical calculations. The tuning procedure is a key step in the validation of generators, because it allows to check that the different details entering the





**Fig. 20.** Acollinearity distribution for the process  $e^+e^- \rightarrow \gamma\gamma$ , according to the PS matched with  $\mathcal{O}(\alpha)$  corrections denoted as exp (solid line), the exact  $\mathcal{O}(\alpha)$  calculation (dashed line) and the pure all-order PS as in BabaYaga v3.5 (dashed-dotted line). Inset: relative effect (in per cent) of multiple photon corrections (solid line) and of non-log contributions of the matched PS algorithm (dashed line). From [253].

complex structure of the generators, e.g. implementation of radiative corrections, event selection routines, MC integration and event generation, are under control and to fix possible mistakes.

The tuned comparisons discussed in the following were performed switching off the vacuum polarization correction to the Bhabha scattering cross section. Actually, the generators implement the non-perturbative hadronic contribution to the running of  $\alpha$  according to different parameterizations, which differently affect the cross section prediction (see Section 6 for discussion). Hence, this simplification is introduced to avoid possible bias in the interpretation of the results and allows to disentangle the effect of pure QED corrections. Also, the comparisons take into account *realistic* event selection cuts, in order to provide useful results for the experiments.

The present Section is a merge of results available in the literature [227] with those of new studies. The results refer to the Bhabha process at the energies of  $\phi$ ,  $\tau$ -charm and  $B$  factories. No tuned comparisons for the two photon production process have been carried out.

### 2.7.1 $\phi$ and $\tau$ -charm factories

First we show comparisons between BabaYaga@NLO and BHWIDE according to the KLOE selection cuts of Eq. (14), considering also the angular range  $20^\circ \leq \vartheta_\pm \leq 160^\circ$  for cross section results. The predictions of the two codes are reported in Table 7 for the two acceptance cuts together with their relative deviations. As can be seen the agreement is excellent, the relative deviations being well below the 0.1%. Comparisons between BabaYaga@NLO

**Table 7.** Cross section predictions [nb] of BabaYaga@NLO and BHWIDE for the Bhabha cross section corresponding to two different angular acceptances, for the KLOE experiment at DAΦNE, and their relative differences (in per cent).

angular acceptance	BabaYaga@NLO	BHWIDE	$\delta(\%)$
$20^\circ \div 160^\circ$	6086.6(1)	6086.3(2)	0.005
$55^\circ \div 125^\circ$	455.85(1)	455.73(1)	0.030

and BHWIDE at the level of differential distributions are given in Fig. 21 and Fig. 22 where the inset shows the relative deviations between the predictions of the two codes. As can be seen there is very good agreement between the two generators, as the predicted distributions appear at a first sight almost indistinguishable. Looking in more detail, there is a relative difference of a few per mille for the acollinearity distribution (Fig. 22) and of a few per cent for the invariant mass (Fig. 21), but only in the very hard tails, where the fluctuations observed are due to limited MC statistics. These configurations however contribute to the integrated cross section negligibly, their contribution being a factor  $10^3 \div 10^4$  smaller than that around the very dominant peak regions. In fact these differences on differential distributions translate into agreement on the cross section values well below the one per mille, as shown in Table 7.

Similar tuned comparisons were performed between the results of BabaYaga@NLO, BHWIDE and MCGPJ in the presence of cuts modeling the event selection criteria of the CMD-2 experiment at the VEPP-2M collider, for a c.m. energy of  $\sqrt{s} = 900$  MeV. The cuts used in this case are

$$\begin{aligned}
 |\theta_- + \theta_+ - \pi| &\leq \Delta\theta, & 1.1 \leq (\theta_+ - \theta_- + \pi)/2 \leq \pi - 1.1, \\
 ||\phi_- + \phi_+| - \pi| &\leq 0.15, \\
 p_- \sin(\theta_-) &\geq 90 \text{ MeV}, & p_+ \sin(\theta_+) \geq 90 \text{ MeV}, \\
 (p_- + p_+)/2 &\geq 90 \text{ MeV}, & (82)
 \end{aligned}$$


where  $\theta_-$ ,  $\theta_+$  are the electron/positron polar angles, respectively,  $\phi_\pm$  their azimuthal angles, and  $p_\pm$  the moduli of their three-momenta.  $\Delta\theta$  stands for an acollinearity cut.

Fig. 23 shows the relative differences between the results of BHWIDE and MCGPJ according to the criteria of Eq. (82), as a function of the acollinearity cut  $\Delta\theta$ . The relative deviations between the results of BabaYaga@NLO and MCGPJ for the same cuts are given in Fig. 24. It can be seen that the predictions of the three generators lie within a 0.2% band with differences of  $\sim 0.3\%$  for extreme values of the acollinearity cut. This agreement can be considered satisfactory since for the acollinearity cut of real experimental interest ( $\Delta\theta \approx 0.2$  rad) the generators agree within the one per mille.


A number of comparisons was also performed for a c.m. energy of 3.5 GeV relevant to the experiments at  $\tau$ -charm factories. An example is given in Table 8 where the predictions of BabaYaga@NLO and MCGPJ are compared, using cuts similar to those of Eq. (82) and for an

**Table 8.** Cross section predictions [nb] of BabaYaga@NLO and MCGPJ for the Bhabha cross section at  $\tau$ -charm factories ( $\sqrt{s} = 3.5$  GeV) and their relative difference (in per cent).

BabaYaga@NLO	MCGPJ	$\delta(\%)$
35.20(2)	35.181(5)	0.06




**Fig. 21.** Invariant mass distribution of the Bhabha process according to BHWIDE and BabaYaga@NLO, for the KLOE experiment at DAΦNE, and relative differences of the program predictions (inset). From [227].




**Fig. 22.** Acolinearity distribution of the Bhabha process according to BHWIDE and BabaYaga@NLO, for the KLOE experiment at DAΦNE, and relative differences of the program predictions (inset). From [227].

acollinearity cut of  $\Delta\theta = 0.25$  rad. The agreement between the two codes is below one per mille. Comparisons between the two codes were also done at the level of differential cross sections, showing satisfactory agreement in the statistically relevant phase space regions. Preliminary results [254] for a c.m. energy on top of the  $J/\Psi$  resonance show good agreement between BabaYaga@NLO and BHWIDE predictions too.



**Fig. 23.** Relative differences between BHWIDE and MCGPJ Bhabha cross sections as a function of the acollinearity cut, for the CMD-2 experiment at VEPP-2M.




**Fig. 24.** Relative differences between BabaYaga@NLO and MCGPJ Bhabha cross sections as a function of the acollinearity cut, for the CMD-2 experiment at VEPP-2M.

## 2.7.2 $B$ -factories

Concerning the  $B$ -factories, a considerable effort was done to establish the level of agreement between the generators BabaYaga@NLO and BHWIDE in comparison with BabaYaga v3.5 too. This study made use of the realistic luminosity cuts quoted in Section 2.3.3 for the BaBar experiment. The cross sections predicted by BabaYaga@NLO and BHWIDE are shown in Table 9, together with the corresponding relative differences as a function of the considered angular range. The latter are also shown in Fig. 25, where the  $1\sigma$  numerical error due to MC statistics is also

**Table 9.** Cross section predictions [nb] of BabaYaga@NLO and BHWIDE for the Bhabha cross section as a function of the angular selection cuts for the BaBar experiment at PEP-II and absolute value of their relative differences (in per cent).

angular range (c.m.s.)	BabaYaga@NLO	BHWIDE	$ \delta(\%) $
$15^\circ \div 165^\circ$	119.5(1)	119.53(8)	0.025
$30^\circ \div 150^\circ$	24.17(2)	24.22(2)	0.207
$40^\circ \div 140^\circ$	11.67(3)	11.660(8)	0.086
$50^\circ \div 130^\circ$	6.31(3)	6.289(4)	0.332
$60^\circ \div 120^\circ$	1.928(2)	1.931(3)	0.141
$70^\circ \div 110^\circ$	3.554(6)	3.549(3)	0.155
$80^\circ \div 100^\circ$	0.824(2)	0.822(1)	0.243




**Fig. 25.** Relative differences between BabaYaga@NLO and BHWIDE Bhabha cross sections as a function of the angular acceptance cut for the BaBar experiment at PEP-II. From [42].

quoted. As it can be seen, the two codes agree nicely, the predictions for the central value being in general in agreement at the 0.1% level or statistically compatible whenever a two to three per mille difference is present.


To further investigate how the two generators compare with each other a number of differential cross sections was studied. The results of this study are shown in Fig. 26 and Fig. 27 for the distribution of the electron energy and the polar angle, respectively, and in Fig. 28 for the acollinearity. For both the energy and scattering angle distribution the two programs agree within the statistical errors showing deviations not above the 0.5%. For the acollinearity dependence of the cross section, BabaYaga@NLO and BHWIDE agree within  $\sim 1\%$ . Therefore, the level of the agreement between the two codes around 10 GeV is the same as that observed at the  $\phi$  factories.

The main conclusions emerging from the tuned comparisons discussed in the present Section can be summarized as follows:


- The predictions for the Bhabha cross section of most precise tools, i.e. BabaYaga@NLO, BHWIDE, MCGPJ, generally agree within 0.1%. If (slightly) larger differences are present they show up for particular tight cuts or are due to limited MC statistics. With



**Fig. 26.** Electron energy distributions according to BHWIDE, BabaYaga@NLO and BabaYaga v3.5 for the BaBar experiment at PEP-II and relative differences of the predictions of the programs. From [42].



**Fig. 27.** Electron polar angle distributions according to BHWIDE, BabaYaga@NLO and BabaYaga v3.5 for the BaBar experiment at PEP-II and relative differences of the predictions of the programs. From [42].



**Fig. 28.** Acollinearity distributions according to BHWIDE, BabaYaga@NLO and BabaYaga v3.5 for the BaBar experiment at PEP-II and relative differences of the predictions of the programs. From [42].

statistically meaningful discrepancies are observed they can be ascribed to the different theoretical recipes for the treatment of radiative corrections and their technical implementation. For example, as already emphasized, BabaYaga@NLO and BHWIDE adopt a fully factorized prescription for the matching of NLO and HO corrections, whereas MCGPJ implement some pieces of the radiative corrections in additive form. This can give rise to discrepancies between the programs predictions especially in the presence of tight cuts enhancing the effect of soft radiation. Furthermore different choices are adopted in the generators for the scale entering the collinear logarithms in HO corrections beyond  $\mathcal{O}(\alpha)$ , that are another possible source of the observed differences. To go beyond the present situation a further non trivial effort should be done by comparing, for instance, the programs in the presence of NLO corrections only (technical test) and by analyzing their different treatment of the exponentiation of soft and collinear logarithms. This would certainly shed light on the origin of the (small) discrepancies still registered at present.

- Also the distributions predicted by the generators agree well, with relative differences below the 1% level. Slightly larger discrepancies are only seen in sparsely populated phase space regions corresponding to very hard photon emission and which do not influence noticeably the luminosity measurement.

## 2.8 Theoretical accuracy

As discussed in Section 2.1, the total luminosity error crucially depends on the theoretical accuracy of the MC programs used by the experimentalists. As emphasized in Section 2.5, some of these generators like BHAGENF, BabaYaga v3.5 and BKQED miss theoretical ingredients which are unavoidable for cross section calculation with a precision at the per mille level. Therefore, they are inadequate for a highly accurate luminosity determination. BabaYaga@NLO, BHWIDE and MCGPJ include, however, both NLO and multiple photon corrections and their accuracy aims at a precision tag of 0.1%. But also these generators are affected by uncertainties which must be carefully considered in the light of the very stringent criteria of per mille accuracy. The most important components of the theoretical error of BabaYaga@NLO, BHWIDE and MCGPJ, mainly due to approximate or partially included pieces of radiative corrections, come from the following sources:

1. the non-perturbative light quark contribution to the running of  $\alpha$ . It can be reliably evaluated only using the data of the hadron cross section at low energies. Hence, the vacuum polarization correction receives a data driven error which affects in turn the prediction of the Bhabha cross section, as emphasized in Section 6.
2. the complete set of  $\mathcal{O}(\alpha^2)$  QED corrections. In spite of the impressive progress in this area, as reviewed in Section 2.3, an important piece of NNLO corrections, i.e. the exact NLO SV QED corrections to the single hard bremsstrahlung process  $e^+e^- \rightarrow e^+e^-\gamma$ , is still missing for the full  $s+t$  Bhabha process<sup>9</sup>. However, partial results obtained for  $t$ -channel small-angle Bhabha scattering [255,39] and large-angle annihilation processes are available [256,257].
3. the  $\mathcal{O}(\alpha^2)$  contribution due to real and virtual (lepton and hadron) pairs. The virtual contributions originate from the NNLO electron, heavy flavor and hadronic loop corrections discussed in Section 2.3, while the real corrections are due to the conversion of an external photon into pairs. The latter, as discussed in Section 2.3.3, gives rise to a final state with four particles, two of which to be considered as undetected to contribute to the Bhabha signature.

The uncertainty relative to the first point can be estimated by using the routines available in the literature for the calculation of the non-perturbative hadronic contribution  $\Delta\alpha_{\text{hadr}}^{(5)}(q^2)$  to the vacuum polarization. Actually these routines return, in addition to  $\Delta\alpha_{\text{hadr}}^{(5)}(q^2)$ , an error  $\delta_{\text{hadr}}$  on its value. Therefore an estimate of the induced error can be simply obtained by computing the Bhabha cross section with  $\Delta\alpha_{\text{hadr}}^{(5)}(q^2) \pm \delta_{\text{hadr}}$  and taking the difference

<sup>9</sup> As already remarked and further discussed in the following the complete calculation of the NLO corrections to hard photon emission in Bhabha scattering was performed during the completion of this report [93].



as the theoretical uncertainty due to the hadronic contribution to vacuum polarization. In Table 10, the Bhabha cross sections, as obtained in the presence of the vacuum polarization correction according to the parameterizations of [251, 252, 4] (denoted as J) and of [156] (denoted as HMNT), respectively, are shown for  $\phi$ ,  $\tau$ -charm and  $B$  factories. The applied angular cuts refer to the typically adopted acceptance  $55^\circ \leq \theta_\pm \leq 125^\circ$ .

**Table 10.** Bhabha scattering cross section in the presence of the vacuum polarization correction, according to [251, 252, 4] (J) and [156] (HMNT), at meson factories. The notation  $J_-/\text{HMNT}_-$ ,  $J/\text{HMNT}$  and  $J_+/\text{HMNT}_+$  indicates minimum, central and maximum value of the two parametrizations.

Parametrization	$\phi$	$\tau$ -charm	$B$
$J_-$	542.662(4)	46.9600(1)	5.85364(2)
J	542.662(4)	46.9658(1)	5.85529(2)
$J_+$	542.662(4)	46.9715(1)	5.85693(2)
HMNT $_-$	542.500(5)	46.9580(1)	5.85496(1)
HMNT	542.391(5)	46.9638(1)	5.85621(1)
HMNT $_+$	542.283(5)	46.9697(1)	5.85746(2)

From Table 10 it can be seen that the two treatments of  $\Delta\alpha_{\text{had}}^{(5)}(q^2)$  induce effects on the Bhabha cross section in very good agreement, the relative differences between the central values being 0.05% ( $\phi$ -factories), 0.005% ( $\tau$ -charm factories) and 0.02% ( $B$ -factories). This can be understood in terms of the dominance of  $t$ -channel exchange for large-angle Bhabha at meson factories. Indeed, the two routines provide results in excellent agreement for space-like momenta, as we explicitly checked, whereas different predictions show up for time-like momenta which, however, contribute to the Bhabha cross section only marginally. Also the spread between the minimum/maximum values and the central one as returned by the two routines agrees rather well, also a consequence of the dominance of  $t$ -channel exchange. This spread amounts to a few units in  $10^{-4}$  and is presented in detail in Table 11 in the next Section.

Concerning the second point a general strategy to evaluate the size of missing NNLO corrections consists in deriving a cross section expansion up to  $\mathcal{O}(\alpha^2)$  from the theoretical formulation implemented in the generator of interest. It can be cast in general into the following form

$$\sigma^{\alpha^2} = \sigma_{\text{SV}}^{\alpha^2} + \sigma_{\text{SV,H}}^{\alpha^2} + \sigma_{\text{HH}}^{\alpha^2}, \quad (83)$$


where in principle each of the  $\mathcal{O}(\alpha^2)$  contributions is affected by an uncertainty to be properly estimated. In Eq. (83) the first contribution is the cross section including  $\mathcal{O}(\alpha^2)$  SV corrections, whose uncertainty can be evaluated through a comparison with some of the available NNLO calculations reviewed in Section 2.3. In particular, in [227] the  $\sigma_{\text{SV}}^{\alpha^2}$  of the BabaYaga@NLO generator was compared with the calculation of photonic corrections by Penin [127, 128] and the calculations by Bonciani *et al.* [132, 133, 143,

144, 145] who computed two-loop fermionic corrections (in the one-family approximation  $N_F = 1$ ) with finite mass terms and the addition of soft bremsstrahlung and real pair contributions<sup>10</sup>. The results of such comparisons are shown in Fig. 29 and Fig. 30 for realistic cuts at the  $\phi$ -factories. In Fig. 29  $\delta\sigma$  is the difference between  $\sigma_{\text{SV}}^{\alpha^2}$  of BabaYaga@NLO and the cross sections of the two  $\mathcal{O}(\alpha^2)$  calculations, denoted as photonic (Penin) and  $N_F = 1$  (Bonciani *et al.*), as a function of the logarithm of the infrared regulator  $\epsilon$ . It can be seen that the differences are given by flat functions, demonstrating that such differences are infrared-safe, as expected, as a consequence of the universality and factorization properties of the infrared divergences. In Fig. 30,  $\delta\sigma$  is shown as a function of the logarithm of a fictitious electron mass and for a fixed value of  $\epsilon = 10^{-5}$ . Since the difference with the calculation by Penin is given by a straight line, this indicates that the soft plus virtual two-loop photonic corrections missing in BabaYaga@NLO are  $\mathcal{O}(\alpha^2 L)$  contributions, as already remarked. On the other hand, the difference with the calculation by Bonciani *et al.* is fitted by a quadratic function, showing that the electron two-loop effects missing in BabaYaga@NLO are of the order of  $\alpha^2 L^2$ . However, it is important to emphasize that, as shown in detail in [227], the sum of the relative differences with the two  $\mathcal{O}(\alpha^2)$  calculations does not exceed the  $2 \times 10^{-4}$  level for experiments at  $\phi$ - and  $B$ -factories.


The second term in Eq. (83) is the cross section containing the one-loop corrections to single hard photon emission and its uncertainty can be estimated by relying on partial results existing in the literature. Actually the exact perturbative expression of  $\sigma_{\text{SV,H}}^{\alpha^2}$  is not yet available for full  $s + t$  Bhabha scattering but using the results valid for small-angle Bhabha scattering [255, 39] and large-angle annihilation processes [256, 257] the relative uncertainty of the theoretical tools in the calculation of  $\sigma_{\text{SV,H}}^{\alpha^2}$  can be conservatively estimated at the level of 0.05%. Indeed the papers [255, 39, 256, 257] show that a YFS matching of NLO and HO corrections gives SV one-loop results for the  $t$ -channel process  $e^+e^- \rightarrow e^+e^-\gamma$  and  $s$ -channel annihilation  $e^+e^- \rightarrow f\bar{f}\gamma$  ( $f = \text{fermion}$ ) differing from the exact perturbative calculations by a few units in  $10^{-4}$  at most. This conclusion holds also when photon energy cuts are varied. It is worth noting that during the completion of the present work a complete calculation of the NLO QED corrections to hard bremsstrahlung emission in full  $s + t$  Bhabha scattering appeared in the literature [93], along the lines described in Section 2.3.2. Explicit comparisons between the results of such an exact calculation with the predictions of the most accurate MC tools according to the typical luminosity cuts used at meson factories would

<sup>10</sup> To provide meaningful results the contribution of the vacuum polarization was switched off in BabaYaga@NLO to compare with the calculation by Penin consistently. For the same reason the real soft and some pieces of virtual electron pair corrections were neglected in the comparison with the calculation by Bonciani *et al.*.





**Fig. 29.** Absolute differences (in nb) between the  $\sigma_{\text{SV}}^{\alpha^2}$  prediction of BabaYaga@NLO and the NNLO calculations of the photonic corrections [127, 128] (photonic) and of the electron loop corrections [132, 133, 143, 144, 145] ( $N_F = 1$ ) as a function of the infrared regulator  $\epsilon$  for typical KLOE cuts. From [227].



**Fig. 30.** Absolute differences (in nb) between the  $\sigma_{\text{SV}}^{\alpha^2}$  prediction of BabaYaga@NLO and the NNLO calculations of the photonic corrections [127, 128] (photonic) and of the electron loop corrections [132, 133, 143, 144, 145] ( $N_F = 1$ ) as a function of a fictitious electron mass for typical KLOE cuts. From [227].

be worthwhile to make the present error estimate related to the calculation of  $\sigma_{\text{SV,H}}^{\alpha^2}$  more robust.

The third contribution in Eq. (83) is the double hard bremsstrahlung cross section whose uncertainty can be directly evaluated by explicit comparison with the exact  $e^+e^- \rightarrow e^+e^-\gamma\gamma$  cross section. In [227] was shown that the differences between  $\sigma_{\text{HH}}^{\alpha^2}$  as in BabaYaga@NLO and the matrix element calculation which exactly describes the contribution of two hard photons are really negligible, being at the  $10^{-5}$  level.

The relative effect due to lepton ( $e, \mu, \tau$ ) and hadron ( $\pi$ ) pairs has been numerically analyzed in Section 2.3.3, in the presence of realistic selection cuts. This evaluation makes use of the complete NNLO virtual corrections combined with an exact matrix element calculation of the four-particle production processes. It supersedes previous approximate estimates which underestimated the impact of those corrections. According to this new evalua-

tion, the pair contribution dominated by the electron pair correction amounts to about 0.3% for KLOE and 0.1% for BaBar. These contributions are partially included in the BabaYaga@NLO code, as well as in other generators, through the insertion of the vacuum polarization correction in the NLO diagrams and detailed comparisons between the exact calculation and the BabaYaga@NLO predictions are in progress [258].

## 2.9 Conclusions and open issues

During the last few years a remarkable progress occurred in reducing the error of the luminosity measurements at flavour factories.

Dedicated event generators like BabaYaga@NLO and MCGPJ were developed in 2006 to provide predictions for the cross section of the large-angle Bhabha process, as well as for other QED reactions of interest, with a theoretical accuracy at the level of 0.1%. In parallel codes well-known since the time of LEP/SLC operation such as BHWIDE were extensively used by the experimentalists in data analyses. These MC programs all include, albeit according to different formulations, exact  $\mathcal{O}(\alpha)$  QED corrections matched with LL contributions describing multiple photon emission. Such ingredients together with the vacuum polarization correction are strictly necessary to achieve a physical precision down to the per mille level. Indeed when considering typical selection cuts the NLO photonic corrections amount to about 15÷20%, the vacuum polarization contributes at the several per cent level and HO effects lie between 1÷2%.

The generators mentioned are, however, affected by an uncertainty due to HO effects neglected in their formulation such as light pair corrections or exact perturbative contributions present in NNLO calculations. From this point of view the great progress in the calculation of two-loop corrections to the Bhabha scattering cross section was essential to establish the theoretical accuracy of the existing generators and will be crucial if an improvement of the precision below the one per mille will be required.

A particular effort was done to compare the predictions of the generators consistently in order to assess the technical precision obtained by the implementation of radiative corrections and related computational details. These comparisons were performed in the presence of realistic event selection criteria and at different c.m. energies. For the KLOE and CMD-2 experiments around the  $\phi$ -resonance, where the statistics of Bhabha events is the highest and the experimental luminosity error at a few per mille level, the cross section results of BabaYaga@NLO, BHWIDE and MCGPJ agree within  $\sim 0.1\%$ . If (slightly) larger discrepancies are observed, they show up only for particularly tight cuts or exclusive distributions in specific phase space regions which do not influence the luminosity determination. Very similar results were obtained for  $\tau$ -charm and  $B$ -factories. The main conclusion of the work on tuned comparisons is that the technical precision of MC programs is well under control, the discrepancies being due to different details in the treatment of the same sources

of radiative corrections and their technical implementation. For example, BabaYaga@NLO and BHWIDE adopt a fully factorized prescription for the matching of NLO and HO corrections, whereas MCGPJ implement some radiative corrections pieces in additive form. This can give rise to some discrepancies between their predictions especially in the presence of tight cuts enhancing the effect of soft radiation. Furthermore different choices are adopted in the generators for the energy scale in the treatment of HO corrections beyond  $\mathcal{O}(\alpha)$ , that are another possible source of the observed differences. To go beyond the present situation a further, non trivial effort should be done by comparing, for instance, the programs in the presence of NLO corrections only (technical test) and for the specific effect due to the exponentiation of soft and collinear logarithms. This would certainly shed light on the origin of the (minor) discrepancies still registered at present.

On the theoretical side a new exact evaluation of lepton and hadron pair corrections to the Bhabha scattering cross section was carried out taking into account realistic cuts. This calculation provides results in substantial agreement with estimates based on singlet SF but supersedes previous evaluations in the soft-photon approximation. The results of the new exact calculation were preliminarily compared with the predictions of BabaYaga@NLO, that includes the bulk of such correction (due to reducible contributions) through the insertion of the vacuum polarization correction in the NLO diagrams but neglects the effect of real pair radiation and two-loop form factors. It turns out that the error induced by the approximate treatment of pair corrections amount to a few units in  $10^{-4}$  both at KLOE and BaBar. Further work is in progress to arrive at a more solid and quantitative error estimate for these corrections when considering other selection criteria and c.m. energies too [258]. Also, the contribution induced by the uncertainty related to the non-perturbative contribution to the running of  $\alpha$  was revisited making use of and comparing the two independent parameterizations derived in [251,252,4] and [156].

A summary of the different sources of theoretical error and their relative impact on the Bhabha cross section is given in Table 11. In Table 11,  $|\delta_{\text{VP}}^{\text{err}}|$  is the error induced by the hadronic component of the vacuum polarization,  $|\delta_{\text{pairs}}^{\text{err}}|$  the error due to missing pair corrections,  $|\delta_{\text{SV}}^{\text{err}}|$  the uncertainty coming from SV NNLO corrections,  $|\delta_{\text{HH}}^{\text{err}}|$  the uncertainty in the calculation of the double hard bremsstrahlung process and  $|\delta_{\text{SV,H}}^{\text{err}}|$  the error estimate for one-loop corrections to single hard bremsstrahlung. As it can be seen pair corrections and exact NLO corrections to  $e^+e^- \rightarrow e^+e^-\gamma$  are the dominant sources of error.

The total theoretical uncertainty as obtained by summing the different contributions linearly is  $0.12 \div 0.14\%$  at the  $\phi$  factories,  $0.18\%$  at the  $\tau$ -charm factories and  $0.11 \div 0.12\%$  at the  $B$  factories. The slightly larger uncertainty at the  $\tau$ -charm factories is mainly due, as can be seen, to the pair contribution error presently based on a very preliminary evaluation and for which a deeper analysis is ongoing [258]. The total uncertainty is slightly

**Table 11.** Summary of different sources of theoretical uncertainty for the most precise generators used for luminosity measurements and the corresponding total theoretical errors for the calculation of the large-angle Bhabha cross section at meson factories.

Source of error (%)	$\phi$	$\tau$ -charm	$B$
$ \delta_{\text{VP}}^{\text{err}} $ [251,252,4]	0.00	0.01	0.03
$ \delta_{\text{VP}}^{\text{err}} $ [156]	0.02	0.01	0.02
$ \delta_{\text{SV}}^{\text{err}} $	0.02	0.02	0.02
$ \delta_{\text{HH}}^{\text{err}} $	0.00	0.00	0.00
$ \delta_{\text{SV,H}}^{\text{err}} $	0.05	0.05	0.05
$ \delta_{\text{pairs}}^{\text{err}} $	0.05	0.1	0.02
$ \delta_{\text{total}}^{\text{err}} $	$0.12 \div 0.14$	0.18	$0.11 \div 0.12$

affected by the particular choice of the routine for the calculation of  $\Delta\alpha_{\text{hadr}}^{(5)}(q^2)$ , since the two parameterizations considered here give rise to similar errors, with the exception of the  $\phi$ -factories for which the two recipes return uncertainties differing by  $2 \times 10^{-4}$ . However the “parametric” error induced by the hadronic contribution to the vacuum polarization may become a relevant source of uncertainty when considering predictions for a c.m. energy on top of and closely around very narrow resonances. For such a specific situation of interest for instance for the BES experiment, the appropriate treatment of  $\alpha$  running in the calculation of the Bhabha cross section should be deeper scrutinized because of the differences observed between the predictions for  $\Delta\alpha_{\text{hadr}}^{(5)}(q^2)$  obtained by means of the available parameterization routines (see Section 6 for a more detailed discussion).

Albeit the theoretical uncertainty quoted in Table 11 could be put on firmer grounds thanks to further studies in progress, it appears to be quite robust and sufficient for present and planned precision luminosity measurements at meson factories, the experimental error currently being about a factor of two or three larger. Adopting the strategy followed during LEP/SLC operation one could arrive at a more aggressive error estimate by summing the relative contributions in quadrature. However, for the time being this does not seem to be necessary in the light of the current experimental errors.

Concluding the precision presently reached by large-angle Bhabha programs used in the luminosity measurement at meson factories is comparable with that achieved about ten years ago for luminosity monitoring through small-angle Bhabha scattering at LEP/SLC.

Some issues are still left open. In the context of tuned comparisons no effort was done to compare the available codes for the process of photon pair production. Since it contributes relevantly to the luminosity determination and precise predictions for its cross section can be obtained by means of the codes BabaYaga@NLO and MCGPJ this work should be definitely carried out. This would lead to a better understanding of the luminosity on the experimental side. In the framework of new theoretical advances an evaluation of NNLO contributions to the process  $e^+e^- \rightarrow \gamma\gamma$  would be worthwhile to better assess

the precision of the generators which do not include for the time being such corrections exactly. More important the exact one-loop corrections to the radiative process  $e^+e^- \rightarrow e^+e^-\gamma$  should be calculated going beyond the partial results scattered in the literature (and referring to selection criteria valid for high-energy  $e^+e^-$  colliders) or limited to the soft-photon approximation<sup>11</sup>. Furthermore the radiative Bhabha process at one-loop should be evaluated taking into account the typical experimental cuts used at meson factories to get a better control of the theoretical uncertainty in the sector of NNLO corrections to Bhabha scattering. Incidentally this calculation would be also of interest for other studies at  $e^+e^-$  colliders of moderately high energy, such as the search for new physics phenomena (e.g., dark matter candidates), for which radiative Bhabha scattering is a very important background.

### 3 R measurement from energy scan

In this section we will consider some theoretical and experimental aspects of direct measurements of the process  $e^+e^- \rightarrow \text{hadrons}$  and related quantities. The cross section of  $e^+e^-$  annihilation into hadrons is required to solve various problems of particle physics and, in particular, for the evaluation of the hadronic contribution to vacuum polarisation (VP). In precision tests of the Standard Model (SM), the electromagnetic running coupling constant at the  $Z$  boson mass scale,  $\alpha(M_Z)$ , and the anomalous magnetic moment of muon,  $a^\mu = (g_\mu - 2)/2$ , are of fundamental importance. The dominant uncertainties in both quantities are due to the effects of hadronic vacuum polarisation which cannot be reliably calculated in the low energy region. Due to this fact  $\alpha(M_Z)$  remains known with the least precision among all the SM parameters in the electroweak sector. The uncertainty in the QED coupling becomes crucial in many tests of the Standard Model. In particular, the uncertainty in the hadronic VP was one of the main factors limiting the accuracy of luminosity measurements at LEP [40].

One of the main motivations for high precision studies of the hadronic cross section is the observed difference of about three standard deviations between the BNL experimental result [23] and the predicted theoretical value of  $a_\mu$  in the frame of the Standard Model. In fact, the main uncertainty of the theoretical prediction for  $a_\mu^{\text{had}}$  comes from the contribution of hadronic VP, related via a dispersion integral to the cross section of  $e^+e^-$  annihilation into hadrons.

<sup>11</sup> As already remarked in Section 2.8 during the completion of the present work a complete calculation of the NLO QED corrections to hard bremsstrahlung emission in full  $s+t$  Bhabha scattering was performed in [93]. However, explicit comparisons between the predictions of this new calculation and the corresponding results of the most precise luminosity tools are still missing and would be needed to better assess the theoretical error induced by such contribution in the calculation of the luminosity cross section.

The ratio of the radiatively-corrected hadronic cross sections to the calculated lowest-order cross section for muon pair production at the given centre-of-mass energy square  $E_{CM}^2 = s$  is usually denoted as

$$R \equiv R(s) = \frac{\sigma(e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons})}{\sigma_0(e^+e^- \rightarrow \gamma^* \rightarrow \mu^+\mu^-)}, \quad (84)$$

where  $\sigma_0(e^+e^- \rightarrow \mu^+\mu^-) \equiv 4\pi\alpha^2/3s$  is the total QED cross section of annihilation into muons at the Born level.

The  $R$  quantity has been measured in many experiments in different energy regions from the pion pair production threshold up to the  $Z$  mass. Practically all electron-positron colliders contributed to the global data set on the hadronic annihilation cross section [259]. In the high energy region ( $s \gg \Lambda_{QCD}^2$ ), away from resonances, the experimentally determined  $R$  values are in a good agreement with predictions of the perturbative QCD confirming in particular the hypothesis of three colour degrees of freedom for quarks. On the other hand, the contributions at lower energy scales involve non-perturbative QCD effects.

$R(s)$  is widely used for various QCD tests as well as for calculations of the dispersion integrals. At high energies  $R(s)$  is usually calculated within perturbative QCD, while for the low energy range the direct measurement [259, 260] of the  $e^+e^- \rightarrow \text{hadrons}$  cross section is necessary<sup>12</sup>. Matching between the two regions is performed at energies of a few GeV, where both the approaches of  $R$  definition are in a fair agreement.

The knowledge of  $R(s)$  with high accuracy is required for the evaluation of the hadronic contribution  $a_\mu^{\text{had}}$  to the anomalous magnetic moment of the muon  $(g-2)/2$  (see Ref. [262, 19, 17] and references therein). The main contribution to the value  $a_\mu^{\text{had}}$  comes from the energy range of the VEPP-2M electron-positron collider. From this energy region comes the major contribution both to the hadronic part of the anomalous magnetic moment itself and to its uncertainty [28, 19, 17]. This uncertainty is dominated by the systematic errors of the experimental values of  $R(s)$  and it is used as an input to the integral with the proper smooth kernel function [263]:

$$\begin{aligned} a_\mu^{\text{had}} &= \left(\frac{\alpha m_\mu}{3\pi}\right)^2 \int_{m_\pi^2}^{\infty} \frac{R(s)K(s)}{s^2} ds \\ &= \frac{m_\mu^2}{12\pi^3} \int_{4m_\pi^2}^{\infty} \frac{\sigma(s)K(s)}{s} ds, \end{aligned} \quad (85)$$

where  $K(s)$  is the QED kernel and  $\sigma(s)$  is the cross section of  $e^+e^- \rightarrow \text{hadrons}$ . One has to use a so-called *undressed* cross section which does not include vacuum polarisation corrections. For the best possible compilation of  $R(s)$ , data from different channels and different experiments have to be combined. For  $\sqrt{s} \leq 1.4$  GeV, the total hadronic cross section is a sum of about 25 exclusive final states. At the

<sup>12</sup> Lattice QCD computations (see, e.g., Ref. [261]) of the hadronic vacuum polarisation are in progress, but they are not yet able to provide the required precision.

present level of precision, careful treatment of the radiative corrections is necessary. As mentioned above, VP effects must be removed from the observed cross sections, but the final state radiation off hadrons should be kept. The major contribution to the uncertainty comes from the systematic error of the  $R(s)$  measurement at low energies ( $s < 2 \text{ GeV}^2$ ), which, in turn, is dominated by the systematic error of the measured cross section  $e^+e^- \rightarrow \pi^+\pi^-$ .

### 3.1 Leading-order annihilation cross sections

In this Section we present and discuss the lowest-order expressions for the processes of electron-positron annihilation into pairs of muons, pions and kaons.

For the muon production channel

$$e^-(p_-) + e^+(p_+) \rightarrow \mu^-(p'_-) + \mu^+(p'_+) \quad (86)$$

within the Standard Model at the Born level we have

$$\frac{d\sigma_0^{\mu\mu}}{d\Omega_-} = \frac{\alpha^2 \beta_\mu}{4s} (2 - \beta_\mu^2 (1 - c^2)) (1 + K_W^{\mu\mu}), \quad (87)$$

$$s = (p_- + p_+)^2 = 4\varepsilon^2, \quad c = \cos \theta_-, \quad \theta = \widehat{\mathbf{p}_- \mathbf{p}'_-},$$

where  $\beta_\mu = \sqrt{1 - m_\mu^2/\varepsilon^2}$  is the muon velocity. Small terms suppressed by the factor  $m_e^2/s$  are omitted. Here  $K_W^{\mu\mu}$  represents contributions due to  $Z$ -boson intermediate states including  $Z - \gamma$  interference, see *e.g.* Refs. [264, 265]:

$$K_W^{\mu\mu} = \frac{s^2(2 - \beta_\mu^2(1 - c^2))^{-1}}{(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} \left\{ (2 - \beta_\mu^2(1 - c^2)) \right. \\ \times \left( c_v^2 \left( 3 - 2 \frac{M_Z^2}{s} \right) + c_a^2 \right) - \frac{1 - \beta_\mu^2}{2} (c_a^2 + c_v^2) \\ \left. + c\beta_\mu \left[ 4 \left( 1 - \frac{M_Z^2}{s} \right) c_a^2 + 8c_a^2 c_v^2 \right] \right\}, \\ c_a = -\frac{1}{2 \sin 2\theta_W}, \quad c_v = c_a(1 - 4 \sin^2 \theta_W), \quad (88)$$

where  $\theta_W$  is the weak mixing angle.

The contribution of  $Z$  boson exchange is suppressed in the energy range under consideration by a factor  $s/M_Z^2$  which reaches the per mill level only at  $B$ -factories.

In the Born approximation the differential cross section of the process

$$e^+(p_+) + e^-(p_-) \rightarrow \pi^+(q_+) + \pi^-(q_-) \quad (89)$$

has the form

$$\frac{d\sigma_0^{\pi\pi}}{d\Omega}(s) = \frac{\alpha^2 \beta_\pi^3}{8s} \sin^2 \theta |F_\pi(s)|^2, \quad (90)$$

$$\beta_\pi = \sqrt{1 - m_\pi^2/\varepsilon^2}, \quad \theta = \widehat{\mathbf{p}_- \mathbf{q}_-}.$$

The pion form factor  $F_\pi(s)$  takes into account non-perturbative vertex virtual corrections due to strong interactions [266, 248]. We will call it a dressed form factor at difference to undressed form factor where the vacuum polarisation contributions are not included. We would like to emphasise

that in the approach under discussion the final state QED corrections are not included into  $F_\pi(s)$ . The form factor is extracted from the experimental data of the same process as discussed below.

The annihilation process with three pions in the final state was considered in Refs. [267, 268] including radiative corrections relevant for the energy region close to the threshold. A stand-alone Monte Carlo event generator for this channel is available [267]. This channel was also included in the MCGPJ generator [228] on the same footing as other processes under consideration.

In the case of  $K_L K_S$  meson pair production the differential cross-section in the Born approximation reads

$$\frac{d\sigma_0(s)^{K_L K_S}}{d\Omega_L} = \frac{\alpha^2 \beta_K^3}{4s} \sin^2 \theta |F_{LS}(s)|^2. \quad (91)$$

Similarly to the pion pair production, we assume here that the form factor  $F_{LS}$  includes the vacuum polarisation operator of the virtual photon. The quantity  $\beta_K = \sqrt{1 - 4m_K^2/s}$  is the  $K$ -meson c.m.s. velocity, and  $\theta$  is the angle between the directions of motion of the long living kaon and the initial electron.

In the case of  $K^+ K^-$  mesons production near threshold the Sakharov-Sommerfeld factor for the Coulomb final state interaction should be taken into account in addition:

$$\frac{d\sigma_0(s)^{K^+ K^-}}{d\Omega_-} = \frac{\alpha^2 \beta_K^3}{4s} \sin^2 \theta |F_K(s)|^2 \frac{Z}{1 - \exp(-Z)},$$

$$Z = \frac{2\pi\alpha}{v}, \quad v = 2\sqrt{\frac{s - 4m_K^2}{s}} \left( 1 + \frac{s - 4m_K^2}{s} \right)^{-1} \quad (92)$$

where  $v$  is the relative velocity of kaons [269], which has the proper non-relativistic and ultra-relativistic limits. When  $s = m_\phi^2$ , we have  $v \approx 0.5$  and the final state interaction correction gives about 5% enhancement in the cross section of charged kaon production. In principle the Sakharov-Sommerfeld factor should be added also for in the case of the pion- and muon-pair production, but as in practice the threshold regions are not studied in those cases, one can drop it without compromising the accuracy.

### 3.2 QED Radiative Corrections

One-loop radiative corrections (RC) for the processes given in Eq.(86) and Eq.(89) can be separated into two natural parts according to the parity with respect to the substitution  $c \rightarrow -c$ .

The  $c$ -even part of the one-loop soft and virtual contribution to the muon annihilation channel can be composed from the well known Dirac and Pauli form factors and the



soft photon contributions. It reads

$$\frac{d\sigma_{\mu\mu\text{-even}}^{B+S+V}}{d\Omega} = \frac{d\sigma_0^{\mu\mu}}{d\Omega} \frac{1}{|1 - \Pi(s)|^2} \left\{ 1 + \frac{2\alpha}{\pi} \left[ \left[ L - 2 + \frac{1 + \beta_\mu^2}{2\beta_\mu} l_\beta \right] \ln \frac{\Delta\varepsilon}{\varepsilon} + \frac{3}{4}(L - 1) + K_{\text{even}}^{\mu\mu} \right] \right\}, \quad (93)$$

$$K_{\text{even}}^{\mu\mu} = \frac{\pi^2}{6} - \frac{5}{4} + \rho \left( \frac{1 + \beta_\mu^2}{2\beta_\mu} - \frac{1}{2} + \frac{1}{4\beta_\mu} \right) + \ln \frac{1 + \beta_\mu}{2} \left( \frac{1}{2\beta_\mu} + \frac{1 + \beta_\mu^2}{\beta_\mu} \right) \quad (94)$$

$$- \frac{1 - \beta_\mu^2}{2\beta_\mu} \frac{l_\beta}{2 - \beta_\mu^2(1 - c^2)} + \frac{1 + \beta_\mu^2}{2\beta_\mu} \left[ \frac{\pi^2}{6} + 2\text{Li}_2 \left( \frac{1 - \beta_\mu}{1 + \beta_\mu} \right) + \rho \ln \frac{1 + \beta_\mu}{2\beta_\mu^2} + 2 \ln \frac{1 + \beta_\mu}{2} \ln \frac{1 + \beta_\mu}{2\beta_\mu^2} \right],$$

$$l_\beta = \ln \frac{1 + \beta_\mu}{1 - \beta_\mu}, \quad \rho = \ln \frac{s}{m_e^2}, \quad L = \ln \frac{s}{m_e^2},$$

where  $\Delta\varepsilon \ll \varepsilon$  is the maximum energy of soft photons in the centre-of-mass system.  $\Pi(s)$  is the vacuum polarisation operator. Here we again see that the terms with the large logarithm  $L$  dominate numerically. The superscript  $B + S + V$  stands for Born+Soft+Virtual contributions.

The  $c$ -odd part of the one-loop correction comes from the interference of Born and box Feynman diagrams and from the interference part of the soft photon emission contribution. It causes the charge asymmetry of the process:

$$\eta = \frac{d\sigma(c) - d\sigma(-c)}{d\sigma(c) + d\sigma(-c)} \neq 0. \quad (95)$$

The  $c$ -odd part of the differential cross section has the following form [237]:

$$\frac{d\sigma_{\text{odd}}^{S+V}}{d\Omega} = \frac{d\sigma_0^{\mu\mu}}{d\Omega} \frac{2\alpha}{\pi} \left[ 2 \ln \frac{\Delta\varepsilon}{\varepsilon} \ln \frac{1 - \beta_c}{1 + \beta_c} + K_{\text{odd}}^{\mu\mu} \right], \quad (96)$$

The expression for the  $c$ -odd form factor can be found in Ref. [237]. Note that in most cases the experiments have a symmetric angular acceptance, so that the odd part of the cross section does not contribute to the measured quantities.

Consider now the process of hard photon emission

$$e^+(p_+) + e^-(p_-) \rightarrow \mu^+(q_+) + \mu^-(q_-) + \gamma(k). \quad (97)$$

It was studied in detail in Ref. [237, 270, 271]. The photon energy is assumed to be larger than  $\Delta\varepsilon$ . The differential cross section has the form

$$d\sigma^{\mu\mu\gamma} = \frac{\alpha^3}{2\pi^2 s^2} R d\Gamma, \quad (98)$$

$$d\Gamma = \frac{d^3 q_- d^3 q_+ d^3 k}{q_-^0 q_+^0 k^0} \delta^{(4)}(p_+ + p_- - q_- - q_+ - k),$$

$$R = \frac{s}{16(4\pi\alpha)^3} \sum_{\text{spins}} |M|^2 = R_e + R_\mu + R_{e\mu}.$$

The quantities  $R_i$  are found directly from the matrix element, they read

$$\begin{aligned} R_e &= \frac{s}{\chi_- \chi_+} B - \frac{m_e^2}{2\chi_-^2} \frac{(t_1^2 + u_1^2 + 2m_\mu^2 s_1)}{s_1^2} \\ &\quad - \frac{m_e^2}{2\chi_+^2} \frac{(t^2 + u^2 + 2m_\mu^2 s_1)}{s_1^2} + \frac{m_\mu^2}{s_1^2} \Delta_{s_1 s_1}, \\ R_{e\mu} &= B \left( \frac{u}{\chi_- \chi'_+} + \frac{u_1}{\chi_+ \chi'_-} - \frac{t}{\chi_- \chi'_-} - \frac{t_1}{\chi_+ \chi'_+} \right) + \frac{m_\mu^2}{s s_1} \Delta_{s s_1}, \\ R_\mu &= \frac{s_1}{\chi'_- \chi'_+} B + \frac{m_\mu^2}{s^2} \Delta_{s s}, \\ B &= \frac{u^2 + u_1^2 + t^2 + t_1^2}{4 s s_1}, \\ \Delta_{s_1 s_1} &= - \frac{(t + u)^2 + (t_1 + u_1)^2}{2 \chi_- \chi_+}, \\ \Delta_{s s} &= - \frac{u^2 + t_1^2 + 2 s m_\mu^2}{2 (\chi'_-)^2} - \frac{u_1^2 + t^2 + 2 s m_\mu^2}{2 (\chi'_+)^2} \\ &\quad + \frac{1}{\chi'_- \chi'_+} (s s_1 - s^2 + t u + t_1 u_1 - 2 s m_\mu^2), \\ \Delta_{s s_1} &= \frac{s + s_1}{2} \left( \frac{u}{\chi_- \chi'_+} + \frac{u_1}{\chi_+ \chi'_-} - \frac{t}{\chi_- \chi'_-} - \frac{t_1}{\chi_+ \chi'_+} \right) + \frac{2(u - t_1)}{\chi'_-} + \frac{2(u_1 - t)}{\chi'_+}, \end{aligned}$$

where

$$\begin{aligned} s_1 &= (q_+ + q_-)^2, \quad t = -2p_- q_-, \quad t_1 = -2p_+ q_+, \\ u &= -2p_- q_+, \quad u_1 = -2p_+ q_-, \quad \chi_\pm = p_\pm k, \quad \chi'_\pm = q_\pm k. \end{aligned}$$

The bulk of the hard photon contribution comes from collinear initial state radiation. If we consider only the emission into two cones along the beam axis with restrictions

$$\widehat{p_\pm k} = \theta \leq \theta_0 \ll 1, \quad \theta_0 \gg \frac{m_e}{\varepsilon}, \quad (99)$$

we see that the corresponding contribution takes the factorized form

$$\left( \frac{d\sigma^{\mu\mu}}{d\Omega} \right)_{\text{coll}} = C_e + D_e, \quad (100)$$

$$C_e = \frac{\alpha}{2\pi} \left( \ln \frac{s}{m_e^2} - 1 \right) \int_{\Delta}^1 dx \frac{1 + (1 - x)^2}{x} A_0,$$

$$D_e = \frac{\alpha}{2\pi} \int_{\Delta}^1 dx \left\{ x + \frac{1 + (1 - x)^2}{x} \ln \frac{\theta_0^2}{4} \right\} A_0,$$

$$A_0 = \left[ \frac{d\tilde{\sigma}_0(1 - x, 1)}{d\Omega_-} + \frac{d\tilde{\sigma}_0(1, 1 - x)}{d\Omega_-} \right],$$

where the *shifted* Born differential cross section describes the process  $e^+(p_+(1 - x_2)) + e^-(p_-(1 - x_1)) \rightarrow \mu^+(q_+) +$



$\mu^-(q_-)$ ,

$$\begin{aligned} \frac{d\tilde{\sigma}_0^{\mu\mu}(z_1, z_2)}{d\Omega_-} &= \frac{\alpha^2}{4s} \\ &\times \frac{y_1[z_1^2(Y_1 - y_1c)^2 + z_2^2(Y_1 + y_1c)^2 + 8z_1z_2m_\mu^2/s]}{z_1^3z_2^3[z_1 + z_2 - (z_1 - z_2)cY_1/y_1]}, \\ y_{1,2}^2 &= Y_{1,2}^2 - \frac{4m_\mu^2}{s}, \quad Y_{1,2} = \frac{q_{-,+}^0}{\varepsilon}, \quad z_{1,2} = 1 - x_{1,2}, \\ Y_1 &= \frac{4m_\mu^2}{s}(z_2 - z_1)c \left[ 2z_1z_2 \right. \\ &\quad \left. + \sqrt{4z_1^2z_2^2 - 4(m_\mu^2/s)((z_1 + z_2)^2 - (z_1 - z_2)^2c^2)} \right]^{-1} \\ &\quad + \frac{2z_1z_2}{z_1 + z_2 - c(z_1 - z_2)}. \end{aligned} \quad (101)$$

One can recognise that the large logs related to the collinear photon emission appear in  $C_e$  in agreement with the structure function approach discussed in Section 2. In analogy to the definition of the QCD structure functions, one can move the factorized logarithmic corrections  $C_e$  to the QED electron structure function. Adding the higher-order radiative corrections in the leading logarithmic approximation to the complete one-loop result, the resulting cross section can be written as follows:

$$\begin{aligned} \frac{d\sigma^{e^+e^- \rightarrow \mu^+\mu^-}(\gamma)}{d\Omega_-} &= \int_{z_{\min}}^1 \int_{z_{\min}}^1 dz_1 dz_2 \frac{\mathcal{D}(z_1, s)\mathcal{D}(z_2, s)}{|1 - \Pi(s z_1 z_2)|^2} \\ &\times \frac{d\tilde{\sigma}_0^{\mu\mu}(z_1, z_2)}{d\Omega_-} \left( 1 + \frac{\alpha}{\pi} K_{\text{odd}}^{\mu\mu} + \frac{\alpha}{\pi} K_{\text{even}}^{\mu\mu} \right) \\ &+ \left\{ \frac{\alpha^3}{2\pi^2 s^2} \int_{\substack{k^0 > \Delta\varepsilon \\ k_{\perp} > \theta_0}} \frac{R_e|_{m_e=0}}{|1 - \Pi(s_1)|^2} \frac{d\Gamma}{d\Omega_-} + \frac{D_e}{|1 - \Pi(s_1)|^2} \right\} \\ &+ \left\{ \frac{\alpha^3}{2\pi^2 s^2} \int_{k^0 > \Delta\varepsilon} \left( \text{Re} \frac{R_{e\mu}}{(1 - \Pi(s_1))(1 - \Pi(s))^*} \right. \right. \\ &\quad \left. \left. + \frac{R_\mu}{|1 - \Pi(s)|^2} \right) \frac{d\Gamma}{d\Omega_-} + \text{Re} \frac{C_{e\mu}}{(1 - \Pi(s_1))(1 - \Pi(s))^*} \right. \\ &\quad \left. + \frac{C_\mu}{|1 - \Pi(s)|^2} \right\}, \end{aligned} \quad (102)$$


$$C_\mu = \frac{2\alpha}{\pi} \frac{d\sigma_0^{\mu\mu}}{d\Omega_-} \ln \frac{\Delta\varepsilon}{\varepsilon} \left( \frac{1 + \beta^2}{2\beta} \ln \frac{1 + \beta}{1 - \beta} - 1 \right),$$

$$C_{e\mu} = \frac{4\alpha}{\pi} \frac{d\sigma_0^{\mu\mu}}{d\Omega_-} \ln \frac{\Delta\varepsilon}{\varepsilon} \ln \frac{1 - \beta c}{1 + \beta c},$$

where  $D_e$ ,  $C_{e\mu}$  and  $C_\mu$  are compensating terms, which provide cancellation of auxiliary parameters  $\Delta$  and  $\theta_0$  inside curly brackets. In the first term, containing  $\mathcal{D}$  functions, we collect all the leading logarithmic terms. A part of non-leading terms proportional to the Born cross section is written as the  $K$ -factor. The remaining non-leading contributions are written as two additional terms. The compensating term  $D_e$  (see Eq. (100)) comes from the integration in the collinear region of hard photon emission.

The quantities  $C_\mu$  and  $C_{e\mu}$  come from the even and odd parts of the differential cross section (arising due to soft and virtual corrections), respectively. Here we consider the phase space of two ( $d\Omega_-$ ) and three ( $d\Gamma$ ) final particles as those, which already include all required experimental cuts. Using given experimental conditions one can determine the lower limits of the integration over  $z_1$  and  $z_2$  instead of the kinematic limit  $z_{\min} = 2m_\mu/(2\varepsilon - m_\mu)$ .

Matching of the complete  $\mathcal{O}(\alpha)$  RC with higher order leading logarithmic corrections can be performed in different schemes. The above approach is implemented in the MCGPJ event generator [228]. The solution of the QED evolution equations in the form of parton showers (see the Luminosity Section) matched again with the first order corrections is implemented in the BabaYaga@NLO generator [226]. Another possibility is realized in the KKMC code [272, 273] with the Yennie-Frautschi-Suura exponentiated representation of the photonic higher order corrections. A good agreement was obtained in [228] for various differential distributions for the  $\mu^+\mu^-$  channel between MCGPJ, BabaYaga@NLO, KKMC, see Fig. 31 for example.



**Fig. 31.** Comparison of the  $e^+e^- \rightarrow \mu^+\mu^-$  total cross sections computed by MCGPJ and KKMC generators versus the centre-of-mass energy.

In the ratio of the measured cross sections of annihilation into hadrons and into muons as well as in the pion form factor (106) the corrections to the initial  $e^+e^-$  state are the same for both channels and cancel out in the ratio. Nevertheless, experimental conditions and systematics are different for the muon and hadron channels. Therefore the processes have to be analysed separately and the corrections to the initial states have to be included in the analysis.

For the  $\pi^+\pi^-$  channel the complete set of  $\mathcal{O}(\alpha)$  corrections matched with the leading logarithmic electron structure functions can be found in Ref. [274, 275]. The calculation of RC was performed there within scalar QED.

Taking into account only final state corrections calculated within scalar QED, it is convenient to introduce the bare  $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$  cross section as

$$\sigma_{\pi\pi(\gamma)}^0 = \frac{\pi\alpha^2}{3s} \beta_\pi^3 |F_\pi(s)|^2 |1 - \Pi(s)|^2 \left(1 + \frac{\alpha}{\pi} \Lambda(s)\right), \quad (103)$$

where the factor  $|1 - \Pi(s)|^2$  with a polarisation operator  $\Pi(s)$  gives the effect of leptonic and hadronic vacuum polarisation. The final state radiation (FSR) correction is denoted by  $\Lambda(s)$ . For an inclusive measurement without cuts it reads [276, 277]

$$\begin{aligned} \Lambda(s) = & \frac{1 + \beta_\pi^2}{\beta_\pi} \left\{ 4\text{Li}_2\left(\frac{1 - \beta_\pi}{1 + \beta_\pi}\right) + 2\text{Li}_2\left(-\frac{1 - \beta_\pi}{1 + \beta_\pi}\right) \right. \\ & - \left[ 3\ln\left(\frac{2}{1 + \beta_\pi}\right) + 2\ln\beta_\pi \right] \ln\frac{1 + \beta_\pi}{1 - \beta_\pi} \left. \right\} - 3\ln\left(\frac{4}{1 - \beta_\pi^2}\right) \\ & - 4\ln\beta_\pi + \frac{1}{\beta_\pi^3} \left[ \frac{5}{4}(1 + \beta_\pi^2)^2 - 2 \right] \ln\frac{1 + \beta_\pi}{1 - \beta_\pi} \\ & + \frac{3}{2} \frac{1 + \beta_\pi^2}{\beta_\pi^2}. \end{aligned} \quad (104)$$

For the neutral kaon channel the corrected cross section has the form

$$\frac{d\sigma^{e^+e^- \rightarrow K_L K_S}(s)}{d\Omega_L} = \int_0^\Delta dx \frac{d\sigma_0^{e^+e^- \rightarrow K_L K_S}(s(1-x))}{d\Omega_L} F(x, s).$$

The radiation factor  $F$  takes into account radiative corrections to the initial state within the leading logarithmic approximation with exponentiation of the numerically important contribution of soft photon radiation, see Ref. [220]:

$$\begin{aligned} F(x, s) = & bx^{b-1} \left[ 1 + \frac{3}{4}b + \frac{\alpha}{\pi} \left( \frac{\pi^2}{3} - \frac{1}{2} \right) - \frac{b^2}{24} \left( \frac{1}{3}L - 2\pi^2 \right. \right. \\ & \left. \left. - \frac{37}{4} \right) \right] - b \left( 1 - \frac{x}{2} \right) + \frac{1}{8}b^2 \left[ 4(2-x)\ln\frac{1}{x} \right. \\ & \left. + \frac{1}{x}(1 + 3(1-x)^2)\ln\frac{1}{1-x} - 6 + x \right] \\ & + \left( \frac{\alpha}{\pi} \right)^2 \left\{ \frac{1}{6x} \left( x - \frac{2m_e}{\varepsilon} \right)^b \left[ (2-2x+x^2) \left( \ln\frac{sx^2}{m_e^2} - \frac{5}{3} \right) \right. \right. \\ & \left. \left. + \frac{b}{3} \left( \ln\frac{sx^2}{m_e^2} - \frac{5}{3} \right)^3 \right] + \frac{1}{2}L^2 \left[ \frac{2}{3} \frac{1-(1-x)^3}{1-x} \right. \right. \\ & \left. \left. + (2-x)\ln(1-x) + \frac{x}{2} \right] \right\} \Theta\left(x - \frac{2m_e}{\varepsilon}\right). \end{aligned}$$

Radiative corrections to the  $K^+K^-$  channel in the point-like particle approximation are the same as for the charged pion pair production (with the substitution  $m_\pi \rightarrow m_K$ ). Usually for the kaon channel we deal with the energy

range close to  $\phi$  mass. There one may choose the maximal energy of a radiated photon as

$$\omega \leq \Delta E = m_\phi - 2m_K \ll m_K, \quad \Delta \equiv \frac{\Delta E}{m_K} \approx \frac{1}{25}. \quad (105)$$

For these photons one can use the soft photon approximation.

### 3.3 Estimate of the theoretical accuracy

Let us discuss the accuracy of the description of the processes under consideration. This uncertainty can be subdivided into two major parts: the technical and the theoretical one. The first one is related to the precision in the actual computer codes and takes into account that usually the codes do not include all known contributions in the *best* approximation. The technical precision can be verified by special tests within a given code (*e.g.*, by looking at the numerical cancellation of dependence on auxiliary parameters) and tuned comparisons of different codes.

The pure theoretical precision consists of unknown higher-order corrections, of uncertainties in the treatment of photon radiation off hadrons, and of errors in the phenomenological definition of such quantities as the hadronic vacuum polarisation and the pion form factor.

Many of the codes used at meson factories do not include contributions of weak interactions even at the Born level. As discussed above, these contributions are suppressed at least by a factor  $s/M_Z^2$  and do not spoil the precision up to the energies of  $B$ -factories.

Matching the complete one-loop QED corrections with the higher-order corrections in the leading logarithmic approximations, certain parts of the second-order next-to-leading corrections are taken into account [227]. For the case of Bhabha scattering, where, *e.g.*, soft and virtual photonic corrections in  $\mathcal{O}(\alpha^2 L)$  are known analytically, one can see that their contribution in the relevant kinematic region does not exceed 0.1%<sup>13</sup>.

The uncertainty coming from the hadronic vacuum polarisation has been estimated [7] to be of order 0.04%. For measurements performed with the centre-of-mass energy of a narrow resonance (like at the  $\phi$ -meson factories) a systematic error in the determination of the resonance contribution into vacuum polarisation has to be added. The updated discussion on the accuracy of the hadronic vacuum polarisation can be found in Section 6.

Another point concerns non-leading terms of the order  $(\alpha/\pi)^2 L$ . There are several sources of them. One is the emission of two extra hard photons: one in the collinear region and the second one at large angles. Others are related to virtual and soft-photon radiative corrections to single hard photon emission and Born processes. Most of these contributions were not considered up to now. Nevertheless, we can estimate the coefficient in front of the quantity  $(\alpha/\pi)^2 L \approx 1 \cdot 10^{-4}$  to be of order of unity. That

<sup>13</sup> The proper choice of the factorisation scale [238] is important here.

was indirectly confirmed by our complete calculations of these terms for the case of small-angle Bhabha scattering.

Considering all mentioned above sources of uncertainties as independent, we conclude that the systematic error of our formulae is about 0.2% or better both for muons and pions. For the former it is a rather safe estimate. Comparisons between different codes which treat higher-order QED corrections in different ways typically show agreement at the 0.1% level. Such comparisons test the technical and partially the theoretical uncertainties. For the  $\pi^+\pi^-$  and two kaon channels, the uncertainty is enhanced due to the presence of form factors and the application of the point-like approximation for the final state hadrons.

### 3.4 Experimental Treatment of Hadronic Cross Sections and R

For older low energy data sets obtained at various  $e^+e^-$  colliders it is difficult to judge if the radiative corrections were applied correctly. As this might lead to uncontrolled systematic errors, it may be reasonable not to include them into physical analyses with the exception of the recent data from CMD-2 and SND.

Both experiments at the VEPP-2M collider in Novosibirsk have delivered independent new measurements. The covered energy range is crucial for  $(g_\mu-2)/2$  of muon and for running  $\alpha$ . For the two-pion channel  $\pi^+\pi^-$ , which gives more than 70% of the total hadronic contribution, both experiments have very good agreement over the whole energy range. The relative deviation “SND - CMD-2” is:  $(-0.3 \pm 1.6)\%$  only, revealing a good agreement within the quoted errors.

The CMD-2 and SND detectors were located in the opposite straight sections of VEPP-2M and were taking data in parallel until year 2000 when the collider was shut down to prepare for the construction of the new collider VEPP-2000. Some important features of the CMD-2 detector allowed one to select a sample of the “clean” collinear back-to-back events. The drift chamber (DC) was used to separate  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\pi^+\pi^-$  and  $K^+K^-$  events from other particles. The Z-chamber allowed to improve significantly the determination of the polar angle of charged particle tracks in DC which provided the detector acceptance with 0.2% precision. The barrel of the electromagnetic calorimeter, based on CsI crystals, helped to extract the Bhabha events from other collinear events.

The electromagnetic calorimeter of the SND detector consisted of three spherical layers of the with 1620 crystals (NaI) with a total weight of 3.6 tons. The solid angle of the calorimeter is about 90% of  $4\pi$  steradians that makes the detector practically hermetic for photons coming from the interaction point. The angular and energy resolution for photons were found to be  $1.5^\circ$  and as  $\sigma(E)/E = 4.2\%/E(\text{GeV})^{1/4}$ . More detail about CMD-2 and SND can be found elsewhere [278, 279].

#### 3.4.1 Analysis of the $\pi^+\pi^-$ channel

The pion form factor determination is crucial for a number of problems in hadronic physics. Detailed experimental data in the annihilation channel allow an accurate extraction of the  $\rho(770)$  meson parameters and of its radial excitations. Extrapolation of the energy dependence of the pion form factor to the point  $s = 0$  provides a value of the pion electromagnetic radius.


From the experimental point of view the form factor can be defined as [260]

$$|F_\pi|^2 = \frac{N_{\pi\pi}}{N_{ee} + N_{\mu\mu}} \frac{\sigma_{ee}(1 + \delta_{ee})\varepsilon_{ee} + \sigma_{\mu\mu}(1 + \delta_{\mu\mu})\varepsilon_{\mu\mu}}{\sigma_{\pi\pi}(1 + \delta_{\pi\pi})(1 + \Delta_N)(1 + \Delta_D)\varepsilon_{\pi\pi}} - \Delta_{3\pi}, \quad (106)$$

where the ratio  $N_{\pi\pi}/(N_{ee} + N_{\mu\mu})$  is derived from the observed numbers of events,  $\sigma$  are the corresponding Born cross sections,  $\delta$  are the radiative corrections (see below),  $\varepsilon$  are the detection efficiencies,  $\Delta_D$  and  $\Delta_N$  are the corrections for the pion losses caused by decays in flight and nuclear interactions respectively, and  $\Delta_{3\pi}$  is the correction for misidentification of  $\omega \rightarrow \pi^+\pi^-\pi^0$  events as  $e^+e^- \rightarrow \pi^+\pi^-$ . In the case of the latter process,  $\sigma_{\pi\pi}$  corresponds to point-like pions. Note that in ratio (106) the contribution of vacuum polarisation in the time-like region cancels out.

The data were collected in the whole energy range of the VEPP-2M and the integrated luminosity of about  $60 \text{ pb}^{-1}$  was recorded by both detectors. The beam energy was controlled and measured with a relative accuracy not worse than  $\sim 10^{-4}$  by using the resonance depolarisation method. Samples of  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\pi^+\pi^-$  events were selected for analysis. For CMD-2, the procedure of the  $e/\mu/\pi$  separation for energies  $2E \leq 600 \text{ MeV}$  was based on momentum measurement in the DC. For these energies the average difference between momenta of  $e/\mu/\pi$  is large enough with respect to the momentum resolution (Fig. 32). For energies  $2E \geq 600 \text{ MeV}$ , the energy deposition of the particles in the calorimeter is quite different and allows to separate electrons from muons and pions (Fig. 33). At the same time muons and pions cannot be separated by their energy depositions in the calorimeter. So, the ratio  $N(\mu^+\mu^-)/N(e^+e^-)$  was fixed according to QED calculations taking into account the detector acceptance and the radiative corrections. Since the selection criteria were the same for all collinear events, many effects of the detector imperfections were partly canceled out. It allowed to measure the cross section of the process  $e^+e^- \rightarrow \pi^+\pi^-$  with better precision than this of the luminosity.

The separation procedure of the  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\pi^+\pi^-$  events was based on the minimisation of the unbinned likelihood function. This method is described in detail elsewhere [280]. To simplify the error calculation of the pion form factor, the likelihood function had the global fit parameters  $(N_{ee} + N_{\mu\mu})$  and  $N_{\pi\pi}/(N_{ee} + N_{\mu\mu})$ , using  $|F_\pi(s)|^2$  given by Eq. (106). The pion form factor measured by CMD-2 has the systematic error of about 0.6-




**Fig. 32.** Two-dimensional plot of the  $e/\mu/\pi$  events. Cosmic events are distributed predominantly along a corridor which extends from the right upper to the left bottom corner. Points in this plot correspond to the momenta of particles for the beam energy of 195 MeV.

0.8% for  $\sqrt{s} \leq 1$  GeV. For energies above 1 GeV the error varies from 1.2% to 4.2%.

Since at low energies all three final states could be separated independently, the cross section of the process  $e^+e^- \rightarrow \mu^+\mu^-$  was also measured, providing an additional consistency test. The experimental value  $\sigma_{\mu\mu}^{exp}/\sigma_{\mu\mu}^{QED} = (0.980 \pm 0.013 \pm 0.007)$  is in a good agreement with the expected value of 1 within 1.4 statistical deviations.

Another method to discriminate electrons and pions from other particles was used in SND. The event separation was based on the difference in longitudinal energy deposition profiles (energy deposition in three calorimeter layers) for these particles. To use the correlations between energy depositions in calorimeter layers in the most complete way, the separation parameter was based on the neural network approach [281, 282]. The network had an input layer consisting of 7 neurons, two hidden layers with 20 neurons each and the output layer with one neuron. As the input data, the network used the energy depositions of the particles in calorimeter layers and the polar angle of one of the particles. The output signal  $R_{e/\pi}$  is a discrimination parameter between different particles. The network was tuned by using simulated events and was checked with experimental  $3\pi$  and  $e^+e^-$  events. The misidentification ratio between electrons and pions was found to be within 0.5% - 1%. The SND measured the  $e^+e^- \rightarrow \pi^+\pi^-$  cross section in the energy range 0.36 - 0.87 GeV with the systematic error of 1.3%.

The Gounaris-Sakurai (GS) parameterisation was used to fit the pion form factor. Results of the fit are shown



**Fig. 33.** Energy deposition of collinear events for the beam energy of 460 MeV.

in Fig. 34. The value of  $\chi^2$  was found to be  $\chi^2/\text{n.d.f.} = 122.9/111$  that corresponds to the probability  $P(\chi^2/\text{n.d.f.}) = 0.21$ . The average deviation between SND[281, 282] and CMD-2[283] data is:  $\Delta(\text{SND} - \text{CMD-2}) \sim (1.3 \pm 3.6)\%$  for the energy range  $\sqrt{s} \leq 0.55$  GeV and  $\Delta(\text{SND} - \text{CMD-2}) \sim (-0.53 \pm 0.34)\%$  for the energy range  $\sqrt{s} \geq 0.55$  GeV. The obtained  $\rho$  meson parameters are:  $\text{CMD-2} - M_\rho = 775.97 \pm 0.46 \pm 0.70$  MeV,  $\Gamma_\rho = 145.98 \pm 0.75 \pm 0.50$  MeV,  $\Gamma_{ee} = 7.048 \pm 0.057 \pm 0.050$  keV,  $\text{Br}(\omega \rightarrow \pi^+\pi^-) = (1.46 \pm 0.12 \pm 0.02)\%$ ;  $\text{SND} - M_\rho = 774.6 \pm 0.4 \pm 0.5$  MeV,  $\Gamma_\rho = 146.1 \pm 0.8 \pm 1.5$  MeV,  $\Gamma_{ee} = 7.12 \pm 0.02 \pm 0.11$  keV,  $\text{Br}(\omega \rightarrow \pi^+\pi^-) = (1.72 \pm 0.10 \pm 0.07)\%$ . The systematic errors were carefully studied and are listed in Table 3.4.1.

The comparison of the fitted  $\rho$  meson parameters with PDG is presented in Fig. 35. Good agreement for all parameters is observed except the branching of  $\omega$  decay into  $\pi^+\pi^-$ , where the difference  $\sim 1.6$  standard deviations is observed.


### 3.4.2 Cross section of the process $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

This channel was studied by SND in the energy range  $\sqrt{s}$  from 0.6 to 1.4 GeV [284, 285], while CMD-2 has reported results of the measurements in vicinity of the  $\omega$  [283] and  $\phi$  meson peaks [286]. For both  $\omega$  and  $\phi$  resonances CMD-2 and SND obtain consistent results for the product of the resonance branching fractions into  $e^+e^-$  and  $\pi^+\pi^-\pi^0$ , which have the world best accuracy (SND for the  $\omega$  and CMD-2 for the  $\phi$  resonance).




Sources of errors	CMD-2 $\sqrt{s} < 1$ GeV	SND	CMD-2 $1.4 > \sqrt{s} > 1$ GeV
Event separation method	0.2-0.4%	0.5%	$0.2 \div 1.5\%$
Fiducial volume	0.2%	0.8%	$0.2 \div 0.5\%$
Detection efficiency	0.2-0.5%	0.6%	$0.5 \div 2\%$
Corrections for pion losses	0.2%	0.2%	0.2%
Radiative corrections	0.3-0.4%	0.2%	$0.5 \div 2\%$
Beam energy determination	0.1-0.3%	0.3%	$0.7 \div 1.1\%$
Other corrections	0.2%	0.5%	$0.6 \div 2.2\%$
The total systematic error	0.6-0.8%	1.3 %	$1.2 \div 4.2\%$

**Table 12.** The main sources of the systematic errors for different energy regions.



**Fig. 34.** The pion form factor data and GS fit. The energy range around the  $\omega$  meson is scaled-up and presented in the inset.




**Fig. 35.** The  $\rho$  meson parameters are compared with the SND results and PDG data. The figures (left to right) are: mass (MeV), width (MeV), lepton width (keV) and branching of the  $\omega$  decay to  $\pi^+\pi^-$  (%).

CMD-2 has also performed a detailed Dalitz plot analysis of the dynamics of  $\phi$  decay to  $\pi^+\pi^-\pi^0$ . Two models of  $3\pi$  production were used: a  $\rho\pi$  mechanism and a contact amplitude. The result obtained for the ratio of the con-

tact and  $\rho\pi$  amplitudes is in good agreement with that of KLOE [287].

The systematic accuracy of the measurements is about 1.3% around  $\omega$  meson energy region, 2.5% in the  $\phi$  region, and about 5.6% for higher energies. The results of different experiments are collected in Fig. 36. The curve is the fit which takes into account the  $\rho, \omega, \phi, \omega'$  and  $\omega''$  mesons.



**Fig. 36.** Cross section of the process  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ .

### 3.4.3 Cross section of the process $e^+e^- \rightarrow 4\pi$

This cross section becomes important for energies above the  $\phi$  meson region. CMD-2 showed that the  $a_1(1260)\pi$  mechanism is dominant for the process  $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$  whereas for the channel  $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$  the intermediate state  $\omega\pi$  is additionally required to describe cross section behaviour vs energy [288]. The SND analysis confirmed these conclusions [289]. The knowledge of the dynamics of  $4\pi$  production allowed to determine with better precision the detector acceptance and efficiencies with respect to the previous measurements.

The cross section of the process  $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$  was measured with the total systematic error of 15% for CMD-2 and 7% for SND. For the channel  $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$  the systematic uncertainty was 15% and 8%, respectively. The CMD-2 reanalysis of the process  $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$  with better procedure of efficiency determination reduced




the systematic error to (5-7)% [290], and these new results are now in good agreement with other experiments.

### 3.4.4 Other modes

CMD-2 and SND have also measured the cross sections of the processes  $e^+e^- \rightarrow K_S K_L$  and  $e^+e^- \rightarrow K^+ K^-$  from threshold and up to 1.38 GeV with much better accuracy than before [291, 292, 293]. These cross sections were studied thoroughly in the vicinity of the  $\phi$  meson and their systematic errors were determined with the precision about 1.7%(SND) and 4%(CMD-2), respectively. The analysis was based on two decay modes of the  $K_S$ :  $K_S \rightarrow \pi^0 \pi^0$  and  $\pi^+ \pi^-$ . As to the process  $e^+e^- \rightarrow K^+ K^-$  the systematic uncertainty was in detail studied and was found to be 2.2%(CMD-2) and 7%(SND).

At energies  $\sqrt{s}$  above 1.04 GeV the cross sections of the processes  $e^+e^- \rightarrow K_S K_L, K^+ K^-$  were measured with the statistical accuracy about 4% and systematic errors about 4-6% and 3%, respectively, and are in good agreement with other experiments.

To summarise, the experiments performed in 1995–2000 with the CMD-2 and SND detectors at VEPP-2M allowed one to measure the exclusive cross sections of  $e^+e^-$  annihilation into hadrons in the energy range  $\sqrt{s} = 0.36 - 1.38$  GeV with larger statistics and smaller systematic errors compared to the previous experiments. The summary results of these measurements are shown in Fig. 37




**Fig. 37.** Hadronic cross sections measured by CMD-2 and SND in the whole energy range of VEPP-2M. The curve represents a smooth spline over all data.

Results of these experiments determine now the current calculation accuracy of the muon anomaly and they are one of the main sources of information about physics of vector mesons at low energies.

### 3.4.5 $R$ measurement at CLEO

Two important measurements of the  $R$  value have been recently reported by the CLEO Collaboration [294, 295].



**Fig. 38.** Comparison of the  $R$  values from CLEO (the inclusive determination) with those from Crystal Ball and BES.


In the energy range just above the open charm threshold, they collected statistics at 13 c.m. energy points from 3.97 to 4.26 GeV [295]. Hadronic cross sections in this region exhibit a rich structure reflecting the production of  $c\bar{c}$  resonances. Two independent measurements have been performed. In one of them they determined a sum of the exclusive cross sections for final states consisting of two charm mesons ( $D\bar{D}, D^*\bar{D}, D^*\bar{D}^*, D_s^+ D_s^-, D_s^{*+} D_s^-,$  and  $D_s^{*+} D_s^{*-}$ ) and for processes in which the charm-meson pair is accompanied by a pion. In the second one they measured the inclusive cross section with a systematic uncertainty between 5.2% and 6.1%. Results of both measurements are in excellent agreement, which brings an important conclusion that in this energy range the sum of the two- and three-body cross sections saturates the total cross section of charm production. In Fig. 38 the inclusive cross section measured by CLEO is compared with the previous measurements by Crystal Ball [296] and BES [297]. Good agreement is observed between data.

CLEO has also performed new measurements of  $R$  at higher energies. They collected statistics at seven c.m. energy points from 6.964 to 10.538 GeV [294] and reached a very small systematic uncertainty of 2% only. Results of their scan are presented in Fig. 39 and they are in good agreement with those of Crystal Ball [296], MD1 [298] and the previous measurement of CLEO [299]. But they are obviously inconsistent with those of the old MARK I measurement [300].

### 3.4.6 $R$ measurement at BES

Above 2 GeV the number of final states becomes too large for completely exclusive measurements, so that the values of  $R$  are measured inclusively.

In 1998, as a test of  $R$  measurement feasibility, BES took data at six c.m. energy points between 2.6 and 5.0 GeV [301]. The integrated luminosity collected at each energy point changed from 85 to 292  $\text{nb}^{-1}$ . The statistical



**Fig. 39.** Top plot: comparison of the  $R$  values from CLEO with those from MARK I, Crystal Ball and MD1; bottom plot: comparison of the new CLEO results with the QCD prediction at  $\Lambda = 0.31$  GeV.

error was around 3% per point and the systematic error ranged from 7% to 10%.

Later, in 1999 BES performed a systematic fine scan over the c.m. energy range from 2 to 4.8 GeV [297]. Data were taken at 85 energy points, with integrated luminosity varying from 9.2 to 135  $\text{nb}^{-1}$  per point. In this experiment, besides the continuum region below charmonium threshold, the high charmonium states from 3.77 to 4.50 GeV were studied [302] in detail. The statistical error was between 2% to 3%, while the systematic error ranged from 5% to 8%, due to improvement on hadronic event selection and Monte Carlo simulation of hadronization processes. The uncertainty error due to luminosity determination was from 2% to 5.8%.

More recently, in 2003 and 2004, before BES-II was shut down for upgrading to BES-III, a high-statistics data sample was taken at 2.6, 3.07 and 3.65 GeV, with integrated luminosity of 1222, 2291 and 6485  $\text{nb}^{-1}$ , respectively [303]. The systematic error, which exceeded the statistical error, was reduced to 3.5% due to further refinement on hadronic event selection and Monte Carlo simulation.

For BES-III, the main goal of the  $R$  measurement is to perform a fine scan over the whole energy region which BEPC-II can cover. For a continuum region (below

3.73 GeV), the step should not exceed 100 MeV, and for the resonance region (above 3.73 GeV), the step should be 10 to 20 MeV. Since the luminosity of BEPC-II is two orders of magnitude higher than at BEPC, the scan of the resonance region will provide precise information on the  $1^{--}$  charmonium states up to 4.6 GeV.


## 4 Radiative return

### 4.1 History and evolution of the radiative return in precision physics

The idea to use *Initial State Radiation* in order to measure hadronic cross sections from the threshold of a reaction up to the centre-of-mass (c.m.) energy of colliders with fixed energies  $\sqrt{s}$ , to reveal reaction mechanisms and to search for new mesonic states consists in exploiting the process  $e^+e^- \rightarrow \text{hadrons} + n\gamma$  to reduce the c.m. energy of the colliding electrons and positrons and consequently the mass squared  $M_{\text{had}}^2 = s - 2\sqrt{s} E_\gamma$  of the hadronic system in the final state by emitting one or more photons. The method is particularly well suited for the modern meson factories like DAΦNE (detector KLOE) running at the  $\phi$ -resonance, BEPC-II (detector BES-III), commissioned in 2008, at the  $J/\psi$  and  $\psi(2S)$ -resonances, PEP-II (detector BaBar) and KEKB (detector Belle) at the  $\Upsilon(4S)$ -resonance with their high luminosities which compensate for the  $\alpha/\pi$  suppression of the emission of a photon. DAΦNE, BEPC-II, PEP-II and KEKB cover the regions in  $M_{\text{had}}$  up to 1.02 GeV, up to 3.8 (maximally 4.6) GeV and up to 10.6 GeV, respectively (restricted for the latter actually up to 4–5 GeV if hard photons are detected). A big advantage of ISR is the low point-to-point systematic errors of the hadronic energy spectra because the luminosity, the energy of the electrons and positrons and many other contributions to the detection efficiencies are determined once for the whole spectrum. As a consequence, the overall normalization error is the same for all energies of the hadronic system. The term *Radiative return* alternately used for ISR refers to the appearance of pronounced resonances (e.g.  $\rho, \omega, \phi, J/\psi, Z$ ) with energies below the collider energy. Reviews and updated results can be found in the Proceedings of the International Workshops in Pisa (2003) [304], Nara (2004) [305], Novosibirsk (2006) [306], Pisa (2006) [307], Frascati (2008) [308], Novosibirsk (2008) [309].

Calculations of ISR date back to the sixties-seventies of the 20<sup>th</sup> century. For example, photon emission for muon pair production in electron-positron collisions has been calculated in Ref. [310], for the  $2\pi$ -final state in Ref. [311, 312], resonances ( $\rho, \omega, \phi$ ) have been implemented in Ref. [312], the excitation of  $\psi(3100)$  and  $\psi'(3700)$  in Ref. [313], and the possibility to determine the pion form factor was discussed in Ref. [314]. The application of ISR to the new high luminosity meson factories, originally aimed at the determination of the hadronic contribution to vacuum polarization, more specifically the pion form factor, has materialized in the late nineties. Early calculations of ISR for the colliders DAΦNE, PEP-II, KEKB can be found in

[315,316,317,318]. In Ref. [274] calculations of radiative corrections for pion and kaon production below energies of 2 GeV have been reported. An impressive example of ISR is the *Radiative Return* to the region of the  $Z$ -resonance at LEP-2 with collider energies around 200 GeV [319,320,321,322] (see Fig. 40).



**Fig. 40.** The reconstructed distribution of  $e^+e^- \rightarrow q\bar{q}$  events as a function of the invariant mass of the quark-antiquark system. The data has been taken for a collider energy range of 182 - 209 GeV. The prominent peak around 90 GeV represents the  $Z$ -resonance, populated after emission of photons in the initial state [320].

ISR became a powerful tool for the analysis of experiments at low and intermediate energies with the development of EVA-PHOKHARA, a Monte Carlo event generator which is user friendly, flexible and easy to implement into the software of the existing detectors [323,324,325,326,327,328,329,330,331,332,333,334,335,336,337,338,339].

EVA and its successor PHOKHARA allow to simulate the process  $e^+e^- \rightarrow \text{hadrons} + \gamma$  for a variety of exclusive final states. As a starting point EVA was constructed [323] to simulate leading order ISR and FSR for the  $\pi^+\pi^-$  channel, additional soft and collinear ISR was included on the basis of structure functions taken from [340]. Subsequently EVA was extended to include the four-pion state [324], however, without FSR. Neglecting FSR and radiative corrections, i. e. including one-photon emission from the initial state only, the cross section for the radiative return can be cast into the product of a radiator function  $H(M_{\text{had}}^2, s)$  and the cross section  $\sigma(M_{\text{had}}^2)$  for the reaction  $e^+e^- \rightarrow \text{hadrons}$ :

$$s d\sigma(e^+e^- \rightarrow \text{hadrons} \gamma)/dM_{\text{had}}^2 = \sigma(M_{\text{had}}^2) H(M_{\text{had}}^2, s).$$

However, for a precise evaluation of  $\sigma(M_{\text{had}}^2)$  the leading logarithmic approximation inherent in EVA is insufficient. Therefore, in the next step, the exact one-loop correction to the ISR process was evaluated analytically, first

for large angle photon emission [325], then for arbitrary, including collinear configurations [326]. This was and is one of the key ingredients of the generator called PHOKHARA [327,328], which also includes soft and hard real radiation, evaluated using exact matrix elements formulated within the framework of helicity amplitudes [327]. FSR in NLO approximation was addressed in [329] and incorporated in [330,331]. The importance of the charge asymmetry, a consequence of interference between ISR and FSR amplitudes, for a test of the (model dependent) description of FSR has been emphasized already in Ref.[323] and was further studied in [331].

Subsequently the generator was extended to allow the generation of many more channels with mesons, like  $K^+K^-$ ,  $K^0\bar{K}^0$ ,  $\pi^+\pi^-\pi^0$ , an improved description of the  $4\pi$  modes [332,333] and improvements in the description of FSR for the  $\mu^+\mu^-$  channel [330,331]. Also the nucleon channels  $p\bar{p}$  and  $n\bar{n}$  were implemented [334] and it was demonstrated that the separation of electric and magnetic proton form factors is feasible for a wide energy range. In fact, for the case of  $\Lambda\Lambda$  and including the polarization sensitive weak decay of  $\Lambda$  into the simulation, it was shown that even the relative phase between the two independent form factors could be disentangled [335].

Starting already with [341] various improvements were made to include the direct decay  $\phi \rightarrow \pi^+\pi^-\gamma$  as a specific aspect of FSR into the generator, a contribution of specific importance for data taken on top of the  $\phi$  resonance.

This was further pursued in the event generators FEVA and FASTERD based on EVA-PHOKHARA. FEVA includes the effects of the direct decay  $\phi \rightarrow \pi^-\pi^+\gamma$  and the decay via the  $\rho$ -resonance  $\phi \rightarrow \rho^\pm\pi^\mp \rightarrow \pi^-\pi^+\gamma$  [342,343,344]. The code FASTERD takes into account *Final State Radiation* in the frameworks of both Resonance Perturbation Theory and sQED, *Initial State Radiation* and their interference and the direct decays  $e^+e^- \rightarrow \phi \rightarrow (f_0; f_0 + \sigma)\gamma \rightarrow \pi^+\pi^-\gamma$ ,  $e^+e^- \rightarrow \phi \rightarrow \rho^\pm\pi^\mp \rightarrow \pi^+\pi^-\gamma$  and  $e^+e^- \rightarrow \rho \rightarrow \omega\pi^0 \rightarrow \pi^0\pi^0\gamma$  [345], with the possibility to include additional models.

EVA-PHOKHARA was applied for the first time to an experiment to determine the cross section  $e^+e^- \rightarrow \pi^+\pi^-$  from the reaction threshold up to the maximum energy of the collider with the detector KLOE at DAΦNE [346,347,348,349,350,351,352,353,354,355,356,357,358,359,360,361,362,363,364,365,366,367,368,369,370] (Section 4.4.1). The motivation was the determination of the  $2\pi$  final state contribution to the hadronic vacuum polarization.

The determination of the hadronic contribution to the vacuum polarization, which arises from the coupling of virtual photons to quark-antiquark-pairs  $\gamma^* \rightarrow q\bar{q} \rightarrow \gamma^*$ , is possible by measuring the cross section of electron positron annihilation into hadrons  $e^+e^- \rightarrow \gamma^* \rightarrow q\bar{q} \rightarrow \text{hadrons}$  applying the optical theorem. It is of great importance for the interpretation of the precision measurement of the anomalous magnetic moment of the muon  $a_\mu$  in Brookhaven (E821) [371,372,23,373] and for the determination of the value of the running fine structure constant at the  $Z^0$  resonance  $\alpha(m_Z^2)$ , contributing to precision tests of the *Standard model* of particle physics, see for details e. g.

Jegerlehner [374], also Davier and Marciano [375], or Teubner *et al.* [376, 18, 377]. The hadronic contribution to  $a_\mu$  below about 2 GeV is dominated by the  $2\pi$  final state, which contributes about 70 % due to the dominance of the  $\rho$ -resonance. Other major contributions come from the three- and four-pion final states. These hadronic final states constitute at present the largest error to the *Standard model* value of  $a_\mu$  and  $\alpha(m_Z^2)$  and can be determined only experimentally because calculations within perturbative QCD are unrealistic, calculations on the lattice are not yet available with necessary accuracy, and calculations in the framework of chiral perturbation theory are restricted to values close to the reaction thresholds. At energies above about 2 to 2.5 GeV perturbative QCD calculations start to become possible and reliable, see for Refs. [378, 379], also [380].

The Novosibirsk groups CMD-2 [306, 260, 291, 381, 283, 382, 383, 384, 385, 386] and SND [285, 281, 387, 282, 293, 292] measured hadronic cross sections below 1.4 GeV by changing the collider energy (*energy scan*, see preceding Section 3). The *Initial State Radiation* method used by KLOE represents an alternative, independent and complementary way to determine hadronic cross sections with different systematic errors. KLOE has determined the cross section for the reaction  $e^+e^- \rightarrow \pi^+\pi^-$  in the energy region between 0.63 and 0.958 GeV by measuring the reaction  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  and applying a radiator function based on PHOKHARA. It obtained for the hadronic contribution to the anomalous magnetic moment of the muon due to the  $2\pi$  final state  $a_\mu^{\pi\pi} = (356.7 \pm 3.1_{\text{stat+syst}}) \cdot 10^{-10}$  [368]. This value is in good agreement with those from SND [292] and CMD-2 [386]:  $a_\mu^{\pi\pi} = (361.0 \pm 5.1_{\text{stat+syst}}) \cdot 10^{-10}$  and  $a_\mu^{\pi\pi} = (361.5 \pm 3.4_{\text{stat+syst}}) \cdot 10^{-10}$ , respectively, leading to an evaluation of  $a_\mu$  [374, 375, 376, 18, 377, 29] which differs by about 3 standard deviations from the BNL experiment [23]. A different evaluation using  $\tau$ -decays into 2 pions results in a reduced discrepancy [375, 29]. The difference between  $e^+e^-$ - and  $\tau$ -based analyses is at present not understood. But one has to be aware that the evaluation with  $\tau$  data needs more theoretical input.

Soon after the application of EVA-PHOKHARA to KLOE [346] the BaBar collaboration also started the measurement of hadronic cross sections exploiting ISR [388] and using PHOKHARA (Section 4.4.2). In recent years a plethora of final states has been studied, starting with the reaction  $e^+e^- \rightarrow J/\psi \gamma \rightarrow \mu^+\mu^- \gamma$  [389]. While detecting a hard photon the upper energy for the hadron cross sections is limited to roughly 4.5 GeV. Final states with 3, 4, 5, 6 charged and neutral pions, 2 pions and 2 kaons, 4 kaons, 4 pions and 2 kaons, with a  $\phi$  and a  $f_0(980)$ ,  $J/\psi$  and 2 pions or 2 kaons, pions and  $\eta$ , kaons and  $\eta$ , but also baryonic final states with protons and antiprotons,  $\Lambda^0$  and  $\bar{\Lambda}^0$ ,  $\Lambda^0$  and  $\bar{\Sigma}^0$ ,  $\Sigma^0$  and  $\bar{\Sigma}^0$ ,  $D\bar{D}$ ,  $D\bar{D}^*$ , and  $D^*\bar{D}^*$  mesons, etc. have been investigated [390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402]. In preparation are final states with 2 pions [403] and 2 kaons. Particularly important final states are those with 4 pions (including  $\omega\pi^0$ ) which contribute significantly to the muon anomalous magnetic moment and which were poorly known before the ISR

measurements. In many of these channels additional insights into isospin symmetry breaking are expected from the comparison between  $e^+e^-$  annihilation and  $\tau$  decays.

More recently also Belle joined the ISR programme with emphasis on final states containing mesons with hidden and open charm:  $J/\psi$  and  $\psi(2S)$ ,  $D$  and  $\bar{D}$ ,  $\Lambda_c^+\Lambda_c^-$  [404, 405, 406, 407, 408, 409, 410, 411] (Section 4.4.3).

A major surprise in recent years was the opening of a totally new field of hadron spectroscopy applying ISR. Several new relatively narrow highly excited states with  $J^{PC} = 1^{--}$ , the quantum numbers of the photon, have been discovered (preliminarily denoted as  $X$ ,  $Y$ ,  $Z$ ) at the B-factories PEP-II and KEKB with the detectors BaBar and Belle, respectively. The first of them was found by BaBar in the reaction  $e^+e^- \rightarrow Y(4260) \gamma \rightarrow J/\psi \pi^+\pi^-\gamma$  [412, 413], a state around 4260 MeV with a width of 90 MeV, later confirmed by Belle via ISR [414, 405, 415] and by CLEO in an direct energy scan [416] and a radiative return [417]. Another state was detected at 2175 MeV by BaBar in the reaction  $e^+e^- \rightarrow Y(2175) \gamma \rightarrow \phi f_0(980) \gamma$  [394]. Belle found new states at 4050, 4360, 4660 MeV in the reactions  $e^+e^- \rightarrow Y \gamma \rightarrow J/\psi \pi^+\pi^-\gamma$  and  $e^+e^- \rightarrow Y \gamma \rightarrow \psi(2S) \pi^+\pi^-\gamma$  [416, 406, 405]. The structure of basically all of these new states (if they will survive) is unknown so far. Four-quark states, e. g. a  $[cs][\bar{c}\bar{s}]$  state for  $Y(4260)$ , a  $[ss][\bar{s}\bar{s}]$  state for  $Y(2175)$ , hybrid and molecular structures are discussed, see also [418].

Detailed analyses allow, in addition, also the identification of intermediate states and consequently a study of reaction mechanisms. For instance, in the case of the final state with 2 charged and 2 neutral pions ( $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma$ ) the dominating intermediate states are  $\omega\pi^0$  and  $a_1(1260)\pi$ , while  $\rho^+\rho^-$  and  $\rho^0 f_0(980)$  contribute significantly less.

Many more highly excited states with quantum numbers, different from those of the photon, have been found in decay chains of the primarily produced heavy mesons at the B-factories PEP-II and KEKB. These analyses without ISR have clearly been triggered and encouraged by the unexpected discovery of highly excited states with  $J^{PC} = 1^{--}$  found with ISR.

Also baryonic final states with protons and antiprotons,  $\Lambda^0$  and  $\bar{\Lambda}^0$ ,  $\Lambda^0$  and  $\bar{\Sigma}^0$ ,  $\Sigma^0$  and  $\bar{\Sigma}^0$  have been investigated using ISR. The effective proton form factor (see Section 4.4.2) shows a strong increase down to the  $p\bar{p}$  threshold and nontrivial structures at invariant  $p\bar{p}$  masses of 2.25 and 3.0 GeV, so far unexplained [392, 419, 420, 421, 422]. Furthermore, it should be possible to disentangle electric and magnetic form factors and thus shed light on discrepancies between different measurements of these quantities in the spacelike region [423].

Prospects for the *Radiative Return* at the Novosibirsk collider VEPP-2000 and BEPC-II are discussed in Sections 4.4.4 and 4.4.5.



## 4.2 Radiative return: A theoretical overview

### 4.2.1 Radiative return at leading order

We consider the  $e^+e^-$  annihilation process


$$e^+(p_1) + e^-(p_2) \rightarrow \text{hadrons} + \gamma(k_1), \quad (107)$$

where the real photon is emitted either from the initial (Fig. 41a) or the final state (Fig. 41b). The former process is denoted initial state radiation (ISR), while the latter is called final state radiation (FSR).

The differential rate for the ISR process can be cast into the product of a leptonic  $L^{\mu\nu}$  and a hadronic  $H^{\mu\nu}$  tensor and the corresponding factorized phase space

$$d\sigma_{\text{ISR}} = \frac{1}{2s} L_{\text{ISR}}^{\mu\nu} H_{\mu\nu} \times d\Phi_2(p_1, p_2; Q, k_1) d\Phi_n(Q; q_1, \cdot, q_n) \frac{dQ^2}{2\pi} \quad (108)$$

where  $d\Phi_n(Q; q_1, \cdot, q_n)$  denotes the hadronic  $n$ -body phase-space with all the statistical factors coming from the hadronic final state included,  $Q = \sum q_i$  and  $s = (p_1 + p_2)^2$ .



**Fig. 41.** Leading order contributions to the reaction  $e^+e^- \rightarrow h \bar{h} + \gamma$  from ISR (a) and FSR (b). Final state particles are pions or muons, or any other multihadron state. The blob represents the hadronic form factor.

For an arbitrary hadronic final state, the matrix element for the diagrams in Fig. 41a is given by

$$\begin{aligned} \mathcal{A}_{\text{ISR}}^{(0)} &= M_{\text{ISR}}^{(0)} \cdot J^{(0)} = \\ &= -\frac{e^2}{Q^2} \bar{v}(p_1) \left( \frac{\not{\epsilon}^*(k_1)[\not{k}_1 - \not{p}_1 + m_e]\gamma^\mu}{2k_1 \cdot p_1} \right. \\ &\quad \left. + \frac{\gamma^\mu[\not{p}_2 - \not{k}_1 + m_e]\not{\epsilon}^*(k_1)}{2k_1 \cdot p_2} \right) u(p_2) J_\mu^{(0)}, \quad (109) \end{aligned}$$

where  $J_\mu$  is the hadronic current. The superscript (0) indicates that the scattering amplitude is evaluated at tree-level. Summing over the polarizations of the final real photon, averaging over the polarizations of the initial  $e^+e^-$  state, and using current conservation,  $Q \cdot J^{(0)} = 0$ , the leptonic tensor

$$L_{\text{ISR}}^{(0),\mu\nu} = \overline{M_{\text{ISR}}^{(0),\mu} (M_{\text{ISR}}^{(0),\nu})^\dagger},$$

can be written in the following form:

$$\begin{aligned} L_{\text{ISR}}^{(0),\mu\nu} &= \frac{(4\pi\alpha)^2}{Q^4} \left[ \left( \frac{2m^2 q^2 (1-q^2)^2}{y_1^2 y_2^2} - \frac{2q^2 + y_1^2 + y_2^2}{y_1 y_2} \right) g^{\mu\nu} \right. \\ &\quad + \left( \frac{8m^2}{y_2^2} - \frac{4q^2}{y_1 y_2} \right) \frac{p_1^\mu p_1^\nu}{s} + \left( \frac{8m^2}{y_1^2} - \frac{4q^2}{y_1 y_2} \right) \frac{p_2^\mu p_2^\nu}{s} \\ &\quad \left. - \left( \frac{8m^2}{y_1 y_2} \right) \frac{p_1^\mu p_2^\nu + p_1^\nu p_2^\mu}{s} \right], \quad (110) \end{aligned}$$

with

$$y_i = \frac{2k_1 \cdot p_i}{s}, \quad m^2 = \frac{m_e^2}{s}, \quad q^2 = \frac{Q^2}{s}. \quad (111)$$

The leptonic tensor is symmetric under the exchange of the electron and the positron momenta. Expressing the bilinear products  $y_i$  by the photon emission angle in the c.m. frame

$$y_{1,2} = \frac{1-q^2}{2} (1 \mp \beta \cos \theta), \quad \beta = \sqrt{1-4m^2},$$

and rewriting the two-body phase space

$$d\Phi_2(p_1, p_2; Q, k_1) = \frac{1-q^2}{32\pi^2} d\Omega, \quad (112)$$

it is evident that expression (110) contains several singularities: soft singularities for  $q^2 \rightarrow 1$  and collinear singularities for  $\cos \theta \rightarrow \pm 1$ . The former are avoided by requiring a minimal photon energy. The latter are regulated by the electron mass. For  $s \gg m_e^2$ , the expression (110) can be nevertheless safely taken in the limit  $m_e \rightarrow 0$  if the emitted real photon lies far from the collinear region. In general, however, one encounters spurious singularities in the phase space integrations if powers of  $m^2 = m_e^2/s$  are prematurely neglected.

Physics of the hadronic system, whose description is model dependent, enters through the hadronic tensor

$$H_{\mu\nu} = J_\mu^{(0)} (J_\nu^{(0)})^\dagger, \quad (113)$$

where the hadronic current has to be parameterized through form factors. For two charged pions in the final state, the current

$$J_{\pi^+\pi^-}^{(0),\mu} = ie F_{2\pi}(Q^2) (q_1 - q_2)^\mu, \quad (114)$$

where  $q_1$  and  $q_2$  are the momenta of the  $\pi^+$  and  $\pi^-$ , respectively, is determined by only one function, the pion form factor  $F_{2\pi}$ . The current for the  $\mu^+\mu^-$  final state is defined obviously by QED:

$$J_{\mu^+\mu^-}^{(0),\mu} = ie \bar{u}(q_2) \gamma^\mu v(q_1). \quad (115)$$

Integrating the hadronic tensor over the hadronic phase space, one gets

$$\int H^{\mu\nu} d\Phi_n(Q; q_1, \cdot, q_n) = \frac{e^2}{6\pi} (Q^\mu Q^\nu - g^{\mu\nu} Q^2) R(Q^2), \quad (116)$$



where  $R(Q^2) = \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma_0(e^+e^- \rightarrow \mu^+\mu^-)$ , with


$$\sigma_0(e^+e^- \rightarrow \mu^+\mu^-) = \frac{4\pi\alpha^2}{3Q^2} \quad (117)$$

the tree-level muonic cross section in the limit  $Q^2 \gg 4m_\mu^2$ . After the additional integration over the photon angles, the differential distribution

$$Q^2 \frac{d\sigma_{\text{ISR}}}{dQ^2} = \frac{4\alpha^3}{3s} R(Q^2) \left\{ \frac{s^2 + Q^4}{s(s - Q^2)} (L - 1) \right\}, \quad (118)$$

with  $L = \log(s/m_e^2)$  is obtained. If instead the photon polar angle is restricted to be in the range  $\theta_{\min} < \theta < \pi - \theta_{\min}$ , this differential distribution is given by

$$Q^2 \frac{d\sigma_{\text{ISR}}}{dQ^2} = \frac{4\alpha^3}{3s} R(Q^2) \left\{ \frac{s^2 + Q^4}{s(s - Q^2)} \log \frac{1 + \cos \theta_{\min}}{1 - \cos \theta_{\min}} - \frac{s - Q^2}{s} \cos \theta_{\min} \right\}. \quad (119)$$



**Fig. 42.** The suppression of the FSR contributions to the cross section by a suitable choice of angular cuts. Results from the PHOKHARA generator. No cuts (upper curves) and suitable cuts applied (lower curves).

In the later case, the electron mass can be taken equal to zero before integration, since the collinear region is excluded by the angular cut. The contribution of the two-pion exclusive channel can be calculated from Eq.(118) and Eq.(119) with

$$R_{\pi^+\pi^-}(Q^2) = \frac{1}{4} \left( 1 - \frac{4m_\pi^2}{Q^2} \right)^{3/2} |F_{2\pi}(Q^2)|^2, \quad (120)$$

and the corresponding muonic contribution with

$$R_{\mu^+\mu^-}(Q^2) = \sqrt{1 - \frac{4m_\mu^2}{Q^2}} \left( 1 + \frac{2m_\mu^2}{Q^2} \right). \quad (121)$$

A potential complication for the measurement of the hadronic cross section from the radiative return may arise from the interplay between photons from ISR and FSR [323]. Their relative strength is strongly dependent on the photon angle relative to the beam and to the direction of the final state particles, the c.m. energy of the reaction and the invariant mass of the hadronic system. While ISR is independent of the hadronic final state, FSR is not. Moreover, it cannot be predicted from first principles and thus has to be modeled.

The amplitude for FSR (Fig. 41b) factorizes as well as

$$\mathcal{A}_{\text{FSR}}^{(0)} = M^{(0)} \cdot J_{\text{FSR}}^{(0)}, \quad (122)$$

where

$$M_\mu^{(0)} = \frac{e}{s} \bar{v}(p_1) \gamma_\mu u(p_2). \quad (123)$$

Assuming that pions are point-like, the FSR current for two pions in scalar QED (sQED) reads

$$J_{\text{FSR}}^{(0),\mu} = -i e^2 F_{2\pi}(s) \times \left[ -2g^{\mu\sigma} + (q_1 + k_1 - q_2)^\mu \frac{(2q_1 + k_1)^\sigma}{2k_1 \cdot q_1} - (q_1 - k_1 - q_2)^\mu \frac{(2q_2 + k_1)^\sigma}{2k_1 \cdot q_2} \right] \epsilon_\sigma^*(k_1). \quad (124)$$

Due to momentum conservation,  $p_1 + p_2 = q_1 + q_2 + k_1$ , and current conservation, this expression can be simplified further to

$$J_{\text{FSR}}^{(0),\mu} = 2i e^2 F_{2\pi}(s) \left[ g^{\mu\sigma} + \frac{q_2^\mu q_1^\sigma}{k_1 \cdot q_1} + \frac{q_1^\mu q_2^\sigma}{k_1 \cdot q_2} \right] \epsilon_\sigma^*(k_1). \quad (125)$$

This is the basic model adopted in EVA [323] and in PHOKHARA [325, 326, 327, 328, 329, 330, 331, 332, 335, 424] to simulate FSR off charged pions. The corresponding FSR current for muons is given by QED.

The fully differential cross section describing photon emission at leading order can be split into three pieces


$$d\sigma^{(0)} = d\sigma_{\text{ISR}}^{(0)} + d\sigma_{\text{FSR}}^{(0)} + d\sigma_{\text{INT}}^{(0)}, \quad (126)$$

which originate from the squared ISR and FSR amplitudes respectively, plus the interference term. The ISR–FSR interference, is odd under charge conjugation,


$$d\sigma_{\text{INT}}^{(0)}(q_1, q_2) = -d\sigma_{\text{INT}}^{(0)}(q_2, q_1), \quad (127)$$

and its contribution vanishes after angular integration. It gives rise, however, to a relatively large charge asymmetry and, correspondingly, to a forward–backward asymmetry

$$A(\theta) = \frac{N_h(\theta) - N_h(\pi - \theta)}{N_h(\theta) + N_h(\pi - \theta)}. \quad (128)$$



**Fig. 43.** Angular distributions of  $\pi^+$  and  $\mu^+$  at  $\sqrt{s} = 1.02$  GeV with and without FSR for different angular cuts.




**Fig. 44.** At  $\sqrt{s}=10.6$  GeV, angular distribution of  $\pi^+$  ( $\text{ISR} \simeq \text{FSR}+\text{ISR}$ ) and of  $\mu^+$  for various  $Q^2$  cuts.

The asymmetry can be used for calibration of the FSR amplitude, and fits to the angular distribution  $A(\theta)$  can test details of its model dependence [323].

The second option to disentangle ISR from FSR exploits the markedly different angular distribution of the photon from the two processes. This observation is completely general and does not rely on any model like sQED for FSR. FSR is dominated by photons collinear to the final state particles, while ISR is dominated by photons collinear to the beam direction. This suggests that we should consider only events with photons well separated from the charged final state particles and preferentially close to the beam [323, 327, 328].

This is illustrated in Fig. 42, which has been generated running PHOKHARA at leading order (LO). After introducing suitable angular cuts the contamination of events with FSR is easily reduced to less than a few per mille. The price to pay, however, is a suppression of the threshold region too. To have access to that region photons at large angles need to be tagged, and a better control of FSR is required. In Fig. 43 the angular distribution of  $\pi^+$  and  $\mu^+$  at DAΦNE energies,  $\sqrt{s} = 1.02$  GeV, are shown for



**Fig. 45.** Typical kinematic configuration of the radiative return at low and high energies.

different angular cuts. The angles are defined with respect to the incoming positron. If no angular cut is applied, the angular distribution in both cases is highly asymmetric as a consequence of the ISR–FSR interference contribution. If cuts suitable to suppress FSR, and therefore the ISR–

FSR interference, are applied, the distributions become symmetric.

Two complementary analyses are therefore possible (for details see Section 4.4.1): The *small photon angle* analysis, where the photon is untagged and FSR can be suppressed below some reasonable limit. This analysis is suitable for intermediate values of the invariant mass of the hadronic system. And the *large photon angle* analysis, giving access to the threshold region, where FSR is more pronounced and the charge asymmetry is a useful tool to probe its model dependence.

These considerations apply, however, to low beam energies, around 1 GeV. At high energies, e.g. at  $B$ -factories, very hard tagged photons are needed to access the region with low hadronic invariant masses and the hadronic system is mainly produced back to back to the hard photon. The suppression of FSR is naturally accomplished and no special angular cuts are needed. This kinematical situation is illustrated in Fig. 45. The suppression of FSR contributions to  $\pi^+\pi^-\gamma$  events is also a consequence of the rapid decrease of the form factor above 1 GeV. The relative size of FSR is of the order of a few per mil (see Fig. 44). For  $\mu^+\mu^-$  in the final state, the amount of FSR depends on the invariant mass of the muons. For  $\sqrt{Q^2} < 1$  GeV FSR is still tiny, and becomes more relevant for larger values of  $Q^2$  (see Fig. 44).

#### 4.2.2 Structure functions

The original and default version of EVA [323], simulating the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  at LO, allowed for additional initial state radiation of soft and collinear photons by the structure function (SF) method [425, 340].

In the leading logarithmic approximation (LL), the multiple emission of collinear photons off an electron is described by the convolution integral

$$\sigma(e^-X \rightarrow Y + n\gamma) = \int_0^1 dx f_e(x, Q^2) \sigma(e^-X \rightarrow Y) , \quad (129)$$

where  $f_e(x, Q^2)$  is the probability distribution of the electron with longitudinal momentum fraction  $x$ , and  $Q$  is the transverse momentum of the collinear photons. The function  $f_e(x, Q^2)$  fulfills the evolution equation

$$\begin{aligned} \frac{d}{d \log Q} f_e(x, Q^2) &= \int_x^1 \frac{dz}{z} \\ &\frac{\alpha}{\pi} \left( \frac{1+z^2}{(1-z)_+} + \frac{3}{2} \delta(1-z) \right) f_e\left(\frac{x}{z}, Q^2\right) , \end{aligned} \quad (130)$$

with initial conditions

$$f_e(x, Q^2)|_{Q^2=m_e^2} = \delta(1-x) , \quad (131)$$

and the  $+$  prescription defined as

$$\int_0^1 dx \frac{f(x)}{(1-x)_+} = \int_0^1 dx \frac{f(x) - f(1)}{(1-x)} . \quad (132)$$

The analytic solution to Eq.(130) provided in Ref. [425, 340], allows to resum soft photons to all orders in perturbation theory, accounting for large logarithms of collinear origin,  $L = \log(s/m_e^2)$ , up to two loops. The resummed cross section

$$\sigma_{\text{SF}} = \int_0^1 dx_1 \int_0^1 dx_2 D(x_1) D(x_2) \sigma_{e^+e^- \rightarrow \text{had.}+\gamma}(x_1 x_2 s) , \quad (133)$$

is thus obtained by convoluting the Born cross section of the hard photon emission process  $e^+e^- \rightarrow \text{hadrons} + \gamma$  with the SF distribution [425, 340]

$$\begin{aligned} D(x) &= [1 + \delta_N]^{1/2} \frac{\beta_e}{2} (1-x)^{\frac{\beta_e}{2}-1} \\ &\times \left\{ \frac{1}{2}(1+x^2) + \frac{1}{2} \frac{(1-x)^2}{L-1} \right. \\ &\left. + \frac{\beta_e}{8} \left( -\frac{1}{2}(1+3x^2) \log x - (1-x)^2 \right) \right\} \end{aligned} \quad (134)$$

with

$$\beta_e = 2 \frac{\alpha}{\pi} (L-1) , \quad (135)$$


and

$$\begin{aligned} \delta_N &= \frac{\alpha}{\pi} \left( \frac{3}{2}L + \frac{\pi^2}{3} - 2 \right) \\ &+ \beta_e^2 \frac{\pi^2}{8} + \left( \frac{\alpha}{\pi} \right)^2 \left( \frac{11}{8} - \frac{2\pi^2}{3} \right) L^2 . \end{aligned} \quad (136)$$

In the SF approach, the additional emission of collinear photons reduces the effective c.m. energy of the collision to  $\sqrt{x_1 x_2 s}$ . Momentum conservation is not accomplished because the extra radiation is integrated out. In order to reduce the kinematic distortion of the events, a minimal invariant mass of the observed particles, hadrons plus the tagged photon, was required in [323], introducing then a cut dependence. The SF predictions are, thus, not accurate enough for a high precision measurement of the hadronic cross section from the radiative return. A next-to-leading order (NLO) calculation is in order. The NLO prediction contains the large logarithms  $L = \log(s/m_e^2)$  at order  $\alpha^3$  and additional subleading terms, which are not taken into account within the SF method. Furthermore, it allows for a better control of the kinematical configurations because of momentum conservation. A comparison between SF and NLO predictions can be found in [327].

#### 4.2.3 Radiative return at NLO

At NLO, the  $e^+e^-$  annihilation process in Eq.(107) receives contributions from one-loop corrections and from the emission of a second real photon (see Fig. 46). After renormalization, the one-loop matrix elements still contain infrared divergences. These are canceled by adding to the one-loop corrections the two-photon contributions. There are several well established methods to perform this cancellation. The slicing method, where amplitudes are



**Fig. 46.** Typical subamplitudes describing virtual and real corrections to the reaction  $e^+e^- \rightarrow h\bar{h} + \gamma(\gamma^*)$ , where  $h = \pi^-, \mu^-$ . The superscripts (0) and (1) denote tree-level and one-loop quantities, respectively. ISR and FSR indicate that real photons are emitted from the initial or final state. The last two diagrams, with exchange of two virtual photons are non-factorizable. Permutations are omitted.

evaluated in dimensional regularization and the two photon contribution is integrated analytically in phase space for one of the photon energies up to an energy cutoff  $E_\gamma < w\sqrt{s}$  far below  $\sqrt{s}$ , was used in [325,326] to calculate the NLO corrections to ISR. The sum of the virtual and soft contributions is finite although depends on the soft photon cutoff. The contribution from the emission of the second photon with energy  $E_\gamma > w\sqrt{s}$ , which is eval-

uated numerically, completes the calculation and cancels this dependence.

The size and sign of the NLO corrections do depend on the particular choice of the experimental cuts. Hence, only using a Monte Carlo event generator one can realistically compare theoretical predictions with experiment. This is the main motivation behind PHOKHARA [325,326,327,328,329,330,331,332,335,424].

The full set of scattering amplitudes at tree-level and one-loop can be constructed from the subamplitudes depicted in Fig. 46. The one-loop amplitude with emission of a single photon is given by

$$\begin{aligned} \mathcal{A}_{1\gamma}^{(1)} &= \mathcal{A}_{\text{ISR}}^{(1)} + \mathcal{A}_{\text{FSR}}^{(1)} \\ &+ M^{(1)} \cdot J_{\text{FSR}}^{(0)} + M_{\text{ISR}}^{(0)} \cdot J^{(1)} \\ &+ \mathcal{A}_{\text{ISR}}^{2\gamma^*} + \mathcal{A}_{\text{FSR}}^{2\gamma^*}, \end{aligned} \quad (137)$$

where

$$\mathcal{A}_{\text{ISR}}^{(1)} = M_{\text{ISR}}^{(1)} \cdot J^{(0)}, \quad \mathcal{A}_{\text{FSR}}^{(1)} = M^{(0)} \cdot J_{\text{FSR}}^{(1)}, \quad (138)$$

while the amplitude with emission of two real photons reads

$$\begin{aligned} \mathcal{A}_{2\gamma}^{(0)} &= \mathcal{A}_{2\text{ISR}}^{(0)} + \mathcal{A}_{2\text{FSR}}^{(0)} \\ &+ \left( M_{\text{ISR}}^{(0)}(k_1) \cdot J_{\text{FSR}}^{(0)}(k_2) + (k_1 \leftrightarrow k_2) \right), \end{aligned} \quad (139)$$

where

$$\mathcal{A}_{2\text{ISR}}^{(0)} = M_{2\text{ISR}}^{(0)} \cdot J^{(0)}, \quad \mathcal{A}_{2\text{FSR}}^{(0)} = M^{(0)} \cdot J_{2\text{FSR}}^{(0)}. \quad (140)$$

PHOKHARA includes the full LO amplitudes and the most relevant C-even NLO contributions:

$$d\sigma = d\sigma^{(0)} + d\sigma_{\text{ISR}}^{(1)} + d\sigma_{\text{IFS}}^{(1)}, \quad (141)$$

where  $d\sigma^{(0)}$  is the LO differential cross section (Eq.(126)),

$$\begin{aligned} d\sigma_{\text{ISR}}^{(1)} &= \frac{1}{2s} \left[ 2\text{Re} \left\{ \mathcal{A}_{\text{ISR}}^{(1)} \left( \mathcal{A}_{\text{ISR}}^{(0)} \right)^\dagger \right\} d\Phi_3(p_1, p_2; q_1, q_2, k_1) \right. \\ &\quad \left. + \left| \mathcal{A}_{2\text{ISR}}^{(0)} \right|^2 d\Phi_4(p_1, p_2; q_1, q_2, k_1, k_2) \right], \end{aligned} \quad (142)$$

is the second order radiative corrections to ISR, and

$$\begin{aligned} d\sigma_{\text{IFS}}^{(1)} &= \frac{1}{2s} \left[ 2\text{Re} \left\{ M_{\text{ISR}}^{(0)} \cdot J^{(1)} \left( \mathcal{A}_{\text{ISR}}^{(0)} \right)^\dagger \right. \right. \\ &\quad \left. \left. + M^{(1)} \cdot J_{\text{FSR}}^{(0)} \left( \mathcal{A}_{\text{FSR}}^{(0)} \right)^\dagger \right\} d\Phi_3(p_1, p_2; q_1, q_2, k_1) \right. \\ &\quad \left. + \left( \left| M_{\text{ISR}}^{(0)}(k_1) \cdot J_{\text{FSR}}^{(0)}(k_2) \right|^2 + (k_1 \leftrightarrow k_2) \right) \right. \\ &\quad \left. \times d\Phi_4(p_1, p_2; q_1, q_2, k_1, k_2) \right], \end{aligned} \quad (143)$$

is the contribution of events with simultaneous emission of one photon from the initial state and another one from the final state, together with ISR amplitudes with final state



one-loop vertex corrections, and FSR amplitudes with initial state one-loop vertex corrections. We denote these corrections as IFS.

Vacuum polarization corrections are included in the hadronic currents multiplicatively:

$$\begin{aligned} J^{(i)} &\rightarrow C_{\text{VP}}(Q^2) J^{(i)} , \\ J_{\text{FSR}}^{(i)}(k_j) &\rightarrow C_{\text{VP}}((Q + k_j)^2) J_{\text{FSR}}^{(i)}(k_j) , \\ J_{2\text{FSR}}^{(0)} &\rightarrow C_{\text{VP}}(s) J_{2\text{FSR}}^{(0)} . \end{aligned} \quad (144)$$

The virtual photon propagator is included by definition in the leptonic subamplitudes  $M^{(i)}$  and  $M_{\text{ISR}}^{(i)}$ , and  $M_{2\text{ISR}}^{(0)}$ :

$$\begin{aligned} M^{(i)} &\sim \frac{1}{s} , \\ M_{\text{ISR}}^{(i)}(k_j) &\sim \frac{1}{(p_1 + p_2 + k_j)^2} \\ M_{2\text{ISR}}^{(0)} &\sim \frac{1}{Q^2} . \end{aligned} \quad (145)$$

Neither diagrams where two photons are emitted from the final state nor final-state vertex corrections with associated real radiation from the final state are included. These constitute radiative corrections to FSR and will give non-negligible contributions only for those cases, where at least one photon is collinear with one of the final state particles. Box diagrams with associated real radiation from the initial- or the final-state leptons, as well as pentagon diagrams, are also neglected. As long as one considers charge symmetric observables only, their contribution is neither divergent in the soft nor the collinear limit and is thus of order  $\alpha/\pi$  without any enhancement factor. One should stress that PHOKHARA includes only C-even gauge invariant sets of diagrams at NLO. The missing contributions are either small, or do not contribute for charge symmetric cuts. Their implementation is, however, underway.

The calculation of the NLO corrections to ISR,  $d\sigma_{\text{ISR}}^{(1)}$ , is independent of the final state. These corrections are included by default for all the final state channels implemented in PHOKHARA, and can be easily added for any other new channel, with the sole substitution of the tree-level final state current. The radiative corrections of the IFS process depend on the final state. The latest version of PHOKHARA (version 6.0 [335]) includes these corrections for two charged pions, kaons and muons.

#### Virtual and soft corrections to ISR

The virtual and soft QED corrections to ISR in  $e^+e^-$  annihilation were originally implemented in PHOKHARA through the leptonic tensor. For future applications, however, it will be more convenient to implement those corrections directly at the amplitude level (in preparation).

In terms of subamplitudes, the leptonic tensor is given by

$$\begin{aligned} L_{\text{ISR}}^{\mu\nu} &= L_{\text{ISR}}^{(0),\mu\nu} + M_{\text{ISR}}^{(1),\mu} \left( M_{\text{ISR}}^{(0),\nu} \right)^\dagger + M_{\text{ISR}}^{(0),\mu} \left( M_{\text{ISR}}^{(1),\nu} \right)^\dagger \\ &\quad + \frac{1}{2(2\pi)^{d-1}} \int_0^{w\sqrt{s}} E^{d-3} dE d\Omega M_{2\text{ISR}}^{(0),\mu} \left( M_{2\text{ISR}}^{(0),\nu} \right)^\dagger , \end{aligned} \quad (146)$$

where  $E$  and  $\Omega$  are the energy and the solid angle of the soft photon, respectively, and  $d = 4 - 2\epsilon$  is the number of dimensions in dimensional regularization. The leptonic tensor has the following general form:

$$\begin{aligned} L_{\text{ISR}}^{\mu\nu} &= \frac{(4\pi\alpha)^2}{Q^4} \left[ a_{00} g^{\mu\nu} + a_{11} \frac{p_1^\mu p_1^\nu}{s} + a_{22} \frac{p_2^\mu p_2^\nu}{s} \right. \\ &\quad \left. + a_{12} \frac{p_1^\mu p_2^\nu + p_2^\mu p_1^\nu}{s} + i\pi a_{-1} \frac{p_1^\mu p_2^\nu - p_2^\mu p_1^\nu}{s} \right] , \end{aligned} \quad (147)$$

where the scalar coefficients  $a_{ij}$  and  $a_{-1}$  allow the following expansion:

$$a_{ij} = a_{ij}^{(0)} + \frac{\alpha}{\pi} a_{ij}^{(1)} , \quad a_{-1} = \frac{\alpha}{\pi} a_{-1}^{(1)} . \quad (148)$$

The imaginary antisymmetric piece, which is proportional to  $a_{-1}$ , appears for the first time at second order, and is particularly relevant for those cases where the hadronic current receives contributions from different amplitudes with non-trivial relative phases. This is possible, e.g. for final states with three or more mesons or for  $p\bar{p}$  production.

The LO coefficients  $a_{ij}^{(0)}$  can be read directly from Eq.(110)

$$\begin{aligned} a_{00}^{(0)} &= \frac{2m^2 q^2 (1 - q^2)^2}{y_1^2 y_2^2} - \frac{2q^2 + y_1^2 + y_2^2}{y_1 y_2} , \\ a_{11}^{(0)} &= \frac{8m^2}{y_2^2} - \frac{4q^2}{y_1 y_2} , \quad a_{22}^{(0)} = a_{11}^{(0)}(y_1 \leftrightarrow y_2) , \\ a_{12}^{(0)} &= -\frac{8m^2}{y_1 y_2} . \end{aligned} \quad (149)$$

The NLO coefficients  $a_{ij}^{(1)}$  and  $a_{-1}^{(1)}$  are obtained by combining the one-loop and the soft contributions. It is convenient to split the coefficients  $a_{ij}^{(1)}$  into a part that contributes at large photon angles and a part proportional to  $m_e^2$  and  $m_e^4$  which is relevant only in the collinear regions. These coefficients are denoted by  $a_{ij}^{(1,0)}$  and  $a_{ij}^{(1,m)}$  respectively:

$$\begin{aligned} a_{ij}^{(1)} &= a_{ij}^{(0)} \left[ -\log(4w^2)[1 + \log(m^2)] \right. \\ &\quad \left. - \frac{3}{2} \log\left(\frac{m^2}{q^2}\right) - 2 + \frac{\pi^2}{3} \right] + a_{ij}^{(1,0)} + a_{ij}^{(1,m)} . \end{aligned} \quad (150)$$

The factor proportional to the LO coefficients  $a_{ij}^{(0)}$  contains usual soft and collinear logarithms. The quantity  $w$  denotes the dimensionless value of the soft photon energy cutoff:  $E_\gamma < w\sqrt{s}$ . It is enough to present four out of the

five coefficients because exchanging the positron with the electron momenta leads to the symmetric relationship

$$a_{22}^{(1)} = a_{11}^{(1)}(y_1 \leftrightarrow y_2) . \quad (151)$$

The large-angle contributions have been calculated in Ref. [325]. The coefficient proportional to  $g^{\mu\nu}$  reads

$$a_{00}^{(1,0)} = \frac{1}{y_1 y_2} \left[ -\frac{q^2(1-q^2)}{2} - y_1 y_2 - \left[ q^2 + \frac{2y_1 y_2}{1-q^2} \right] \log(q^2) \right. \\ \left. + \left\{ \frac{y_1}{2} \left[ 4 - y_1 - \frac{3(1+q^2)}{1-y_2} \right] \log\left(\frac{y_1}{q^2}\right) \right. \right. \\ \left. \left. - \left[ 1 + (1-y_2)^2 + \frac{y_1 q^2}{y_2} \right] L(y_1) + (y_1 \leftrightarrow y_2) \right\} \right] . \quad (152)$$

The coefficient in front of the tensor structure  $p_1^\mu p_1^\nu$ , is given by

$$a_{11}^{(1,0)} = \frac{1}{y_1 y_2} \left[ (1+q^2)^2 \left( \frac{1}{1-y_1} - \frac{1}{1-q^2} \right) - \frac{4(1-y_2)y_1}{1-q^2} \right. \\ \left. - \frac{2q^2}{1-q^2} \left[ (1-y_2) \left( \frac{1}{y_2} + \frac{q^2}{y_1} + \frac{2y_1}{1-q^2} \right) \right. \right. \\ \left. \left. + \frac{2q^2}{1-q^2} \right] \log(q^2) - q^2 \left[ 1 + \frac{2}{y_2} \right] \log\left(\frac{y_1}{q^2}\right) \right. \\ \left. - q^2 \left[ \frac{(2-3y_1)(1-y_2)^2}{y_1(1-y_1)^2} \right] \log\left(\frac{y_2}{q^2}\right) \right. \\ \left. - 2q^2 \left[ 1 + \frac{1}{y_2^2} \right] L(y_1) - 2q^2 \left[ 3 + \frac{2q^2}{y_1} + \frac{q^4}{y_1^2} \right] L(y_2) \right] . \quad (153)$$

For the symmetric tensor structure  $(p_1^\mu p_2^\nu + p_2^\mu p_1^\nu)$ , one gets

$$a_{12}^{(1,0)} = \frac{1}{y_1 y_2} \left[ -\frac{4q^2 + (y_1 - y_2)^2}{1-q^2} \right. \\ \left. - 2q^2 \left[ \frac{q^2}{y_1 y_2} + \frac{1+q^2-2y_1 y_2}{(1-q^2)^2} \right] \log(q^2) + \left\{ \frac{q^2}{1-y_1} \right. \right. \\ \left. \left. - \frac{2q^2}{1-y_2} \left[ 1 - y_1 + \frac{q^2}{y_2} - \frac{q^2}{2(1-y_2)} \right] \log\left(\frac{y_1}{q^2}\right) \right. \right. \\ \left. \left. - 2q^2 \left[ 1 + \frac{q^2}{y_2} + \frac{q^2}{y_2^2} \right] L(y_1) + (y_1 \leftrightarrow y_2) \right\} \right] . \quad (154)$$

Finally, the antisymmetric coefficient  $a_{-1}$ , accompanying  $(p_1^\mu p_2^\nu - p_2^\mu p_1^\nu)$ , reads

$$a_{-1}^{(1,0)} = \frac{q^2}{y_1 y_2} \left[ \frac{2 \log(1-y_1)}{y_1} + \frac{1-q^2}{1-y_1} + \frac{q^2}{(1-y_1)^2} \right] \\ - (y_1 \leftrightarrow y_2) . \quad (155)$$

The mass-suppressed coefficients  $a_{ij}^{(1,m)}$  are given by [326]

$$a_{00}^{(1,m)} = \frac{m^2 q^2}{y_1^2} \left[ \log(q^2) \log\left(\frac{y_1^4}{m^4 q^2}\right) + 4\text{Li}_2(1-q^2) \right. \\ \left. + \text{Li}_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right] - \frac{m^2(1-q^2)}{y_1^2} \left[ 1 - \log\left(\frac{y_1}{m^2}\right) \right. \\ \left. + \frac{m^2}{y_1} \left( \text{Li}_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right) \right] + \frac{q^2}{2} n(y_1, \frac{1-3q^2}{q^2}) \\ + (y_1 \leftrightarrow y_2) , \quad (156)$$

and

$$a_{11}^{(1,m)} = \frac{q^2}{1-q^2} \left\{ \frac{4m^2}{y_1^2} \left[ 1 - \log\left(\frac{y_1}{m^2}\right) \right. \right. \\ \left. \left. + \frac{m^2}{y_1} \left( \text{Li}_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right) \right] - n(y_1, 1) \right. \\ \left. + \frac{2m^2 q^2}{y_1(m^2(1-q^2) - y_1)} \left[ \frac{1}{q^2} \log\left(\frac{y_1}{m^2}\right) + \frac{\log(q^2)}{1-q^2} \right. \right. \\ \left. \left. + \left( 1 + \frac{m^2}{m^2(1-q^2) - y_1} \right) N(y_1) \right] \right\} + \\ + \frac{1}{1-q^2} \left\{ \frac{4m^2(1-q^2)}{y_2^2} \left[ \log(q^2) \log\left(\frac{y_2^4}{m^4 q^2}\right) \right. \right. \\ \left. \left. + 4\text{Li}_2(1-q^2) + 2 \left( \text{Li}_2\left(1 - \frac{y_2}{m^2}\right) - \frac{\pi^2}{6} \right) \right] \right. \\ \left. + \frac{4m^2 q^2}{y_2^2} \left[ 1 - \log\left(\frac{y_2}{m^2}\right) + \left( 1 + \frac{m^2}{y_2} \right) \left( \text{Li}_2\left(1 - \frac{y_2}{m^2}\right) \right. \right. \right. \\ \left. \left. - \frac{\pi^2}{6} \right) \right] - \frac{1-2q^4}{q^2} n(y_2, \frac{3-8q^2+6q^4}{1-2q^4}) \right. \\ \left. + \frac{2m^2}{y_2(m^2(1-q^2) - y_2)} \left[ \frac{1}{q^2} \log\left(\frac{y_2}{m^2}\right) + \frac{\log(q^2)}{1-q^2} \right. \right. \\ \left. \left. + \left( 3 + \frac{m^2}{m^2(1-q^2) - y_2} \right) N(y_2) \right] \right\} , \quad (157)$$

and

$$a_{12}^{(1,m)} = \frac{q^2}{1-q^2} \left\{ \frac{4m^2}{y_1^2} \left[ 1 - \log\left(\frac{y_1}{m^2}\right) \right. \right. \\ \left. \left. + \left( \frac{1}{2} + \frac{m^2}{y_1} \right) \left( \text{Li}_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right) \right] \right. \\ \left. - \frac{1-q^2}{q^2} n(y_1, \frac{1}{1-q^2}) + \frac{2m^2}{y_1(m^2(1-q^2) - y_1)} \right. \\ \left. \times \left[ \frac{1}{q^2} \log\left(\frac{y_1}{m^2}\right) + \frac{\log(q^2)}{1-q^2} \right. \right. \\ \left. \left. + \left( 2 + \frac{m^2}{m^2(1-q^2) - y_1} \right) N(y_1) \right] \right\} + (y_1 \leftrightarrow y_2) . \quad (158)$$

The asymmetric coefficient does not get mass corrections:

$$a_{-1}^{(1,m)} = 0 . \quad (159)$$

These results are written in terms of the function

$$L(y_i) = \text{Li}_2\left(-\frac{y_i}{q^2}\right) - \text{Li}_2\left(1 - \frac{1}{q^2}\right) + \log(q^2 + y_i) \log\left(\frac{y_i}{q^2}\right), \quad (160)$$

where  $\text{Li}_2$  is the Spence or dilogarithm function, and the functions  $n(y_i, z)$  and  $N(y_i)$ , which are defined through

$$n(y_i, z) = \frac{m^2}{y_i(m^2 - y_i)} \left[ 1 + z \log\left(\frac{y_i}{m^2}\right) \right] + \frac{m^2}{(m^2 - y_i)^2} \log\left(\frac{y_i}{m^2}\right), \quad (161)$$

and

$$N(y_i) = \log(q^2) \log\left(\frac{y_i}{m^2}\right) + \text{Li}_2(1 - q^2) + \text{Li}_2\left(1 - \frac{y_i}{m^2}\right) - \frac{\pi^2}{6}. \quad (162)$$

The apparent singularity of the function  $n(y_i, z)$  inside the limits of phase space is compensated by the zero in the numerator. In the region  $y_i$  close to  $m^2$  it behaves as

$$n(y_i, z)|_{y_i \rightarrow m^2} = \frac{1}{y_i} \left[ 1 + z \log\left(\frac{y_i}{m^2}\right) \right] - \frac{1}{m^2} \sum_{n=0} \left( \frac{1}{n+2} + \frac{z}{n+1} \right) \left( 1 - \frac{y_i}{m^2} \right)^n. \quad (163)$$

Similarly, the function  $N(y_i)$  guarantees that the coefficients  $a_{ij}^{(1)}$  are finite in the limit  $y_i \rightarrow m^2(1 - q^2)$ :

$$\frac{m^2 N(y_i)}{m^2(1 - q^2) - y_i} \Big|_{y_i \rightarrow m^2(1 - q^2)} = -\frac{\log(1 - q^2)}{q^2} - \frac{\log(q^2)}{1 - q^2}. \quad (164)$$

#### Virtual and soft corrections to IFS

The virtual plus soft photon corrections of the initial-state and final-state vertex (see Eq.(143)) to FSR and ISR, respectively, can be written as [271, 426]

$$d\sigma_{\text{IFS}}^{\text{V+S}} = \frac{\alpha}{\pi} \left[ \delta^{\text{V+S}}(w) d\sigma_{\text{FSR}}^{(0)}(s) + \eta^{\text{V+S}}(s', w) d\sigma_{\text{ISR}}^{(0)}(s') \right], \quad (165)$$

where  $d\sigma_{\text{FSR}}^{(0)}$  and  $d\sigma_{\text{ISR}}^{(0)}$  are the leading order FSR and ISR differential cross sections, respectively,  $w = E_\gamma^{\text{cut}}/\sqrt{s}$ , with  $E_\gamma^{\text{cut}}$  the maximal energy of the soft photon in the  $e^+e^-$  c.m. rest frame, and  $s'$  corresponds to the squared mass of the  $h\bar{h}\gamma$  system. The function  $\delta^{\text{V+S}}(w)$  is independent of the final state. In the limit  $m_e^2 \ll s$ :

$$\delta^{\text{V+S}}(w) = 2 \left[ (L - 1) \log(2w) + \frac{3}{4}L - 1 + \frac{\pi^2}{6} \right], \quad (166)$$

where  $L = \log(s/m_e^2)$ . For two pions in the final state, the function  $\eta^{\text{V+S}}(s', w)$  is given by

$$\begin{aligned} \eta^{\text{V+S}}(s', w) = & -2 \left[ \frac{1 + \beta_\pi^2}{2\beta_\pi} \log(t_\pi) + 1 \right] \\ & \times \left[ \log(2w) + 1 + \frac{s'}{s' - s} \log\left(\frac{s}{s'}\right) \right] + \log\left(\frac{m_\pi^2}{s'}\right) \\ & - \frac{1 + \beta_\pi^2}{\beta_\pi} \left[ 2\text{Li}_2(1 - t_\pi) + \log(t_\pi) \log(1 + t_\pi) - \frac{\pi^2}{2} \right] \\ & - \frac{2 + \beta_\pi^2}{\beta_\pi} \log(t_\pi) - 2, \end{aligned} \quad (167)$$

where

$$\beta_\pi = \sqrt{1 - \frac{4m_\pi^2}{s'}}, \quad t_\pi = \frac{1 - \beta_\pi}{1 + \beta_\pi}. \quad (168)$$

The function  $\eta^{\text{V+S}}(s', w)$  is equivalent to the familiar correction factor derived in [427] for the reaction  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  in the framework of sQED (see also [277]) in the limit  $s \rightarrow s'$ :

$$\log(2w) + 1 + \frac{s'}{s' - s} \log\left(\frac{s}{s'}\right) \Big|_{s \rightarrow s'} = \log(2w') \quad (169)$$

with  $w' = E_\gamma^{\text{cut}}/\sqrt{s'}$ . The factor in the right hand side of Eq.(169) for  $s \neq s'$  arises from defining the soft photon cutoff in the  $e^+e^-$  laboratory frame.

Correspondingly, the function  $\eta^{\text{V+S}}(s', w)$  for two muons in the final state reads

$$\begin{aligned} \eta^{\text{V+S}}(s', w) = & -2 \left[ \frac{1 + \beta_\mu^2}{2\beta_\mu} \log(t_\mu) + 1 \right] \\ & \times \left[ \log(2w) + 1 + \frac{s'}{s' - s} \log\left(\frac{s}{s'}\right) \right] + \log\left(\frac{m_\mu^2}{s'}\right) \\ & - \frac{1 + \beta_\mu^2}{2\beta_\mu} \left[ 4\text{Li}_2(1 - t_\mu) - 2\log(t_\mu) \log\left(\frac{1 + \beta_\mu}{2}\right) - \pi^2 \right] \\ & - \frac{1}{\beta_\mu} \left[ \frac{3}{3 - \beta_\mu^2} + \beta_\mu^2 \right] \log(t_\mu) - 2, \end{aligned} \quad (170)$$

where

$$\beta_\mu = \sqrt{1 - \frac{4m_\mu^2}{s'}}, \quad t_\mu = \frac{1 - \beta_\mu}{1 + \beta_\mu}. \quad (171)$$

#### Real corrections

Matrix elements for the emission of two real photons

$$e^+(p_1) + e^-(p_2) \rightarrow \text{hadrons}(Q) + \gamma(k_1) + \gamma(k_2), \quad (172)$$

are calculated in PHOKHARA following the helicity amplitude method with the conventions introduced in [428, 429]. The Weyl representation for fermions is used where the Dirac matrices

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma_\mu^+ \\ \sigma_\mu^- & 0 \end{pmatrix}, \quad \mu = 0, 1, 2, 3, \quad (173)$$

are given in terms of the unit  $2 \times 2$  matrix  $I$  and the Pauli matrices  $\sigma_i, i = 1, 2, 3$ , with  $\sigma_\pm^\mu = (I, \pm \sigma_i)$ . The contraction of any four-vector  $a^\mu$  with the  $\gamma^\mu$  matrices has the form

$$\not{a} = a_\mu \gamma^\mu = \begin{pmatrix} 0 & a^+ \\ a^- & 0 \end{pmatrix}, \quad (174)$$

where the  $2 \times 2$  matrices  $a^\pm$  are given by

$$a^\pm = a^\mu \sigma_\mu^\pm = \begin{pmatrix} a^0 \mp a^3 & \mp(a^1 - ia^2) \\ \mp(a^1 + ia^2) & a^0 \pm a^3 \end{pmatrix}. \quad (175)$$

The helicity spinors  $u$  and  $v$  for a particle and an antiparticle of four-momentum  $p = (E, \mathbf{p})$  and helicity  $\lambda = \pm 1/2$  are given by

$$\begin{aligned} u(p, \lambda = \pm 1/2) &= \begin{pmatrix} \sqrt{E \mp |\mathbf{p}|} \chi(\mathbf{p}, \pm) \\ \sqrt{E \pm |\mathbf{p}|} \chi(\mathbf{p}, \pm) \end{pmatrix} \equiv \begin{pmatrix} u_I \\ u_{II} \end{pmatrix}, \\ v(p, \lambda = \pm 1/2) &= \begin{pmatrix} \mp \sqrt{E \pm |\mathbf{p}|} \chi(\mathbf{p}, \mp) \\ \pm \sqrt{E \mp |\mathbf{p}|} \chi(\mathbf{p}, \mp) \end{pmatrix} \equiv \begin{pmatrix} v_I \\ v_{II} \end{pmatrix}. \end{aligned} \quad (176)$$

The helicity eigenstates  $\chi(\mathbf{p}, \lambda)$  can be expressed in terms of the polar and azimuthal angles of the momentum vector  $\mathbf{p}$  as

$$\begin{aligned} \chi(\mathbf{p}, +) &= \begin{pmatrix} \cos(\theta/2) \\ e^{i\phi} \sin(\theta/2) \end{pmatrix}, \\ \chi(\mathbf{p}, -) &= \begin{pmatrix} -e^{-i\phi} \sin(\theta/2) \\ \cos(\theta/2) \end{pmatrix}. \end{aligned} \quad (177)$$

Finally, complex polarization vectors in the helicity basis are defined for the real photons:

$$\begin{aligned} \varepsilon^\mu(k_i, \lambda_i = \pm) &= \frac{1}{\sqrt{2}} (0, \mp \cos \theta_i \cos \phi_i + i \sin \phi_i, \\ &\quad \mp \cos \theta_i \sin \phi_i - i \cos \phi_i, \pm \sin \theta_i) \end{aligned} \quad (178)$$

with  $i = 1, 2$ .

### Phase space

One of the key ingredients of any Monte Carlo simulation is an efficient generation of the phase space. The generation of the multiparticle phase space in PHOKHARA is based on the following Lorentz-invariant representation:

$$\begin{aligned} d\Phi_{m+n}(p_1, p_2; k_1, \cdot, k_m, q_1, \cdot, q_n) &= \\ d\Phi_m(p_1, p_2; Q, k_1, \cdot, k_m) d\Phi_n(Q; q_1, \cdot, q_n) \frac{dQ^2}{2\pi}, \end{aligned} \quad (179)$$

where  $p_1$  and  $p_2$  are the four-momenta of the initial particles,  $k_1 \dots k_m$  are the four momenta of the emitted photons and  $q_1 \dots q_n$ , with  $Q = \sum q_i$ , label the four-momenta of the final state hadrons.

When two particles of the same mass are produced in the final state,  $q_i^2 = M^2$ , their phase space is given by

$$d\Phi_2(Q; q_1, q_2) = \frac{\sqrt{1 - \frac{4M^2}{Q^2}}}{32\pi^2} d\Omega, \quad (180)$$

where  $d\Omega$  is the solid angle of one of the final state particles at, for instance, the  $Q^2$  rest frame.

Single photon emission is described by the corresponding leptonic part of phase space

$$d\Phi_2(p_1, p_2; Q, k_1) = \frac{1 - q^2}{32\pi^2} d\Omega_1, \quad (181)$$

with  $q^2 = Q^2/s$  and  $d\Omega_1$  the solid angle of the emitted photon at the  $e^+e^-$  rest frame. The polar angle  $\theta_1$  is defined with respect to the positron momentum  $p_1$ . In order to make the Monte Carlo generation more efficient, the following substitution is performed:

$$\cos \theta_1 = \frac{1}{\beta} \tanh(\beta t_1), \quad t_1 = \frac{1}{2\beta} \log \frac{1 + \beta \cos \theta_1}{1 - \beta \cos \theta_1}, \quad (182)$$

with  $\beta = \sqrt{1 - 4m_e^2/s}$ , which accounts for the collinear emission peaks

$$\frac{d \cos \theta_1}{1 - \beta^2 \cos^2 \theta_1} = dt_1. \quad (183)$$

Then, the azimuthal angle and the new variable  $t_1$  are generated flat.

Considering the emission of two real photons in the c.m. of the initial particles, the four-momenta of the positron, the electron and the two emitted photons are given by

$$\begin{aligned} p_1 &= \frac{\sqrt{s}}{2} (1, 0, 0, \beta), \quad p_2 = \frac{\sqrt{s}}{2} (1, 0, 0, -\beta), \\ k_1 &= w_1 \sqrt{s} (1, \sin \theta_1 \cos \phi_1, \sin \theta_1 \sin \phi_1, \cos \theta_1), \\ k_2 &= w_2 \sqrt{s} (1, \sin \theta_2 \cos \phi_2, \sin \theta_2 \sin \phi_2, \cos \theta_2), \end{aligned} \quad (184)$$

respectively. The polar angles  $\theta_1$  and  $\theta_2$  are again defined with respect to the positron momentum  $p_1$ . Both photons are generated with energies larger than the soft photon cutoff:  $w_i > w$  with  $i = 1, 2$ . At least one of these exceeds the minimal detection energy:  $w_1 > E_\gamma^{\min}/\sqrt{s}$  or  $w_2 > E_\gamma^{\min}/\sqrt{s}$ . In terms of the solid angles  $d\Omega_1$  and  $d\Omega_2$  of the two photons and the normalized energy of one of them, e.g.  $w_1$ , the leptonic part of phase space reads

$$\begin{aligned} d\Phi_3(p_1, p_2; Q, k_1, k_2) &= \frac{1}{2!} \frac{s}{4(2\pi)^5} \\ &\quad \times \frac{w_1 w_2^2}{1 - q^2 - 2w_1} dw_1 d\Omega_1 d\Omega_2, \end{aligned} \quad (185)$$

where the limits of the phase space are determined from the constraint

$$q^2 = 1 - 2(w_1 + w_2) + 2w_1 w_2 (1 - \cos \chi_{12}), \quad (186)$$

with  $\chi_{12}$  being the angle between the two photons

$$\cos \chi_{12} = \sin \theta_1 \sin \theta_2 \cos(\phi_1 - \phi_2) + \cos \theta_1 \cos \theta_2. \quad (187)$$

Again, the matrix element squared contains several peaks, soft and collinear, which should be softened by



choosing suitable substitutions in order to achieve an efficient Monte Carlo generator. The leading behaviour of the matrix element squared is given by  $1/(y_{11} y_{12} y_{21} y_{22})$ , where

$$y_{ij} = \frac{2k_i \cdot p_j}{s} = w_i(1 \mp \beta \cos \theta_i) . \quad (188)$$

In combination with the leptonic part of phase space, we have

$$\frac{d\Phi_3(p_1, p_2; Q, k_1, k_2)}{y_{11} y_{12} y_{21} y_{22}} \sim \frac{dw_1}{w_1(1 - q^2 - 2w_1)} \times \frac{d\Omega_1}{1 - \beta^2 \cos^2 \theta_1} \frac{d\Omega_2}{1 - \beta^2 \cos^2 \theta_2} . \quad (189)$$

The collinear peaks are then flattened with the help of Eq.(182), with one change of variables for each photon polar angle. The remaining soft peak,  $w_1 \rightarrow w$ , is reabsorbed with the following substitution

$$w_1 = \frac{1 - q^2}{2 + e^{-u_1}} , \quad u_1 = \log \frac{w_1}{1 - q^2 - 2w_1} , \quad (190)$$

or

$$\frac{dw_1}{w_1(1 - q^2 - 2w_1)} = \frac{du_1}{1 - q^2} , \quad (191)$$

where the new variable  $u_1$  is generated flat. Multi-channeling is used to absorb simultaneously the soft and collinear peaks, and the peaks of the form factors.


#### NLO cross section and theoretical uncertainty

The LO and NLO predictions for the differential cross section of the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma)$  at DAΦNE energies,  $\sqrt{s} = 1.02$  GeV, are presented in Fig. 47 as a function of the invariant mass of the hadronic system  $M_{\pi\pi}$ . We choose the same kinematical cuts as in the small angle analysis of KLOE [368]; pions are restricted to be in the central region,  $50^\circ < \theta_\pi < 130^\circ$ , with  $|p_T| > 160$  MeV or  $|p_z| > 90$  MeV, the hard photon is not tagged and the sum of the momenta of the two pions, which flows in the opposite direction to the photon's momenta, is close to the beam ( $\theta_{\pi\pi} < 15^\circ$  or  $\theta_{\pi\pi} > 165^\circ$ ). The track mass, which is calculated from the equation

$$\left( \sqrt{s} - \sqrt{|\mathbf{p}_{\pi^+}|^2 + M_{\text{trk}}^2} - \sqrt{|\mathbf{p}_{\pi^-}|^2 + M_{\text{trk}}^2} \right)^2 - (\mathbf{p}_{\pi^+} + \mathbf{p}_{\pi^-})^2 = 0 , \quad (192)$$

lies within the limits  $130 \text{ MeV} < M_{\text{trk}} < 220 \text{ MeV}$  and  $M_{\text{trk}} < (250 - 105\sqrt{1 - (M_{\pi\pi}^2/0.85)^2}) \text{ MeV}$ , with  $M_{\pi\pi}$  in GeV, in order to reject  $\mu^+\mu^-$  and  $\pi^+\pi^-\pi^0$  events. The cut on the track mass, however, does not have any effect for single photon emission, as obviously  $M_{\text{trk}} = m_\pi$  for such events.

The lower plot in Fig. 47 shows the relative size, with respect to the LO prediction, of FSR at LO, ISR corrections at NLO, and IFS contributions. The NLO ISR radiative corrections are almost flat and of the order of



**Fig. 47.** Differential cross section for the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  at LO and NLO for  $\sqrt{s} = 1.02$  GeV. The cuts are the same as in the small angle analysis of KLOE, including the cut on the track mass. The lower plot shows the relative size of FSR at LO, ISR at NLO and IFS contributions with respect to the full LO prediction.

−8%, FSR is clearly below 1%, while IFS corrections are also small although they become of the order of a few per cent at high values of  $M_{\pi\pi}$ .

To estimate the systematic uncertainty of the NLO prediction, we observe that leading logarithmic two-loop  $\mathcal{O}(\alpha^2)$  corrections and the associate real emission are not included. For samples with untagged photons the process  $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$  might also become a sizable background. This process, however, can be simulated with the Monte Carlo event generator EKHARA [216, 215]. Its contribution depends on the pion pair invariant mass, ranges from 0.1 – 0.8% for the KLOE event selection, and has been taken into account in the KLOE analysis [368].

From naïve exponentiation one expects that LL corrections at next-to-next-to-leading order (NNLO) are of the order of  $\frac{1}{2}(\frac{3}{2}(\alpha/\pi) \log(s/m_e^2))^2 \approx 0.1\text{--}0.2\%$  for inclusive observables. For less inclusive distributions, a larger error is expected. The conservative estimate of the accuracy of PHOKHARA from ISR is 0.5%. This has been confirmed

by comparisons with KKMC [257, 256], where the biggest observed difference is about 0.3% in the invariant mass regions, which are not close to the nominal energies of the experiments. Improving the accuracy of PHOKHARA below 0.5%, however, will be required to meet the growing experimental requirements in the near future.

#### 4.2.4 FSR beyond sQED\*VMD model

The model of the FSR emission from pions described in details in Sections 4.2.1 and 4.2.3 will be called for short the sQED\*VMD model. A question arises how well it can reflect the data. As it is shown in [311], the first two terms in the expansion of the FSR amplitude as a function of  $k^0/\sqrt{Q^2}$  (i.e. the divergence and the constant) are fully given by the pion form factor. Thus only for a hard photon emission one could expect that going beyond this approximation is necessary. Moreover, the pion form factor is extremely big in the  $\rho$  resonance region and thus the validity of this approximation is further extended. In the kinematical regions, where there are resonance contributions not contained in the pion form factor and also near the  $\pi^+\pi^-$  threshold, where the emitted photon is hard and the pion form factor is relatively small, going beyond the sQED\*VMD model is necessary and one needs a more general description of the amplitude  $M(\gamma^*(Q) \rightarrow \gamma(k) + \pi^+(q_1) + \pi^-(q_2))$ .

In the general case the amplitude of the reaction  $\gamma^*(Q) \rightarrow \gamma(k) + \pi^+(q_1) + \pi^-(q_2)$  depends on three 4-momenta, which can be chosen as  $Q, k$ , and  $l \equiv q_1 - q_2$ . The second-rank Lorentz tensor  $M^{\mu\nu}(Q, k, l)$ , that describes the FSR amplitude, can be decomposed through ten independent tensors [430, 431]. Taking into account the charge conjugation symmetry of the S-matrix element  $(\langle \gamma(k), \pi^+(q_1) \pi^-(q_2) | S | \gamma^*(Q) \rangle =$

$$\langle \gamma(k), \pi^-(q_1) \pi^+(q_2) | S | \gamma^*(Q) \rangle),$$

the photon crossing symmetry ( $Q \leftrightarrow -k$  and  $\mu \leftrightarrow \nu$ ) and the gauge invariance conditions  $Q_\mu M^{\mu\nu}(Q, k, l) = 0$  and  $M_F^{\mu\nu}(Q, k, l) k_\nu = 0$ , the number of the independent tensors decreases to five. For a final real photon, i.e.  $k^2 = 0$  and  $k^\nu \epsilon_\nu = 0$  ( $\epsilon_\nu$  is the polarization vector of the final photon) and the initial virtual photon produced in  $e^+e^-$  annihilation ( $Q^2 \geq 4m_\pi^2$ ), the FSR tensor can be rewritten in the terms of three gauge invariant tensors [430, 431]

$$M^{\mu\nu}(Q, k, l) = \tau_1^{\mu\nu} f_1 + \tau_2^{\mu\nu} f_2 + \tau_3^{\mu\nu} f_3, \quad (193)$$

where the gauge invariant tensors  $\tau_i^{\mu\nu}$  read

$$\begin{aligned} \tau_1^{\mu\nu} &= k^\mu Q^\nu - g^{\mu\nu} k \cdot Q, \\ \tau_2^{\mu\nu} &= k \cdot l (l^\mu Q^\nu - g^{\mu\nu} k \cdot l) + l^\nu (k^\mu k \cdot l - l^\mu k \cdot Q), \\ \tau_3^{\mu\nu} &= Q^2 (g^{\mu\nu} k \cdot l - k^\mu l^\nu) + Q^\mu (l^\nu k \cdot Q - Q^\nu k \cdot l) \end{aligned} \quad (194)$$

It thus follows that the evaluation of the FSR tensor amounts to the calculation of the scalar functions

$$f_i(Q^2, Q \cdot k, k \cdot l) \quad (i = 1, 2, 3).$$

As it is clear from the above discussion, the extraction of the pion form factor from radiative return experiments is a demanding task. The main problem is that

in the same experiments one has to test the models describing the pion-photon interactions (see Section 4.3) and to extract the pion form factor needed for the evaluation of the muon anomalous magnetic moment. Fortunately, there are event selections, which naturally suppress the FSR contributions, independently of their nature. These were already discussed in Section 4.2.1 in the context of the sQED\*VMD model.

Extensive theoretical studies of the role of the FSR emission beyond the sQED\*VMD model were performed [331, 341, 343, 345, 344]. They are important mainly for the KLOE measurements at DAΦNE as at  $B$ -meson factories the FSR is naturally suppressed and the accuracy needed in its modeling is by far less demanding than for the KLOE purposes.

For DAΦNE, running on or near the  $\phi$  resonance, the following mechanisms of the  $\pi^+\pi^-$  final state photon emission have to be considered:

bremsstrahlung process

$$e^+ + e^- \rightarrow \pi^+ + \pi^- + \gamma, \quad (195)$$

which is modelled by sQED\*VMD;

$\phi$  direct decay

$$e^+ + e^- \rightarrow \phi \rightarrow (f_0; f_0 + \sigma) \gamma \rightarrow \pi^+ + \pi^- + \gamma, \quad (196)$$

and double resonance process

$$e^+ + e^- \rightarrow (\phi; \omega') \rightarrow \rho \pi \rightarrow \pi^+ + \pi^- + \gamma. \quad (197)$$

The resonance chiral theory (R $\chi$ T) [432, 433] was used in [343, 344] to estimate the contributions beyond sQED\*VMD. They were implemented at the leading order into an event generator FASTERD [345]. Having in mind that at present the models still await accurate experimental tests, in the event generator FASTERD other models [434, 435] were also implemented. To include both next-to-leading-order radiative corrections and the discussed mechanisms of the FSR production, a part of the FASTERD code, based on models [434, 435], was implemented by O. Shekhovtsova in PHOKHARA6.0 (PHOKHARA6.1 [436]) and the studies presented below are based on this code. The model used there, even if far from an ideal, is the best tested model available in literature.

We shortly describe main features of the models used to describe processes contributing to FSR photon emission listed above. For more detailed description and the calculation of the function  $f_i$  we refer the reader to [331, 341, 345] (see also references therein).

The sQED\*VMD part gives contributions to  $f_1$  and  $f_2$ .

The  $\phi$  direct decay is assumed to proceed through the intermediate scalar meson state:  $\phi \rightarrow (f_0 + \sigma) \gamma \rightarrow \pi \pi \gamma$ . Various models are proposed to describe the  $\phi$ -scalar- $\gamma$  vertex: either it is the direct decay  $\phi \rightarrow (scalar) \gamma$  or the vertex is generated dynamically through the loop of the charged kaons. As shown in [341], in the framework of any model the  $\phi$  direct decay affects only the form factor  $f_1$  of Eq. (193).

The double resonance contribution consists of the off-shell  $\phi$  meson decay into  $(\rho^\pm \pi^\mp)$  and subsequent decay  $\rho \rightarrow \pi\gamma$ . In the energy region around 1 GeV the tail of the excited  $\omega$  meson can also play a role and  $\gamma^* \rightarrow \omega' \rightarrow \rho\pi$  has to be considered. The double resonance mechanism affects all three form factors  $f_i$  of Eq. (193).


Assuming isospin symmetry, this part can be deduced from the measurement of the neutral pion pair production. Various models [434, 435] were confronted with data by KLOE [437] for the neutral mode. The model, which was reproducing the data in the best way was adopted to be used for the charged pion pair production relying on the isospin symmetry [436].

In [331] it was shown that an important tool for testing of the various models of the FSR emission is the charge asymmetry. At the leading order it originates from the fact that the pion pair couples to even (odd) number of photons if the final state photon is emitted from the final (initial) state. The interference diagrams do not give contribution to the integrated cross section for C-even event selections but produce asymmetry in the angular distribution. The definitions and experimental studies based on the charge asymmetry are presented in Section 4.3.2


Few strategies can be adopted how to profit in the best way from the KLOE data taken on and off peak. The 'easiest' part is to look for the event selections, where the FSR contributions are negligible. This was performed by KLOE [368] (see Section 4.4.1), giving important information on the pion form factor relevant for the prediction of the hadronic contributions to the muon anomalous magnetic moment  $a_\mu$ . Typical contributions of the FSR (1%-4%) to the differential cross section (Fig. 47 and 48) allow for excellent control on the accuracy of these corrections. One disadvantage of using this event selection is that it does not allow to perform measurements near the pion production threshold.

The next step, partly discussed in Section 4.3.2, is to confront the models based on the isospin symmetry and the neutral channel data with charged pion data taken off-peak, where the contributions from models beyond the sQED\*VMD approximation are relatively small (Fig. 49). For the off-peak data [438] the region below  $Q^2 = 0.3\text{GeV}^2$  can be covered experimentally, however, the small statistics in this region makes it difficult to perform high-precision tests of the models. For this analysis an accurate knowledge of the pion form factor at the nominal energy of the experiment is important as it defines the sQED\*VMD predictions and the FSR corrections (Fig. 50) are sizable.

The last step, which allows for the most accurate FSR model testing and profits from the knowledge of the pion form factor from previous analysis, is the on-peak large angle measurement. The large FSR corrections coming from sources beyond the sQED\*VMD approximation (Fig. 49 and 50) make these data [439] the most valuable source of information on these models. In this case the accumulated data set is much larger than the off-peak data and one is able to cover also the region below  $Q^2 = 0.3\text{GeV}^2$ .



**Fig. 48.** Relative contribution of the FSR to the differential cross section of the reaction  $e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma)$  for  $\sqrt{s} = m_\phi$  and low invariant masses of pion pairs. KLOE small angle event selection [368] was used and for this event selection the relative contribution of the FSR is almost identical also for off peak cross section. The effect of mass track cut (see Section 4.4.1) is shown. ISRNLO refers to initial state correction at next to leading order (NLO). IFSNLO cross section contains the final states emission at NLO.




**Fig. 49.** The contributions of FSR beyond sQED\*VMD approximation (see eq. 196 and 197) for large angle KLOE event selection [438, 439] for  $\sqrt{s} = m_\phi$  and for  $\sqrt{s} = 1\text{GeV}$ .

### 4.3 Experiment confronting theory

#### 4.3.1 Study of the process $e^+e^- \rightarrow \pi^+\pi^-\gamma$ with FSR with CMD-2 detector at VEPP-2M

A study of the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  with photon radiation from the final state pions can be used to answer the question whether one can treat pions as point-like particles and apply scalar QED in the calculation of the radiative corrections to the cross section. In particular, one can



**Fig. 50.** Relative contribution of FSR to the differential cross section of the reaction  $e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma)$  for  $\sqrt{s} = m_\phi$  and for  $\sqrt{s} = 1\text{GeV}$ . KLOE large angle event selection [438, 439] was used.

compare the photon spectra obtained using scalar QED with the ones found in data.

The radiative corrections due to photon emission in the final state contribute about 1% to the cross section. The hadronic contribution to  $(g-2)/2$  of the muon from the process  $e^+e^- \rightarrow \pi^+\pi^-$  amounts to 50 ppm, while the anomalous magnetic moment of the muon was measured in the E821 experiment at BNL with an accuracy of 0.5 ppm [23]. Therefore the theoretical uncertainty on the cross section calculation of the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  should be better than 1% to be able to neglect the error of this contribution with respect to the 0.5 ppm. These facts are the main motivation to study this process.

In the following, the preliminary results of the analysis of the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  with final state radiation are presented. The analysis is based on an integrated luminosity of about  $1.2\text{ pb}^{-1}$ . The data were collected in the c.m. energy range from 720 MeV to 780 MeV by the CMD-2 detector. Pions were considered as point-like objects and scalar QED was applied to describe the photon radiation in the final state.

#### Event selection

For the analysis, the data were taken in a c.m. energy range from 720 MeV to 780 MeV, with the photon detected in the CsI calorimeter of the CMD-2 experiment. Events from the processes  $e^+e^- \rightarrow e^+e^-\gamma$ ,  $e^+e^- \rightarrow \mu^+\mu^-\gamma$  have a very similar signature in the detector compared to  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  events. In addition, the cross section of the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  with a final state photon is more than ten times smaller than the one for the similar process with a photon radiated in the initial state. On the other hand, the cross section of the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  has a resonant shape in the studied energy region due to the presence of the  $\rho$ -meson. This allows to enhance significantly the fraction of events  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  with final state photons. Selecting events below the  $\rho$ -resonance, the photon radiation from the initial state

will decrease the cross section, whereas the process with a final state photon has practically no energy dependence. Several curves describing the ratio  $\sigma_{\pi^+\pi^-\gamma}^{FSR+ISR}/\sigma_{\pi^+\pi^-\gamma}^{ISR}$  plotted against the c.m. energy are presented in Fig. 51 for different energy thresholds of photon detection in the calorimeter. It is clearly visible that the optimal energy range to be used in this study goes from 720 MeV up to 780 MeV.

It can also be seen that the cross section ratio increases with the energy threshold for photons. The fraction of  $\pi^+\pi^-\gamma$  events with a final state radiation photon increases with photon energy. This allows to enrich the spectrum of  $\pi^+\pi^-\gamma$  events with FSR. Of special interest is the part of the photon spectra in which the photon energy is of the same order as the pion mass or larger.


A typical  $\pi^+\pi^-\gamma$  event in the CMD-2 detector has two tracks in the drift chamber with two associated clusters in the CsI calorimeter and a third cluster representing the radiated photon. To suppress multiphoton events and significantly cut off the collinear  $\pi^+\pi^-$  events the following requirements were applied: The angle between the photon direction and the missing momentum must be larger than 1 rad and the angle between any of the tracks and the photon direction must be larger than 0.2 rad.

To suppress  $e^+e^-\gamma$  events, a parameter  $W = p/E$  was used, in which the particle momentum  $p$  (measured in the drift chamber) is divided by the energy  $E$  (measured in the CsI calorimeter). Simulation results are presented in Fig. 52. The condition  $W < 0.4$  reduces the electron contribution to the level of  $\sim 1\%$ . The squared invariant mass for electrons, muons and pions is plotted in Fig. 53. The condition  $M^2 > 10000\text{ MeV}^2$  rejects additional electrons and muons by a factor of 1.5. About 1% of pion events are lost.


#### Preliminary results of the analysis

The histogram of the photon spectrum from the CMD-2 experimental data is presented in Fig. 54. The histogram represents the simulation, while the points with error bars represent the experimental result. Vertical dotted lines divide the plot area into three zones. The inscription inside each zone indicates the fraction of  $\pi^+\pi^-\gamma$  events with final state radiation with respect to the total number of events. The number of the simulated events was normalized to the experimental one. The average deviation between the two distributions was found to be  $(-2.1 \pm 2.3)\%$ . As a result, one can conclude that there are no indications that photon radiation by pions needs to be described beyond the scalar QED framework. Pions can be treated as point-like objects and the application of the scalar QED is found to be valid within the stated accuracy. Unfortunately, lack of statistics in the energy range under study does not allow to check this assumption with better accuracy. Forthcoming results from VEPP-2000 will significantly improve the statistical error.





(a)



(b)

**Fig. 51.** (a) Ratio  $\sigma_{ISR+FSR}/\sigma_{ISR}$  vs c.m.energy. The set of curves indicates how this ratio depends on the detection threshold energy for photons. The threshold energy in MeV is stated over the curves.


**Fig. 52.** (b) Distributions of the parameter  $W$  for events of the processes  $e^+e^- \rightarrow \pi^+\pi^-\gamma$ ,  $e^+e^- \rightarrow \mu^+\mu^-\gamma$  and  $e^+e^- \rightarrow e^+e^-\gamma$ , for a c.m.-energy of 780 MeV.

#### 4.3.2 Study of the process $e^+e^- \rightarrow \pi^+\pi^-\gamma$ with FSR with KLOE detector


As has been explained in Sec. 4.2, the forward-backward asymmetry

$$A_{FB}(Q^2) = \frac{N(\theta_{\pi^+} > 90^\circ) - N(\theta_{\pi^+} < 90^\circ)}{N(\theta_{\pi^+} > 90^\circ) + N(\theta_{\pi^+} < 90^\circ)}(Q^2) \quad (198)$$

can be used to test the validity of the description of the various mechanisms of the  $\pi^+\pi^-$  final state photon emis-



(a)



(b)

**Fig. 53.** (a) Distributions of the parameter  $M^2$  for events of the processes  $e^+e^- \rightarrow \pi^+\pi^-\gamma$ ,  $e^+e^- \rightarrow \mu^+\mu^-\gamma$  and  $e^+e^- \rightarrow e^+e^-\gamma$  for a c.m.-energy of 780 MeV.

**Fig. 54.** (b) Distribution of the events  $\pi^+\pi^-\gamma$  vs photon energy in relative units. The fraction of the  $\pi^+\pi^-\gamma$  events with final state radiation is stated for each vertical zone.

sion by confronting the output of the Monte Carlo generator with data. In the following studies, the Monte Carlo generator PHOKHARA6.1 [436] was used. The parameters for the pion form factor were taken from [440] based on the parametrization of Kühn and Santamaria [441]. The parameters for the description of the  $\phi$  direct decay and the double resonance contribution were taken from the KLOE analysis of the neutral mode [437].

To suppress higher order effects, for which the interference and thus the asymmetry is not implemented in the Monte Carlo generator, a rather tight cut on the track mass variable (see Sec. 4.4.1 and Fig. 62) of  $|M_{\text{trk}} - M_{\pi^\pm}| < 10$  MeV has been applied in the data in addition to the *large angle* selection cuts described in Sec. 4.4.1. This should reduce events with more than one hard photon emitted and enhance the contribution of the final state radiation processes under study over the dominant ISR process.

The data sets used in the analysis were taken in two different periods:

- The data taken in 2002 was taken with DAΦNE operating at the  $\phi$ -peak, at  $\sqrt{s} = M_\phi$  (240 pb $^{-1}$ )
- The data taken in 2006 was collected with DAΦNE operating 20 MeV below the  $\phi$ -peak, at  $\sqrt{s} = 1000$  MeV (230 pb $^{-1}$ )

Since the 2006 data were taken more than  $4\Gamma_\phi$  below the resonant peak ( $\Gamma_\phi = 4.26$  MeV), one expects the contributions from the  $\phi$  direct decay and the double resonance contribution to be suppressed compared to the data taken on the peak in 2002 (see Fig. 49). In fact, one observes a very different shape of the forward-backward asymmetry for the two different data sets, as can be seen in Fig. 55 and Fig. 56. Especially in the region below 0.4 GeV $^2$  and in the vicinity of the  $f_0(980)$  at 0.96 GeV $^2$  one observes different trends in the asymmetries for the two data sets.

One can also see that qualitatively, the theoretical description used to model the different FSR contributions agrees well with the data, although especially at low  $M_{\pi\pi}^2$  the data statistics becomes poor and the data asymmetry points have large errors. In particular, the *off-peak* data in Fig. 56 shows very good agreement above 0.35 GeV $^2$ . In this case, the asymmetry is dominated fully by the bremsstrahlungs-process, as the other processes do not contribute outside the  $\phi$ -resonance. The assumption of pointlike pions (sQED) used to describe the bremsstrahlung in the Monte Carlo generator seems to be valid above 0.35 GeV $^2$ , while below it is difficult to make a statement due to the large statistical errors of the data points.


However, to obtain a solid quantitative statement on the validity of the models, as it is needed e.g. in the radiative return analyses at the KLOE experiment, one needs to understand how a discrepancy between theory and data in the forward-backward asymmetry affects the cross section, as it is the cross section one wants to measure. This requires further work which at the moment is still in progress.

It should also be mentioned that the KLOE experiment has taken almost 10 times more data in the years 2004-2005 than is shown in Fig. 55, with DAΦNE operating at the  $\phi$ -peak energy. This is unfortunately not the case for the *off-peak* data, which is restricted to the dataset shown in Fig. 56. In future, the larger dataset from 2004-2005 may be used to determine with high precision the parameters of the  $\phi$  direct decay and the double resonance contribution, together with the results from the neutral channel and the assumption of isospin symmetry.

#### 4.4 The use of the radiative return as an experimental tool


##### 4.4.1 Radiative return at KLOE

The KLOE experiment, in operation at the DAΦNE  $e^+e^-$  collider in Frascati between 1999 and 2006, utilizes the radiative return to obtain precise measurements of hadronic



**Fig. 55.** (a) Preliminary Forward-Backward asymmetry for data taken at  $\sqrt{s} = M_\phi$  in 2002, and the corresponding Monte Carlo prediction using the PHOKHARA6.1 generator. (b) Absolute difference between the asymmetries from data and Monte Carlo prediction. Used with permission of the KLOE collaboration.

cross sections in the energy range below 1 GeV. As the DAΦNE machine was designed to operate as a meson factory with collision energy equal to the mass of the  $\phi$ -meson ( $m_\phi = 1.01946$  GeV), with limited possibility to change the energy of the colliding beams while maintaining stable running conditions, the use of events with initial state radiation of hard photons from the  $e^+$  or the  $e^-$  is the only way to access energies below DAΦNE's nominal collision energy. These low-energy cross sections are important in the theoretical evaluation of the muon magnetic moment anomaly  $a_\mu = (g_\mu - 2)/2$  [7], and high precision is needed since the uncertainty on the cross section data enters the uncertainty of the theoretical prediction. The channel  $e^+e^- \rightarrow \pi^+\pi^-$  gives the largest contribution to the hadronic part  $a_\mu^{\text{had}}$  of the anomaly, therefore so far KLOE efforts have concentrated on the derivation of the pion pair-production cross section  $\sigma_{\pi\pi}$  from measurements of the differential cross section  $\frac{d\sigma_{\pi\pi\gamma(\gamma)}}{dM_{\pi\pi}^2}$ , in which  $M_{\pi\pi}^2$  is



**Fig. 56.** (a) Preliminary Forward-Backward asymmetry for data taken at  $\sqrt{s} \simeq 1000$  MeV in 2006, and the corresponding Monte Carlo prediction using the PHOKHARA6.1 generator. (b) Absolute difference between the asymmetries from data and Monte Carlo prediction. Used with permission of the KLOE collaboration.


the invariant mass squared of the di-pion system in the final state.

The KLOE detector (shown in Fig. 57), which consists of a high resolution drift chamber ( $\sigma_p/p \leq 0.4\%$ ) and an electromagnetic calorimeter with excellent time ( $\sigma_t \sim 54 \text{ ps}/\sqrt{E [\text{GeV}]} \oplus 100 \text{ ps}$ ) and good energy ( $\sigma_E/E \sim 5.7\%/\sqrt{E [\text{GeV}]}$ ) resolution, is optimally suited for this kind of analyses.

#### The KLOE $\pi\pi\gamma$ analyses

The KLOE analyses to obtain  $\sigma_{\pi\pi}$  use two different sets of acceptance cuts:

- In the *small angle* analysis, photons are emitted within a cone of  $\theta_\gamma < 15^\circ$  around the beamline (narrow cones in Fig. 57), and the two charged pion tracks have  $50^\circ < \theta_\pi < 130^\circ$ . The photon is not explicitly detected, its direction is reconstructed from the track momenta by closing kinematics:  $\mathbf{p}_\gamma \simeq \mathbf{p}_{\text{miss}} = -(\mathbf{p}_{\pi^+} + \mathbf{p}_{\pi^-})$ . The



**Fig. 57.** KLOE detector with the selection regions for small angle photons (narrow cones) and for pion tracks and large angle photons (wide cones). Used with permission of the KLOE collaboration.

separation of pion- and photon selection regions in this analysis greatly reduces the contamination from the resonant process  $e^+e^- \rightarrow \phi \rightarrow \pi^+\pi^-\pi^0$  in which the  $\pi^0$  mimicks the missing momentum of the photon(s) and from the final state radiation process  $e^+e^- \rightarrow \pi^+\pi^-\gamma_{\text{FSR}}$ . Since ISR-photons are mostly collinear with the beam line, a high statistics for the ISR signal events remains. On the other hand, a high energy photon emitted at angles close to the incoming beams forces the pions also to have a small angle with respect to the beamline (and thus outside the selection cuts), resulting in a kinematical suppression of events with  $M_{\pi\pi}^2 < 0.35 \text{ GeV}^2$ .

- The *large angle* analysis requires both photons and pions to be emitted at  $50^\circ < \theta_{\pi,\gamma} < 130^\circ$  (wide cones in Fig. 57), allowing for a detection of the photons in the barrel of the calorimeter. This analysis allows to reach the  $2\pi$  threshold region, at the price of higher background contributions from the  $\pi^+\pi^-\pi^0$  final state and events with final state radiation. In addition, events from the decays  $\phi \rightarrow f_0\gamma \rightarrow \pi^+\pi^-\gamma$  and  $\phi \rightarrow \pi^\pm\rho^\mp \rightarrow \pi^\pm\pi^\mp\gamma$ , which need to be described by model-dependent parameterisations, contribute to the spectrum of selected events (running at  $\phi$  peak).

Two analyses based on the *small angle* acceptance cuts have been carried out. The first one using  $140 \text{ pb}^{-1}$  of data taken in the year 2001 was published in 2005 [367], the second one, based on  $240 \text{ pb}^{-1}$  of data taken in 2002 was published in 2008 [442].

The differential cross section is obtained from the spectrum of selected events  $N^{\text{sel}}$  subtracting the residual background (mostly  $\mu\mu\gamma(\gamma)$ ,  $\pi\pi\pi$  and radiative Bhabha events)

and dividing by the selection efficiencies and the integrated luminosity:

$$\frac{d\sigma_{\pi\pi\gamma(\gamma)}}{dM_{\pi\pi}^2} = \frac{N^{\text{sel}} - N^{\text{bkg}}}{\Delta M_{\pi\pi}^2} \cdot \frac{1}{\varepsilon_{\text{sel}}} \cdot \frac{1}{\int L dt} \quad (199)$$

$\Delta M_{\pi\pi}^2$  is the bin width used in the analysis (typically 0.01  $\text{GeV}^2$ ), and  $\int L dt$  is the integrated luminosity obtained from Bhabha events detected at large angles ( $55^\circ < \theta_e < 125^\circ$ ) and the reference cross section from the BabaYaga generator [225, 227] (discussed in Sec. 2). The total cross section is then obtained from the formula

$$\sigma_{\pi\pi}(M_{\pi\pi}^2) = s \cdot \frac{d\sigma_{\pi\pi\gamma(\gamma)}}{dM_{\pi\pi}^2} \frac{1}{H(s, M_{\pi\pi}^2)} \quad (200)$$

In this formula,  $s$  is the squared energy at which the DAΦNE collider is operated during the data taking, and  $H(s, M_{\pi\pi}^2)$  is the radiator function describing the emission of photons from the  $e^+$  or the  $e^-$  in the initial state. Note that Eq. 200 does not contain the effects from final state radiation from pions. These effects complicate the analysis, since the KLOE detector can not distinguish whether photons in an event were emitted in the initial or the final state. The PHOKHARA Monte Carlo generator [329], which includes final state radiation at next-to-leading order in the pointlike-pion approximation, is used to properly take into account final state radiation in the analyses. This is important because the *bare* cross section used to evaluate  $a_\mu^{\text{had}}$  via an appropriate dispersion integral should be inclusive with respect to final state radiation, and also needs to be undressed from vacuum polarisation effects present in the virtual photon produced in the  $e^+e^-$  annihilation. For the latter, we use a function provided by F. Jegerlehner [443] (see Section 6), and correct the cross section via

$$\sigma_{\pi\pi}^{\text{bare}}(M_{\pi\pi}^2) = \sigma_{\pi\pi}^{\text{dressed}}(M_{\pi\pi}^2) \left( \frac{\alpha(0)}{\alpha(M_{\pi\pi}^2)} \right)^2 \quad (201)$$


Here  $\alpha(0)$  is the fine structure constant in the limit  $q = 0$ , and  $\alpha(M_{\pi\pi}^2)$  represents its effective value at the specific value of the squared invariant mass of the di-pion system. Since the hadronic contribution to  $\alpha(M_{\pi\pi}^2)$  comes from a dispersion integral which includes the hadronic cross section itself as the integrand (see Sec. 6), the correct procedure has to be iterative and it should include the same data that must be corrected. However, since the correction is at the few percent level, the inclusion of the new KLOE data will not change  $\alpha(M_{\pi\pi}^2)$  at a level which would significantly affect the analyses. We therefore have used the values for  $\alpha(M_{\pi\pi}^2)$  derived from the existing hadronic cross section database. As an example, Fig. 58 shows the KLOE result for  $d\sigma_{\pi\pi\gamma(\gamma)}/dM_{\pi\pi}^2$  obtained from data taken in the year 2002 [442]. Inserting this differential cross section into Eq. 200 and the outcome into Eq. 201, one derives  $\sigma_{\pi\pi}^{\text{bare}}$ . Using the *bare* cross section to get the  $\pi\pi$ -contribution to  $a_\mu^{\text{had}}$  between 0.35 and 0.95  $\text{GeV}^2$  then gives the value (in units of  $10^{-10}$ )

$$a_\mu^{\pi\pi}(0.35 - 0.95 \text{ GeV}^2) = (387.2 \pm 0.5_{\text{stat}} \pm 2.4_{\text{exp}} \pm 2.3_{\text{th}})$$

Reconstruction Filter	negligible
Background subtraction	0.3 %
Trackmass	0.2 %
Particle ID	negligible
Tracking	0.3 %
Trigger	0.1 %
Unfolding	negligible
Acceptance ( $\theta_{\pi\pi}$ )	0.2 %
Acceptance ( $\theta_\pi$ )	negligible
Software Trigger (L3)	0.1 %
Luminosity ( $0.1_{\text{th}} \oplus 0.3_{\text{exp}}$ ) %	0.3 %
$\sqrt{s}$ dep. of $H$	0.2 %
Total exp systematics	0.6 %
Vacuum Polarization	0.1 %
FSR resummation	0.3 %
Rad. function $H$	0.5 %
Total theory systematics	0.6 %

**Table 13.** List of systematic errors on the  $\pi\pi$ -contribution to  $a_\mu^{\text{had}}$  between 0.35 and 0.95  $\text{GeV}^2$  when using the  $\sigma_{\pi\pi}$  cross section measured by the KLOE experiment in the corresponding dispersion integral [442].

Table 13 shows the contributions to the systematic errors on  $a_\mu^{\pi\pi}(0.35 - 0.95 \text{ GeV}^2)$ .




**Fig. 58.** Differential radiative cross section  $d\sigma_{\pi\pi\gamma(\gamma)}/dM_{\pi\pi}^2$ , inclusive in  $\theta_\pi$  and with  $0^\circ < \theta_\gamma < 15^\circ$  or  $165^\circ < \theta_\gamma < 180^\circ$  measured by the KLOE experiment [442]. Used with permission of the KLOE collaboration.

#### Radiative corrections and Monte Carlo tools

The radiator function is a crucial ingredient in this kind of radiative return analyses, it is obtained using the





**Fig. 59.** The dimensionless radiator function  $H(s, M_{\pi\pi}^2)$ , inclusive in  $\theta_{\pi,\gamma}$ . The value used for  $s$  in the Monte Carlo production was  $s = (M_\phi)^2 = (1.019456 \text{ GeV})^2$ .

relation


$$H(s, M_{\pi\pi}^2) = s \cdot \frac{3M_{\pi\pi}^2}{\pi\alpha^2\beta_\pi^3} \cdot \frac{d\sigma_{\pi\pi\gamma(\gamma)}^{ISR}}{dM_{\pi\pi}^2} \Big|_{|F_{2\pi}|^2=1}, \quad (202)$$

in which  $\frac{d\sigma_{\pi\pi\gamma(\gamma)}^{ISR}}{dM_{\pi\pi}^2} \Big|_{|F_{2\pi}|^2=1}$  is evaluated using the PHOKHARA Monte Carlo generator in next-to-leading order ISR-only configuration, with the squared pion form factor  $|F_{2\pi}|^2$  set to 1.  $\beta_\pi = \sqrt{1 - \frac{4m_\pi^2}{M_{\pi\pi}^2}}$  is the pion velocity. While Eq. 202 provides a convenient mechanism to extract the dimensionless quantity  $H(s, M_{\pi\pi}^2)$  also for specific angular regions of pions and photons by applying the relevant cuts to  $\frac{d\sigma_{\pi\pi\gamma(\gamma)}^{ISR}}{dM_{\pi\pi}^2} \Big|_{|F_{2\pi}|^2=1}$ , in the published KLOE analyses,

$H(s, M_{\pi\pi}^2)$  is evaluated fully inclusive for pion and photon angles:  $0^\circ < \theta_{\pi,\gamma} < 180^\circ$ . Fig. 59 shows the radiator function in the range of  $0.35 < M_{\pi\pi}^2 < 0.95 \text{ GeV}^2$ . As can be seen from Table 13, the 0.5% uncertainty of the radiator function quoted by the authors of PHOKHARA translates into an uncertainty of 0.5% in the  $\pi\pi$ -contribution to  $a_\mu^{\text{had}}$  between 0.35 and 0.95  $\text{GeV}^2$ , giving the largest individual contribution and dominating the theoretical systematic error.

The presence of events with final state radiation in the data sample affects the analyses in several ways:

- Passing from  $M_{\pi\pi}^2$  to  $(M_{\pi\pi}^0)^2$  The presence of final state radiation shifts the observed value of  $M_{\pi\pi}^2$  (evaluated from the momenta of the two charged pion tracks in the events) away from the value of the invariant mass squared of the virtual photon produced in the collision of the electron and the positron,  $(M_{\pi\pi}^0)^2$ . The transition from  $M_{\pi\pi}^2$  to  $(M_{\pi\pi}^0)^2$  is performed using a modified version of the PHOKHARA Monte Carlo generator, which allows to (approximately) determine whether a generated photon comes from the initial or the final state [444]. Fig. 60 shows the probability matrix relating  $M_{\pi\pi}^2$  to  $(M_{\pi\pi}^0)^2$  by giving the probability for an event in a bin of  $M_{\pi\pi}^2$  to end up in a bin of  $(M_{\pi\pi}^0)^2$ . It can be seen that the shift is only in one direction,  $(M_{\pi\pi}^0)^2 \geq M_{\pi\pi}^2$ , so events with one photon from initial state radiation and one photon from final state radiation move to a higher value of  $(M_{\pi\pi}^0)^2$ . The



**Fig. 60.** Probability matrix relating the measured quantity  $M_{\pi\pi}^2$  to  $(M_{\pi\pi}^0)^2$ . To produce this plot, a private version of the PHOKHARA Monte Carlo generator was used [444]. The photon angle is restricted to  $\theta_\gamma < 15^\circ$  ( $\theta_\gamma > 165^\circ$ ).

entries lining up above  $(M_{\pi\pi}^0)^2 \simeq 1.03 \text{ GeV}^2$  represent events with two pions and only one photon, emitted in the final state. Events of this type have  $(M_{\pi\pi}^0)^2 = s$ , there is no hard photon from initial state radiation present. Since in the KLOE analyses, the maximum value of  $(M_{\pi\pi}^0)^2$  for which the cross sections are measured (0.95  $\text{GeV}^2$ ) is sufficiently smaller than  $s \simeq M_\phi^2$  of the DAΦNE collider, these *leading-order* final state radiation events need to be taken out in the analysis. By moving these events to  $(M_{\pi\pi}^0)^2 = s$ , the passage from  $M_{\pi\pi}^2$  to  $(M_{\pi\pi}^0)^2$  automatically performs this task. Fig. 61 shows the fraction of events from *leading-order* final state radiation contributing to the total number of events, evaluated with the PHOKHARA event generator. Since in the *small angle* analysis the angular regions for pions and photons are separated, final state radiation, for which the photons are emitted preferably along the direction of the pions, is suppressed to less than 0.5%. Using *large angle* acceptance cuts, the effect is much bigger, especially above and below the  $\rho$ -resonance, where it can reach 20-30%. The correction of the shift in  $M_{\pi\pi}^2$  depends on the implementation of final state radiation in the Monte Carlo generator in terms of model dependence and missing contributions. It also relies on the correct assignment of photons coming from the initial or the final state, however, in case of symmetrical cuts in  $\theta_\gamma$ , interference effects between the two states vanish and the separation of initial and final state amplitudes is feasible.


- The acceptance in  $\theta_\gamma$ . Since the direction of the photons emitted in the final state is peaked along the direction of the pions, and the photons are emitted in the initial state along the  $e^+/e^-$  direction, the choice of the acceptance cuts affects the amount of final state radiation in the analyses. Using the *small angle* analysis cuts, a large part of final state radiation is suppressed by the separation of the pion and photon acceptance regions, and consequently needs to be reintroduced using corrections obtained from Monte Carlo simulations

to arrive at a result which is inclusive with respect to final state radiation (as needed in the dispersion integral for  $a_\mu^{\pi\pi}$ ). Even if in the *large angle* analysis the fraction of events with final state radiation surviving the selection is larger, again the missing part has to be added using Monte Carlo simulations. The acceptance correction for the cut in  $\theta_\gamma$  is evaluated for initial and final state radiation using the PHOKHARA generator, and the small differences found in the comparison of data and Monte Carlo distributions contribute to the systematic uncertainty of the measurement (see Table 13 and [445]).


- The distributions of kinematical variables. Cuts on the kinematical *trackmass* variable  $M_{\text{trk}}$  (see eq. 192), introduced in the analyses to remove background from the process  $e^+e^- \rightarrow \phi \rightarrow \pi^+\pi^-\pi^0$ , take out also a fraction of the events with final state radiation, necessitating a correction to obtain an inclusive result. Fig. 62 shows the effect final state radiation has on the distribution of the trackmass variable. The radiative tail of multi-photon events to the right of the peak at the  $\pi^\pm$  mass increases because the additional radiation moves events from the peak to higher values in  $M_{\text{trk}}$ . The width of the peak at  $M_{\pi^\pm}$  is due to the detector resolution, the plot was produced using the PHOKHARA event generator interfaced with the KLOE detector simulation [446]. Between 150 and 200 MeV, a  $M_{\pi\pi}^2$ -dependent cut is used in the event selection to reject the  $\pi^+\pi^-\pi^0$  events which have a value of  $M_{\text{trk}} > M_{\pi^\pm}$ . In this region, the cut also acts on the signal events. Missing terms concerning final state radiation in the Monte Carlo simulation or the non-validity of the pointlike-pion approximation used in PHOKHARA may affect the shape of the radiative tail in the trackmass variable. To overcome this, in the KLOE analyses, small corrections are applied to the momenta and the angles of the charged particles in the event in the simulation to obtain good agreement in the shape of  $M_{\text{trk}}$  for Monte Carlo simulation and data [445].
- The division by the radiator function  $H(s, M_{\pi\pi}^2)$ . In this case, one assumes perfect factorization between the ISR and the FSR process. This has been tested by performing the analysis in an inclusive and exclusive approach with respect to final state radiation, the assumption was found to be valid within 0.2% [367, 447].

It has been argued that contributions from events with two hard photons in the final state, which are not included in the PHOKHARA generator, may have an effect on the analyses [374].

The effect of the direct decay  $\phi \rightarrow \pi^+\pi^-\gamma$  on the radiative return analysis has been already addressed in [341]. Running at  $\sqrt{s} \simeq 1.02$  GeV, the amplitude for the processes  $\phi \rightarrow (f_0(980) + f_0(600))\gamma \rightarrow \pi^+\pi^-\gamma$  interferes with the amplitude for the final state radiation process. Due to the yet unclear nature of the scalar states  $f_0(980)$  and  $f_0(600)$ , the effect on the  $\pi^+\pi^-\gamma(\gamma)$  cross section depends




**Fig. 61.** (a) Fraction of events with leading order final state radiation in the *small angle* selection:  $50^\circ < \theta_\pi < 130^\circ$  and  $\theta_\gamma < 15^\circ$  ( $\theta_\gamma > 165^\circ$ ). (b) Fraction of events with leading order final state radiation in the *large angle* selection:  $50^\circ < \theta_\pi < 130^\circ$  and  $50^\circ < \theta_\gamma < 130^\circ$ . The PHOKHARA generator was used to produce the plots.



**Fig. 62.** Modification of the distribution of the trackmass variable due to the presence of final state radiation (dark grey triangles) compared to the one with initial state radiation only (light grey triangles). The arrows indicate the region in which the  $M_{\pi\pi}^2$ -dependent cut is applied in the analysis. The plot was created with the PHOKHARA generator interfaced to the KLOE detector simulation [446].

on the model used to describe the scalar mesons. The possibility to simulate  $\phi$  decays together with the processes for initial and final state radiation has been implemented in the PHOKHARA event generator in [331], using two characteristic models for the  $\phi$  decays: the “no structure” model of [448] and the  $K^+K^-$  loop model of [449]. A refined version of the  $K^+K^-$  loop model [435] and the



**Fig. 63.** (a):  $d\sigma_{\pi\pi\gamma}^{(ISR+FSR+f_0+\varrho\pi)}/d\sigma_{\pi\pi\gamma}^{(ISR+FSR)}$  for  $\sqrt{s} = 1.019$  GeV. (b):  $d\sigma_{\pi\pi\gamma}^{(ISR+FSR+f_0+\varrho\pi)}/d\sigma_{\pi\pi\gamma}^{(ISR+FSR)}$  for  $\sqrt{s} = 1$  GeV. Both plots were produced with the PHOKHARA6.1 event generator using *large angle* acceptance regions for pions and photons, with model parameters for the  $f_0$  and  $\varrho\pi$  contributions found in [435, 437].

double vector resonance  $\phi \rightarrow \pi^\pm \varrho^\mp (\rightarrow \pi^\mp \gamma)$  have been included as described in [344]. Using parameter values for the different  $\phi$  decays found in the analysis of the neutral channel  $\phi \rightarrow (f_0(980) + f_0(600))\gamma \rightarrow \pi^0 \pi^0 \gamma$  [435, 437], one can estimate the effect on the different analyses. While in the *small angle* analysis, there is no significant effect due to the choice of the acceptance cuts, in the *large angle* selection, the effect is of the order of several percent, and can reach up to 20% in the vicinity of the  $f_0(980)$ , as shown in Fig. 63, (a). While this allows to study the different models for the direct decays of  $\phi$ -mesons (see also Sec. 4.3.2), it prevents a precise measurement of  $\sigma_{\pi\pi}$  until the model and the parameters are understood with better uncertainty. An obvious way out is to use data taken at a value of  $\sqrt{s}$  outside the narrow peak of the  $\phi$  resonance ( $\Gamma_\phi = 4.26 \pm 0.04$  MeV [259]). In 2006, the KLOE experiment has taken  $\sim 250 \text{ pb}^{-1}$  of data at  $\sqrt{s} = 1$  GeV, 20 MeV below  $M_\phi$ . As can be seen in Fig. 63 (b), this reduces the effect due to contributions from  $f_0\gamma$  and  $\varrho\pi$  decays of the  $\phi$ -meson to within  $\pm 1\%$ .

#### Normalization with muon events

An alternative method to extract the pion form factor is to normalize the differential cross section  $d\sigma_{\pi\pi\gamma(\gamma)}/dM_{\pi\pi}^2$  directly to the process  $e^+e^- \rightarrow \mu^+\mu^-\gamma(\gamma)$ ,  $d\sigma_{\mu\mu\gamma(\gamma)}/dM_{\mu\mu}^2$ , in each bin of  $\Delta M_{\pi\pi}^2 = \Delta M_{\mu\mu}^2$ . Radiative corrections like the effect of vacuum polarisation, the radiator function and also the integrated luminosity  $\int L dt$  cancel out in the


ratio of pions over muons, only the effects from final state radiation (which is different for pions and muons) need to be taken into account consistently. An approach currently under way at KLOE uses the following equation to obtain  $|F_{2\pi}|^2$ :

$$|F_{2\pi}(s')|^2 \cdot (1 + \eta(s')) = \frac{4(1 + 2m_\mu^2/s')\beta_\mu}{\beta_\pi^3} \cdot \frac{(\frac{d\sigma_{\pi\pi\gamma(\gamma)}}{dM_{\pi\pi}^2})^{ISR+FSR}}{(\frac{d\sigma_{\mu\mu\gamma(\gamma)}}{dM_{\mu\mu}^2})^{ISR}} \quad (203)$$

In this formula, the measured differential cross section  $d\sigma_{\pi\pi\gamma(\gamma)}/dM_{\pi\pi}^2$  should be inclusive with respect to pionic final state radiation, while the measured cross section  $d\sigma_{\mu\mu\gamma(\gamma)}/dM_{\mu\mu}^2$  should be exclusive for muonic final state radiation.  $s' = M_{\pi\pi}^2 = M_{\mu\mu}^2$  is the squared invariant mass of the di-pion or the di-muon system after the respective corrections concerning final state radiation. Using this approach one gets on the left-hand side the pion form factor times the factor  $(1 + \eta(s'))$ , which describes the effect of the pionic final state radiation. This *bare* form factor is the quantity needed in the dispersion integral for the  $\pi\pi$ -contribution to  $a_\mu^{\text{had}}$ . While the measurement of  $d\sigma_{\pi\pi\gamma(\gamma)}/dM_{\pi\pi}^2$  and its corrections for pionic final state radiation are very similar to the one using the normalization with Bhabha events already performed at KLOE, the corrections needed to subtract the muonic final state radiation from the  $d\sigma_{\mu\mu\gamma(\gamma)}/dM_{\mu\mu}^2$  cross section are pure QED and can be obtained from the PHOKHARA generator, which includes final state radiation for muon pair production at next-to-leading order [330]. Due to the fact that the KLOE detector does not provide particle IDs, pions and muons have to be separated and identified using kinematical variables (e.g. the aforementioned trackmass variable) [361]. The analysis is in progress and a systematic precision similar to the one obtained in the absolute measurement is expected.

#### 4.4.2 Radiative return at BaBar

The BaBar radiative return program aims for the study of all significant hadronic processes in electron-positron annihilation  $e^+e^- \rightarrow \text{hadrons}$  for energies from threshold to about 4.5 GeV. Moreover, hadron spectroscopy of the initial  $J^{PC} = 1^{--}$  states, which are produced in  $e^+e^-$  collision, and of their decay products is performed. In this chapter BaBar results for processes with 3, 4, 5 and 6 hadrons in the final state, as well as measurements of baryon form factors in the time-like region are reported. A precision analysis of the pion form factor, i.e. of the cross section  $e^+e^- \rightarrow \pi^+\pi^-$ , which is essential for an improved determination of the hadronic contribution to the anomalous magnetic moment of the muon, appeared shortly before the publication of this paper [450]. The results presented in this chapter are based on a total integrated luminosity of  $230 \text{ fb}^{-1}$ , except the  $3\pi$  and  $4$  hadrons channels of ref. [391], which were analysed using a data sample of  $90 \text{ fb}^{-1}$ . The total BaBar data sample collected between the years 1999 to 2008 amounts to  $530 \text{ fb}^{-1}$ . A typical feature common to all radiative return analyses at BaBar



**Fig. 64.** BaBar measurement of the energy dependence of the  $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$  cross section obtained by radiative return in comparison with the world data set.

is a wide coverage of the entire mass range of interest in one single experiment with reduced point-by-point uncertainties compared to previous experiments.


#### $e^+e^- \rightarrow 3 \text{ Pions}$

The  $\pi^+\pi^-\pi^0$  mass spectrum has been measured from 1.05 GeV up to the  $J/\psi$  mass region with a systematic error of  $\sim 5\%$  below 2.5 GeV and up to  $\sim 20\%$  at higher masses [390]. The spectrum is dominated by the  $\omega$ ,  $\phi$  and  $J/\psi$  resonances. The BaBar measurement was able to significantly improve the world knowledge on the excited  $\omega$  states. The spectrum has been fitted up to 1.8 GeV and the following results for the masses and widths of the  $\omega'$  and  $\omega''$  states have been found:  $M(\omega') = (1350 \pm 20 \pm 20)$  MeV,  $\Gamma(\omega') = (450 \pm 70 \pm 70)$  MeV,  $M(\omega'') = (1660 \pm 10 \pm 2)$  MeV,  $\Gamma(\omega'') = (230 \pm 30 \pm 20)$  MeV. Note that below 1.4 GeV results of BaBar are in good agreement with those from SND [284] While above this energy cross sections measured by BaBar are much higher than those from DM2 [451].

#### $e^+e^- \rightarrow 4 \text{ Hadrons}$

The  $\pi^+\pi^-\pi^+\pi^-$ ,  $K^+K^-\pi^+\pi^-$  and  $K^+K^-K^+K^-$  exclusive final states have been measured from threshold up to 4.5 GeV with systematic errors of 5%, 15% and 25%, respectively [391]. The  $K^+K^-K^+K^-$  measurement is the first measurement of this process ever. Fig. 64 shows the mass distribution of the  $\pi^+\pi^-\pi^+\pi^-$  channel. We identify a huge improvement with respect to previous experiments. Background is relatively low for all channels under study (e.g. few percent at 1.5 GeV for  $\pi^+\pi^-\pi^+\pi^-$ ) and is dominated by ISR-events of higher multiplicities and of continuum non-ISR events at higher masses. The  $\pi^+\pi^-\pi^+\pi^-$  final state is dominated by the two-body  $a_1(1260)\pi$  intermediate state; the  $K^+K^-\pi^+\pi^-$  final state shows no significant two-body states, but rich three-body structure, including  $K^*(890)K\pi$ ,  $\phi\pi\pi$ ,  $\rho K K$  and  $K_2^*(1430)K\pi$ .

Fig. 65 shows BaBar preliminary results for the process



**Fig. 65.** Preliminary BaBar data for the  $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$  cross section in comparison with previous experiments.

$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ . The current systematic error of the measurement varies from 8% around the peak of the cross section to 14% at 4.5 GeV. BaBar results are in agreement with SND [452] in the energy range below 1.4 GeV and show a huge improvement for higher energies ( $> 1.4$  GeV). In the energy range above 2.5 GeV this is the first measurement ever. The  $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$  final state is dominated by the  $\omega\pi^0$ ,  $a_1(1260)\pi$  and  $\rho^+\rho^-$  intermediate channels, where the latter channel has been observed for the first time.

A specific analysis was devoted to the intermediate structures in the  $e^+e^- \rightarrow K^+K^-\pi^+\pi^-$  and  $e^+e^- \rightarrow K^+K^-\pi^0\pi^0$  channels [395]. Of special interest is the intermediate state  $\phi f_0(980)$ , where the decays  $f_0(980) \rightarrow \pi^+\pi^-$  and  $f_0(980) \rightarrow \pi^0\pi^0$  have been looked at. A peak is observed in the  $\phi f_0(980)$  channel at a mass  $M = 2175 \pm 18$  MeV and a width  $\Gamma = 58 \pm 2$ . The new state is usually denoted as  $Y(2175)$  and is also clearly visible in the  $K^+K^-f_0$  spectrum.

#### $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0, 2(\pi^+\pi^-)\eta$


The  $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$  cross section has been measured by BaBar from threshold up to 4.5 GeV [397]. A large coupling of the  $J/\psi$  and  $\psi(2S)$  to this channel is observed. The systematic error of the measurement is about 7% around the peak of the mass spectrum. In the  $\pi^+\pi^-\pi^0$  mass distribution the  $\omega$  and  $\eta$  peaks are observed; the rest of the events have a  $3\pi\rho$  structure.

BaBar performed also the first measurement of the  $e^+e^- \rightarrow 2(\pi^+\pi^-)\eta$  cross section. A peak value of about 1.2 nb at about 2.2 GeV is observed, followed by a monotonic decrease towards higher energies. Three intermediate states are seen:  $\eta\rho(1450)$ ,  $\eta'\rho(770)$  and  $f_1(1285)\rho(770)$ .

#### $e^+e^- \rightarrow 6 \text{ Hadrons}$

The 6 hadrons final state has been measured in the exclusive channels  $3(\pi^+\pi^-)$ ,  $2(\pi^+\pi^-)2\pi^0$  and  $K^+K^-2(\pi^+\pi^-)$  [393]. The cross section in the last case has never been mea-






**Fig. 66.** The energy dependence of the  $e^+e^- \rightarrow 3(\pi^+\pi^-)$  (upper plot) and the  $e^+e^- \rightarrow 2(\pi^+\pi^-)2\pi^0$  (lower plot) cross section obtained by BaBar (filled circles) by radiative return in comparison with previous data.

sured before; the precision in the first two cases is  $\sim 20\%$ , which is a large improvement with respect to existing data. Again, the entire energy range from threshold up to 4.5 GeV is measured in one single experiment. The distributions for the final states  $3(\pi^+\pi^-)$  and  $2(\pi^+\pi^-)2\pi^0$  are shown in Fig. 66. A clear dip is visible at about 1.9 GeV in both pion modes. A similar feature was already seen by FOCUS [453] in the diffractive photo-production of six charged pions. The spectra are fitted by BaBar using the sum of a Breit-Wigner resonance function and a Jacob-Slansky continuum shape. For the  $3(\pi^+\pi^-)$  ( $2(\pi^+\pi^-)2\pi^0$ ) mode, BaBar obtains values of  $1880 \pm 30$  MeV ( $1860 \pm 20$  MeV) for the resonance peak,  $130 \pm 30$  MeV ( $160 \pm 20$  MeV) for the resonance width and  $21^\circ \pm 14^\circ$  ( $-3^\circ \pm 15^\circ$ ) for the phase shift between the resonance and continuum.

$$e^+e^- \rightarrow K^+K^-\pi^0, K^+K^-\eta, K_S K^\pm \pi^\mp$$

A recent BaBar ISR-analysis is dedicated to three hadrons in the final state, including a pair of kaons ( $K^+K^-\pi^0$ ,  $KK_S\pi$ ); a peak near 1.7 GeV, which is mainly due to the  $\phi'(1680)$  state, is observed. A Dalitz plot analysis shows that the  $KK^*(892)$  and  $KK_2^*(1430)$  intermediate



**Fig. 67.** The  $e^+e^- \rightarrow p\bar{p}$  cross section measured by BaBar (filled circles) in comparison with data from other  $e^+e^-$  colliders (blue points) and from  $p\bar{p}$  experiments (red points).

states are dominating the  $K\bar{K}\pi$  channel. A fit to the  $e^+e^- \rightarrow K\bar{K}\pi$  cross section assuming the expected contributions from the  $\phi, \phi', \phi'', \rho^0, \rho', \rho''$  states was performed. The parameters of the  $\phi'$  and other excited vector meson states are compatible with PDG values.

*Time-like proton form factor  $e^+e^- \rightarrow p\bar{p}$ , hyperon form factors  $e^+e^- \rightarrow \Lambda^0\bar{\Lambda}^0, \Lambda^0\bar{\Sigma}^0, \Sigma^0\bar{\Sigma}^0$*

BaBar has also performed a measurement of the  $e^+e^- \rightarrow p\bar{p}$  cross section [392]. This time-like process is parametrised by the electric and magnetic form factors  $G_E$  and  $G_M$ :

$$\sigma_{e^+e^- \rightarrow p\bar{p}} = \frac{4\pi\alpha^2\beta C}{3s} \times (|G_M|^2 + \frac{2m_p^2}{s}|G_E|^2),$$

where  $\beta = \sqrt{1 - 4m_p^2/s}$  and the factor  $C = y/(1 - e^{-y})$  (with  $y = \pi\alpha m_p/(\beta\sqrt{s})$ ) accounts for the Coulomb interaction of the final state particles. The proton helicity angle  $\theta_p$  in the  $p\bar{p}$  rest frame can be used to separate the  $|G_E|^2$  and  $|G_M|^2$  terms. Their respective variations are approximately  $\sim \sin^2\theta_p$  and  $\sim (1 + \cos^2\theta_p)$ . By fitting the  $\cos\theta_p$  distribution to a sum of the two terms, the ratio  $|G_E|/|G_M|$  can be extracted. This is done separately in six bins of  $M_{p\bar{p}}$ . The results disagree significantly with previous measurements from LEAR [454] above threshold. BaBar observes a ratio  $|G_E|/|G_M| > 1$  above threshold, while at larger values of  $M_{p\bar{p}}$  the BaBar measurement finds  $|G_E|/|G_M| \approx 1$ . LEAR data on the contrary shows a behaviour  $|G_E|/|G_M| < 1$  above threshold.

In order to compare the cross section measurement with previous data ( $e^+e^-$  and  $p\bar{p}$  experiments), the *effective* form factor  $G$  is introduced:  $G = \sqrt{|G_E|^2 + 2m_p^2/s|G_M|^2}$ . The BaBar measurement of  $G$  is in good agreement with existing results, as can be seen in Fig. 67. The structure



of the form factor is rather complicated; the following observations can be made: (i) BaBar confirms an increase of  $G$  towards threshold as seen before by other experiments; (ii) two sharp drops of the spectrum at  $M_{p\bar{p}} = 2.25$  and  $3.0$  GeV are observed; (iii) data at large values  $M_{p\bar{p}} > 3$  GeV is in good agreement with the prediction from perturbative QCD.

A continuation of the ISR program with baryon final states is the measurement of the  $e^+e^- \rightarrow \Lambda\bar{\Lambda}$  cross section [398]. So far only one data point from DM2 [455] was existing for this channel, which is in good agreement with BaBar data. About 360  $\Lambda\bar{\Lambda}$  events could be selected using the  $\Lambda \rightarrow p\pi$  decay. In two invariant mass bins an attempt has been made to extract the ratio of the electric to magnetic form factor  $|G_E|/|G_M|$ . In the mass range below 2.4 GeV this ratio is above unity - as in the proton case - with a significance of one standard deviation ( $|G_E|/|G_M| = 1.73^{+0.99}_{-0.57}$ ). Above 2.4 GeV the ratio is consistent with unity ( $|G_E|/|G_M| = 0.71^{+0.66}_{-0.71}$ ). Also the  $\Lambda$  polarisation and the phase between  $G_E$  and  $G_M$  was studied using the slope of the angle between the polarisation axis and the proton momentum in the  $\Lambda$  rest frame. The following limit on  $\Lambda$  polarisation is obtained:  $-0.22 < \zeta < 0.28$ ; the relative phase between the two form factors is measured as  $-0.76 < \sin(\phi) < 0.98$ , which is not yet significant due to limited statistics.

Finally the first measurements of the  $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$  and  $e^+e^- \rightarrow \Sigma^0\bar{\Lambda}(\Lambda\bar{\Sigma}^0)$  cross sections were performed. For the detection of the  $\Sigma^0$  baryon, the decay  $\Sigma^0 \rightarrow \Lambda\gamma \rightarrow p\pi\gamma$  was used. About 40 candidate events were selected for the  $\Sigma^0\bar{\Sigma}^0$ , about 20 events for the  $\Lambda\bar{\Sigma}^0$  reaction. All baryon form factors measured by BaBar have a similar size and mass shape, namely a rise towards threshold. The reason for this peculiar behaviour is not understood.

#### 4.4.3 Radiative return at Belle

##### *ISR studies at Belle*

Until now most of the Belle analyses using radiative return focused on studies of the charmonium and charmonium-like states. They can be subdivided into final states with open and hidden charm.

##### *Final states with open charm*

Belle performed a systematic study of various exclusive channels of  $e^+e^-$  annihilation into charmed mesons and baryons using ISR often based on the so called partial reconstruction to increase the detection efficiency and suppress background.

In Ref. [404] they measured the cross sections of the processes  $e^+e^- \rightarrow D^{*\pm}D^{*\mp}$  and  $e^+e^- \rightarrow D^+D^{*-} + c.c.$ . The shape of the former is complicated and has several local maxima and minima. The first two maxima are close to the  $\psi(4040)$  and  $\psi(4160)$  states. The latter shows significant excess of events near the  $\psi(4040)$ .

The cross sections of the processes  $e^+e^- \rightarrow D^+D^-$  and  $e^+e^- \rightarrow D^0\bar{D}^0$  show a signal of the  $\psi(3770)$  as well

as hints of the  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  [407]. There is also an enhancement near 3.9 GeV, which qualitatively agrees with the prediction of the coupled channel model [456].

The cross section of the process  $e^+e^- \rightarrow D^0D^-\pi^+$  has a prominent peak at the energy corresponding to the  $\psi(4415)$  [410]. From a study of the resonant substructure in the decay  $\psi(4415) \rightarrow D^0D^-\pi^+$  they conclude that it is dominated by the intermediate  $D\bar{D}^*(2460)$  mechanism.

In contrast to expectations of some hybrid models predicting  $Y(4260) \rightarrow D^{(*)}\bar{D}^{(*)}\pi$  decays, no clear structures were observed in the cross section of the process  $e^+e^- \rightarrow D^0D^{*-}\pi^+$  [457]. There is only some evidence ( $\sim 3.1\sigma$ ) for the  $\psi(4415)$ .

Finally, they measure the cross section of the reaction  $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$  and observe a significant peak near threshold that they dub  $X(4630)$  [411]. Assuming that the peak is a resonance, they find that its mass and width are compatible within errors with those of the  $Y(4660)$  state found by Belle in the  $\psi(2S)\pi^+\pi^-$  final state via ISR [406]. However, interpretations other than  $X(4630) \equiv Y(4660)$  cannot be excluded. For example, peaks at the baryon-antibaryon threshold are observed in various processes [458]. According to other assumptions, the  $X(4630)$  is a  $5^3S_1$  [459] or  $\psi(6S)$  [460] charmonium state or, for example, a threshold effect, which is due to the  $\psi(3D)$  slightly below the  $\Lambda_c^+\Lambda_c^-$  threshold [461]. Fig. 68 shows all mentioned above cross sections with the vertical lines showing positions of both well established states like  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  and new charmonium-like states  $Y(4008)$ ,  $Y(4260)$ ,  $Y(4360)$  and  $Y(4660)$  discussed below.

Summing the measured cross sections and taking into account not yet observed final states on base of isospin symmetry they find that until about 4.3 GeV the sum of exclusive cross sections almost saturates the total inclusive cross section measured by BES [297].

##### *Final states with hidden charm*

Studying the  $J/\psi\pi^+\pi^-$  final state Belle confirmed the  $Y(4260)$  discovered by BaBar and in addition observed a new structure dubbed  $Y(4008)$  [405], see Fig. 69. They also observe the reaction  $e^+e^- \rightarrow J/\psi K^+K^-$  and find first evidence for the reaction  $e^+e^- \rightarrow J/\psi K_S^0 K_S^0$  [408].


Studying the  $\psi(2S)\pi^+\pi^-$  final state Belle confirmed the  $Y(4360)$  discovered by BaBar and in addition observed a new structure dubbed  $Y(4660)$  [406], see Fig. 70.

It is worth noting that the resonance interpretation of various enhancements discussed above is not unambiguous and can be strongly affected by close thresholds of different final states and rescattering effects.


Various ISR studies performed at the Belle detector in the charmonium region are summarized in Table 14.

##### *ISR studies of light quark states*

In one case the ISR method was used to study the light quark states [462]. In this analysis the cross sections




**Fig. 68.** Cross sections of various exclusive processes measured by Belle: a)  $e^+e^- \rightarrow D\bar{D}$ , b)  $e^+e^- \rightarrow D^+D^{*-} + c.c.$ , c)  $e^+e^- \rightarrow D^{*\pm}D^{*\mp}$ , d)  $e^+e^- \rightarrow D^0D^-\pi^+$ , e)  $e^+e^- \rightarrow D^0D^{*-}\pi^+$ , and f)  $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ . The dashed lines show the position of the  $\psi$  states while the dotted lines correspond to the  $Y(4008)$ ,  $Y(4260)$ ,  $Y(4360)$ , and  $Y(4660)$  states.



**Fig. 69.** The  $J/\psi\pi^+\pi^-$  invariant mass distribution.


of the reactions  $e^+e^- \rightarrow \phi\pi^+\pi^-$  and  $e^+e^- \rightarrow \phi f_0(980)$  are measured from threshold to 3 GeV using a data sample of  $673 \text{ fb}^{-1}$ , see Fig. 71(a,b). In the  $\phi\pi^+\pi^-$  mode the authors observe and measure for the first time the parameters of the  $\phi(1680)$ , they also observe and measure the parameters of the  $\phi(2170)$ . Also selected in this analysis is the  $\phi f_0(980)$  final state, which shows a clear signal of the  $\phi(2170)$ . For Monte Carlo simulation they use a version of PHOKHARA in which the produced resonance decays into  $\phi\pi^+\pi^-$  or  $\phi f_0(980)$  with the subsequent decays



**Fig. 70.** The  $\psi(2S)\pi^+\pi^-$  invariant mass distribution

**Table 14.** Summary of ISR studies in the  $c\bar{c}$  region at Belle

Final state	$\int L dt, \text{fb}^{-1}$	Ref.
$D^{*+}D^{*-}$	547.8	[404]
$D^\pm D^{*\mp}$	547.8	[404]
$D^0 D^0, D^+ D^-$	673	[407]
$D^0 D^-\pi^+$	673	[410]
$D^0 D^{*-}\pi^+$	695	[457]
$\Lambda_c^+ \Lambda_c^-$	695	[411]
$J/\psi\pi^+\pi^-$	548	[405]
$\psi(2S)\pi^+\pi^-$	673	[406]
$J/\psi K^+ K^-$	673	[408]



**Fig. 71.** The cross sections of the processes  $e^+e^- \rightarrow \phi\pi^+\pi^-$  (a) and  $e^+e^- \rightarrow \phi f_0(980)$  (b)

$\phi \rightarrow K^+K^-$  and  $f_0(980) \rightarrow \pi^+\pi^-$ . The  $\pi^+\pi^-$  system is in  $S$ -wave, the  $\pi^+\pi^-$  system and the  $\phi$  are also in a relative  $S$ -wave. The  $\pi^+\pi^-$  mass distribution is generated according to phase space. They assign a 0.1% as the systematic uncertainty of the ISR photon radiator.

In all the ISR studies the Monte Carlo simulation is performed as follows. First, the kinematics of the initial-state radiation is generated using the PHOKHARA 5.0 package for simulation of the process  $e^+e^- \rightarrow V\gamma_{\text{ISR}}(\gamma_{\text{ISR}})$  [332]. Then a  $q\bar{q}$  generator is used to generate  $V$  decays.

#### 4.4.4 Prospects for radiative return at VEPP-2000

It is well known that the main hadronic contribution to  $a_\mu^{\text{had}}$  comes from the energy region below 1 GeV and is dominated by the  $\pi^+\pi^-$  channel. Direct scan experiments with CMD-3 at VEPP-2000 will deliver huge statistics, but the accuracy of the cross section determination will be

determined by systematic effects. So, any other possibility to measure the pion form factor, for example with ISR, will be a valuable instrument to provide a cross check for better understanding of the systematics.

The design luminosity  $\sim 10^{32}\text{cm}^{-2}\text{s}^{-1}$  is expected at  $\sqrt{s} = 2$  GeV. As a result, the statistics similar to that of CMD-2 will be collected. Let us recollect that the ISR method provides a “low energy scan” while data taking occurs at fixed high energy. The threshold region,  $2m_{\pi} - 0.5$  GeV, gives about 13% of the total contribution to the muon anomaly. To overcome the lack of data at threshold energies, the ISR method can serve as very efficient and unique way to measure the pion form factor.


Trigger and reconstruction efficiencies, detector imperfections will be identical for the whole energy range. As a result, some systematic errors will be canceled out at least partially and the total systematic error will be lower. Measurements of the cross section of the process  $e^+e^- \rightarrow \mu^+\mu^-$  must confirm the validity of this method and will allow one to determine an energy scale and some systematic errors. A fit of the  $\omega$  and  $\phi$  resonances will also provide calibration of the energy scale. The ISR method provides an additional instrument to better understand and estimate the systematics. Let us recollect that we are going to achieve a systematic accuracy for the pion form factor of a few pro mille.

In direct scan experiments the data as a rule are collected at fixed energy points. Thus some “empty” regions without data naturally arise. An important feature of experiments with ISR is that the whole energy scale will be covered filling any existing gaps.

Currently, the theoretical error for the cross section of the process  $e^+e^- \rightarrow \pi^+\pi^-\gamma$  is dominated by the uncertainty of the radiator function (0.5%) and there is hope to reduce it to a few pro mille in future by theorists. In the case of the pion form factor extraction from the  $\pi^+\pi^-\gamma/\mu^+\mu^-\gamma$  ratio, the dependence on theory will be significantly reduced since the main uncertainty of the radiator function and vacuum polarization effects cancel out in the ratio. With the integrated luminosity of several inverse femtobarn at 2 GeV one can reach a fractional accuracy on the total error better than 0.5%.

#### 4.4.5 Prospects for radiative return at BES-III

The designed peak luminosity of the BEPC-II is  $1 \times 10^{33}\text{cm}^{-2}\text{s}^{-1}$  at  $\sqrt{s} = 3.77$  GeV, i.e., the  $\psi(3770)$  peak. It has reached 30% of the designed luminosity now and is starting to deliver luminosity to BES-III for physics. Although the physics programs at BES-III are rather rich [43], most of the time, the machine will run at  $\sqrt{s} = 3.77$  GeV and 4.17 GeV for charm physics, since the cross sections of  $J/\psi$  and  $\psi(2S)$  production are large and the required statistics can be accumulated in short time, say, one year at each energy point. An estimation of the BEPC-II running time at  $\sqrt{s} = 3.77$  GeV and 4.17 GeV will be around 8 years, it corresponds to an integrated luminosity of about  $20\text{fb}^{-1}$  at each energy point.




**Fig. 72.** Expected luminosity at low energies due to ISR for  $10\text{fb}^{-1}$  data accumulated on the  $\psi(3770)$  peak.

Data samples at  $\sqrt{s} = 3.77$  GeV and 4.17 GeV can be used for radiative return study, for the center of mass energies of the hadron system between  $\pi^+\pi^-$  threshold to above 2.0 GeV. This will allow a measurement of the pion form factor, the kaon form factor, and the proton form factor, as well as the cross sections of some multi-hadron final states. The good coverage of the muon detector at the BES-III also allows an identification of the  $\mu^+\mu^-$  final state, thus supply as a normalization factor to the other two-body final states.

Fig. 72 shows the expected luminosity at low energies in 10 MeV bin for  $10\text{fb}^{-1}$  data accumulated on the  $\psi(3770)$  peak. In terms of luminosity at the  $\rho^0$  peak, one can see that  $10\text{fb}^{-1}$  data at  $\sqrt{s} = 3.77$  GeV is equivalent to  $70\text{fb}^{-1}$  data at 10.58 GeV, namely, the B-factories.

With Monte Carlo generated  $e^+e^- \rightarrow \gamma_{ISR}\pi^+\pi^-$  data using PHOKHARA [327], after a fast simulation and reconstruction with the BES-III softwares, one found the efficiency for events at the  $\rho^0$  peak is around 5% if one requires the ISR photon is detected, this is higher than the efficiency at the BaBar experiment [463]. Fig. 73 shows the signal for 10,000 generated  $\pi^+\pi^-$  events. One estimates the number of events in each 10 MeV bin is around 20,000 at the  $\rho^0$  peak for  $10\text{fb}^{-1}$  data at  $\sqrt{s} = 3.77$  GeV. This is comparable to the recent BaBar results based on  $232\text{fb}^{-1}$  data at the  $\Upsilon(4S)$  peak [463].



**Fig. 73.** Detected  $\gamma_{\text{ISR}}\pi^+\pi^-$  in 10000 produced events at the  $\psi(3770)$  peak. The sample is generated with PHOKHARA.

The most important work related to the pion form factor measurement is the estimation of the systematic error. Since the cross section of good events at  $\psi(3770)$  peak is not large (around 30 nb for the total hadronic cross section, with about 400 nb cross section for the QED processes) compared to the highest trigger rate energies  $J/\psi$  and  $\psi(2S)$  peaks, a loose trigger to allow the ISR events being recorded is out of question. In principle, the trigger rate for these events could reach 100% with an allowed trigger purity of less than 20%.

With enough  $D\bar{D}$  events accumulated at the same energy, the tracking efficiency and particle ID efficiency can be measured in high precision (as has been done at the CLEO-c [464]); in addition, a huge data sample at  $\psi(2S)$  and the well measured large branching fraction  $\psi(2S)$  transition modes, such as  $\pi^+\pi^-J/\psi$ ,  $J/\psi \rightarrow \mu^+\mu^-$ , can be used to study the tracking efficiency,  $\mu$ -ID efficiency and so on. All these will help greatly in understanding the detector performance, and in pinning down the systematic errors in the form factor measurement.

The kaon and proton form factors can be measured as well since they are even simpler than the measurement of pion form factor. This will allow us a better understanding of the structure close to the threshold and the possible existing high mass structures.

Except for the lowest lying vector states ( $\rho$ ,  $\omega$ , and  $\phi$ ), the parameters of other vector states are poorly known and further investigations are needed. BES-III ISR analyses may reach a bit above 2 GeV, while above that, BEPC-II can run directly by setting beam energy there. This allows BES-III a full reach of the vectors between  $\pi^+\pi^-$  threshold and 4.6 GeV, the highest energy BEPC-II can reach, covering the  $\rho$ ,  $\omega$ ,  $\phi$ , as well as the  $\psi$  sectors. Here one will have chance to study the excited  $\rho$ ,  $\omega$ , and  $\phi$  states between 1 and about 2.5 GeV. The final states include  $\pi^+\pi^-\pi^0$ ,  $K\bar{K}$ , 4 pions,  $\pi\pi K\bar{K}$ , etc. Final states with more than four particles will be hard to study using ISR method, since the  $D\bar{D}$  decay will contribute as background.

## 5 Tau decays

### 5.1 Introduction

After the  $\tau$  lepton discovery, a fundamental lepton, which is heavy enough to decay not only into leptons, but also into dozens of various hadronic final states, it became clear that corresponding Monte Carlo (MC) event generators are needed for various purposes:

- To calculate detector acceptance, efficiencies and various distributions for signal event selection and comparison to data. The acceptance is small (a few percent) and depends on the model; in principle, it is a complicated function of invariant masses, angles, and resolutions. Analysis of publications shows that effects of MC signal modelling are almost always neglected
- To estimate the number of background (BG) events  $N_{\text{ev}}^{\text{BG}}$  and their distributions; in addition to background coming from  $\tau^+\tau^-$  pairs (so called cross-feed), there might be BG events from  $q\bar{q}$  continuum,  $\gamma\gamma$  collisions etc.
- To unfold observed distributions to get rid of detector effects, important while extracting resonance parameters

Various computer packages like, e.g., KORALB [465], KKMC [466], TAUOLA [467, 468, 469], PHOTOS [470] were developed to generate events of  $\tau$  lepton production in  $e^+e^-$  annihilation and their subsequent decay taking into account a possibility of photon emission. These codes became very important tools for experiments at LEP, CLEO, Tevatron and HERA.

Simulation of hadronic decays requires knowledge of hadronic form factors. Various final hadronic states were considered in the 90's, resulting in a large number of specific hadronic currents [471].

However, already experiments at LEP and CLEO showed that with increase of the collected data sets a more precise description is necessary. Some attempts were made to improve the parameterisation of various hadronic currents. One should note the serious efforts of the ALEPH and CLEO Collaborations, which created their own parameterisations of TAUOLA hadronic currents already in the late 90's, or a parameterisation of the hadronic current in the  $4\pi$  decays [472] based on the experimental information on  $e^+e^- \rightarrow 2\pi^+2\pi^-$ ,  $\pi^+\pi^-2\pi^0$  from Novosibirsk [288], which are now implemented in the presently distributed TAUOLA code [473].

### 5.2 Current status of data and MC generators

In this section we will briefly discuss the most precise recent experimental data on  $\tau$  lepton decays showing wherever possible their comparison to the existing MC generators and discussing decay dynamics.



### 5.2.1 $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ at Belle

Recently results of a study of the  $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$  decay by the Belle Collaboration were published [474]. From less than 10% of the data set available the authors selected a huge statistics of 5.4M events, about two orders of magnitude larger than in any previous experiment, determined the branching fraction and after the unfolding obtained the hadronic mass spectrum, in which for the first time three  $\rho$ -like resonances were observed together:  $\rho(770)$ ,  $\rho(1450)$ ,  $\rho(1700)$ . Their parameters were also determined.


Comparison of the obtained missing mass distributions with simulation for different polar angle ranges (Fig. 74) shows that there exist small discrepancies between MC and data.

Fig. 75 shows various background contributions to the di-pion mass distribution (upper) and underlying dynamics (lower) clearly demonstrating a pattern of three interfering resonances:  $\rho(770)$ ,  $\rho(1450)$ ,  $\rho(1700)$ .

### 5.2.2 $\tau^- \rightarrow \bar{K}^0 \pi^- \nu_\tau$ , $K^- \pi^0 \nu_\tau$


Two high-precision studies of the  $\tau$  decay into the  $K\pi\nu_\tau$  final state were recently published. The BaBar Collaboration reported a measurement of the branching fraction of the  $\tau^- \rightarrow K^- \pi^0 \nu_\tau$  decay [475]. They do not study in detail the  $K\pi$  invariant mass distribution noting only that the  $K^*(892)^-$  resonance is seen prominently above the simulated background, see Fig. 76. Near 1.4 GeV/ $c^2$  decays to higher  $K^*$  mesons are expected, such as the  $K^*(1410)^-$  and  $K_0^*(1430)^-$ , but their branching fractions are not yet well measured. These decays are not included in the BaBar simulation of  $\tau$  decays, but seem to be present in the data around 1.4 GeV/ $c^2$ . It is also worth noting that this decay mode is heavily contaminated by cross-feed backgrounds from other  $\tau$  decays. For example, below 0.7 GeV/ $c^2$  the background is dominated by  $K^- \pi^0 \pi^0 \nu_\tau$  and  $K^- K^0 \pi^0 \nu_\tau$  events, for which the branching fractions are only known with large relative uncertainties of  $\approx 37\%$  and  $\approx 13\%$ , respectively. Non-negligible background may also come from the  $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$  decay, which has a large branching fraction and thus should be simulated properly.

Another charge combination of the final state particles, i.e.,  $K_S^0 \pi^- \nu_\tau$ , was studied in the Belle experiment [476]. In this case a detailed analysis of the  $K\pi$  invariant mass distribution has been performed. The authors also conclude that decay dynamics differs from pure  $K^*(892)$ : the best fit includes  $K_0^*(800) + K^*(892) + K^*(1410)/K_0^*(1430)$ , Fig. 77.



**Fig. 74.** Projections to the missing mass and missing direction for  $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$  decay at Belle: (a)-(c) correspond to different ranges of the missing polar angles. The solid circles represent the data and the histogram – MC simulation (signal + background). The open histogram shows the contribution from  $\tau^+ \tau^-$  pairs, the vertical (horizontal) striped area shows that from two-photon leptonic (hadronic) processes; the wide (narrow) hatched area shows that from Bhabha ( $\mu^+ \mu^-$ ), and the shaded area that from the  $q\bar{q}$  continuum.






**Fig. 75.** Invariant-mass-squared distribution for  $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$  decay at Belle. (a) Contributions of different background sources. The solid circles with error bars represent the data, and the histogram represents the MC simulation (signal + background). (b) The fully corrected distribution. The solid curve is the result of a fit to the Gounaris-Sakurai model with the  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$  resonances.

### 5.2.3 $\tau$ decays into three pseudoscalars

Recently a measurement of the branching fractions of various particle combinations in the decay to three charged hadrons (any combination of pions and kaons) was reported by the BaBar Collaboration [477]. A similar study was also performed by the Belle group [478]. However, both groups have not yet analysed in detail the mass spectra. In the  $K^-K^+K^-\nu_\tau$  final state BaBar [477] and Belle [479] reported the observation of the decay mode  $\phi K^-\nu_\tau$  while in the  $K^-K^+\pi^-\nu_\tau$  final state BaBar observed the  $\phi\pi^-\nu_\tau$  decay mode [477]. Belle analysed the spectrum of the  $\phi K^-$  mass and concluded that it might have a complicated dynamics, see Fig. 78.



**Fig. 76.** The  $K\pi$  invariant mass distribution for  $\tau^- \rightarrow K^-\pi^0\nu_\tau$  decay at BaBar. The dots are the data, while the histograms are background MC events with selection and efficiency corrections:  $\tau$  background (dashed line),  $q\bar{q}$  (dash-dotted line),  $\mu^+\mu^-$  (dotted line).

The most detailed previous study of the mass spectra was done by the CLEO group [480]. With the statistics of about 8 000 events they conclude that the  $3\pi$  mass spectrum is dominated by the  $a_1(1260)$  meson and confirmed that the decay of the latter is not saturated by the  $\rho\pi$  intermediate state having in addition a significant  $f_0(600)\pi\pi^-$  component earlier observed in  $e^+e^-$  annihilation into four charged pions [288].

Recently the Belle Collaboration performed a detailed study of various decays with the  $\eta$  meson in the final state [481]. They measured the branching fractions of the following decay modes:  $\tau^- \rightarrow K^-\eta\nu_\tau$ ,  $\tau^- \rightarrow K^-\pi^0\eta\nu_\tau$ ,  $\tau^- \rightarrow \pi^-\pi^0\eta\nu_\tau$ ,  $\tau^- \rightarrow \pi^-K_S^0\eta\nu_\tau$ , and  $\tau^- \rightarrow K^{*-}\eta\nu_\tau$ . They also set upper limits on the branching fractions of the decays into  $K^-K_S^0\eta\nu_\tau$ ,  $\pi^-K_S^0\pi^0\eta\nu_\tau$ ,  $K^-\eta\eta\nu_\tau$ ,  $\pi^-\eta\eta\nu_\tau$ , and non-resonant  $K^-\pi^0\eta\nu_\tau$  final states.

Fig. 79 shows that there is reasonable agreement for  $\eta\pi^-\pi^0\nu_\tau$  (a, b) and a worse one for  $\eta K^-\nu_\tau$  (c) and  $\eta K^{*-}\nu_\tau$  (d).


### 5.2.4 $\tau$ decays to four pions

There are two possible isospin combinations of the final hadronic state,  $2\pi^-\pi^+\pi^0$  and  $\pi^-3\pi^0$ . Both have not yet been studied at  $B$  factories, so the best existing results are based on ALEPH [482] and CLEO [483] results.

Theoretical description of such decays is based on the CVC relations and available low energy  $e^+e^-$  data [484, 324, 472, 333].

### 5.2.5 $\tau^- \rightarrow 3h^-2h^+\nu_\tau$ at BaBar

A new study of the  $\tau^- \rightarrow 3h^-2h^+\nu_\tau$  decay ( $h = \pi, K$ ) has been performed by the BaBar Collaboration [485]. A



**Fig. 79.** The invariant mass distributions: (a)  $\pi\pi^0$  and (b)  $\pi\pi^0\eta$  for  $\tau \rightarrow \pi\pi^0\eta\nu_\tau$ ; (c)  $\eta K$  for  $\tau \rightarrow K\eta\nu_\tau$  and (d)  $\pi K_S^0\eta$  for  $\tau \rightarrow \pi K_S^0\eta\nu_\tau$  at Belle. The points with error bars are the data. The normal and filled histograms indicate the signal and  $\tau^+\tau^-$  background MC distributions, respectively.

large data set of over 34 000 events (two orders of magnitude higher than in the best previous measurement at CLEO [486]) allows one a first search for resonant structure and decay dynamics.

The invariant mass distribution of the five charged particles in Fig. 80 shows clear discrepancy between the data and the MC simulation, which uses the phase space distribution for  $\tau^- \rightarrow 3\pi^- 2\pi^+ \nu_\tau$ .

The mass of the  $h^+h^-$  pair combinations in Fig. 81 up, with a prominent shoulder at  $0.77 \text{ GeV}/c^2$  suggests a strong contribution from the  $\rho$  meson. Note that there are three allowed isospin states for this decay of which two may have a  $\rho$  meson. The mass of  $2h^+2h^-$  combinations in Fig. 81 down, also shows a structure at  $1.285 \text{ GeV}/c^2$  coming from the  $\tau^- \rightarrow f_1(1285)\pi^- \nu_\tau$  decay.

The first attempt to take into account the dynamics of this decay was recently performed in Ref. [487].

### 5.2.6 $\tau$ decays to six pions

The six-pion final state was studied by the CLEO Collaboration [488]. Two charge combinations,  $3\pi^- 2\pi^+ \pi^0$  and  $2\pi^- \pi^+ 3\pi^0$  were observed and it was found that decays are saturated by intermediate states with  $\eta$  and  $\omega$  mesons. Despite rather limited statistics (about 260 events altogether), it became clear that the dynamics of these decays is rather rich.

### 5.2.7 Lepton-Flavour Violating Decays

More than 50 different Lepton-Flavour Violating (LFV) decays have been studied by the CLEO, BaBar and Belle Collaborations. Publications rarely describe how simulation of such decays is performed. Moreover, theoretical papers suggesting LFV in new models usually do not provide differential cross sections. In some experimental papers the authors claim that the production of final state hadrons with a phase space distribution is assumed. However, the real meaning of this statement is not very clear since LFV




Fig. 77. The  $K\pi$  invariant mass distribution for  $\tau^- \rightarrow K^- \pi^0 \nu_\tau$  decay at Belle. Points are experimental data, histograms are spectra expected for different models. (a) shows the fitted result in the model with the  $K^*(892)$  alone. (b) shows the fitted result in the  $K^*(892) + K_0^*(800) + K^*(1410)$  model. Also shown are different types of background.




Fig. 78. The  $\phi K$  invariant mass distribution for  $\tau^- \rightarrow \phi K^- \nu_\tau$  decay at Belle. Points with error bars are the data. The open histogram is the phase-space distributed signal MC, and dotted and dot-dashed histograms indicate the signal MC mediated by a resonance with mass and width of 1650 MeV and 100 MeV, and 1570 MeV and 150 MeV, respectively.




Fig. 80. Invariant mass of five charged particles for  $\tau^- \rightarrow 3h^- 2h^+ \nu_\tau$  at BaBar





Fig. 81. Invariant mass distributions for  $\tau^- \rightarrow 3h^- 2h^+ \nu_\tau$  at BaBar, points with error bars are the data: upper -  $h^+ h^-$ , the unshaded and shaded histograms are the signal and background predicted by MC; lower -  $2\pi^+ 2\pi^-$ . The solid line is a fit to the data using a second-order polynomial (dashed line) for the background and a Breit-Wigner convoluted with a Gaussian for a peak region.

assumes New Physics and, therefore, matrix elements are not necessarily separated into weak and hadronic parts.

However, there exist a few theoretical papers considering differential cross sections. For example, angular correlations for  $\tau^- \rightarrow \mu^- \gamma$ ,  $\mu^- \mu^+ \mu^-$ ,  $\mu^- e^+ e^-$  decays were studied in Ref. [489]. An attempt to classify different types of operators entering New Physics Lagrangians for  $\tau$  decays to three charged leptons was made in [490].

### 5.3 Status of Monte Carlo event generators for $\tau$ production and decays

High-statistics and high-precision experiments as well as searches for rare processes result in a new challenge: Monte Carlo generators based on an adequate theoretical description of energy and angular distributions. In the following we will describe status of the Monte Carlo programs used by experiments. We will review building blocks used in the simulation with the goal in mind to localise the points requiring most urgent attention.

At present, for the production of  $\tau$  pairs the Monte Carlo programs KORALB [465] and KKMC [466] are standard codes to be used. For the generation of bremsstrahlung in decays PHOTOS [470] Monte Carlo is used. Finally,  $\tau$  decays themselves are simulated with TAUOLA [467, 468, 469]. The EvtGen code was written and maintained for simulation of  $B$  meson decays, see

[www.slac.stanford.edu/~lange/EvtGen/](http://www.slac.stanford.edu/~lange/EvtGen/). It offers a unique opportunity to specify at run time a list of the final state particles<sup>14</sup>, without having to change and/or compile the underlying code. In a multi-particle final state dominated by phase space considerations, this generator provides adequate description of the final state momenta, for which the underlying form factor calculation is more involved and not presently available in a closed form. That is why it is used by experiments measuring  $\tau$  decays too.

So far, our discussion has been based on comparison of experimental data and theory embodied into Monte Carlo results treated as a black box. One could see that a typical signature of any given  $\tau$  decay channel is matching rather poorly the publicly available Monte Carlo predictions. This should be of no surprise as efforts to compare data with predictions were completed for the last time in late 90's by the ALEPH and CLEO collaborations. The resulting hadronic currents were afterwards implemented in [473]. Since that time no efforts to prepare a complete parameterisation of  $\tau$  decay simulation for the public use were seriously undertaken.

There is another important message which can be drawn from these comparisons. Starting from a certain precision level, a study of the given decay mode can not be separated from a discussion of others. In the distributions aimed at representing the given decay mode, a contribution from the other  $\tau$  decay modes can be large, up to even 30%.

It may be less clear that experiments differ significantly in the way how they measure individual decay modes. For instance, ALEPH produced  $\tau$  samples free of the non- $\tau$

backgrounds, but, on the other hand, strongly boosted, making more difficult the reconstruction of some angles in the hadronic system. This is important and affects properties of the decay models which will be used for a parameterisation. In particular, when statistics is small, possible fluctuations may affect the picture and there is not enough data to complete an estimate of the systematic errors. In this case details of the hadronic current description, as the inclusion of intermediate resonances, are not important. Let us consider as an example  $\tau^- \rightarrow K_S^0 \pi^- \pi^0 \nu_\tau$ . The matrix element in the ALEPH parameterisation is saturated with  $\rho^- \rightarrow \pi^- \pi^0$  and  $K^{*0} \rightarrow K_S^0 \pi^0$ , and a similar parameterisation is used for  $K^{*-} \rightarrow K_S^0 \pi^-$ . In practice, a contribution of the  $\rho$  is more significant in the ALEPH parameterisation in contrast to the CLEO one where the  $K^*$  dominates. One has to admit that at the time when both collaborations were preparing their parameterisations to be used in TAUOLA, data samples of both experiments were rather small and the differences were of not much significance. This can, however, affect possible estimates of backgrounds for searches of rare decays, e.g., of  $B$  mesons at LHCb<sup>15</sup>.

Let us now go point by point and discuss examples of Monte Carlo programs and fitting strategies. We will focus on subjects requiring most attention and future work. We will review the theoretical constraints which are useful in construction of the models used for the data description.

### 5.4 Phase space

Because of a relatively low multiplicity of final state particles, it is possible to separate a description of  $\tau$  production and decay into segments describing the matrix elements and the phase space. In the phase space no approximations are used, contrary to the matrix elements where all approximations and assumptions reside. The description of the phase space used in TAUOLA is given in detail in [469]. The description of the phase space for  $\tau$  production is given in [466]. Thanks to conformal symmetry it is exact for an arbitrary number of photons. Using exponentiation, see, for example, Yennie-Frautchi-Suura [491], the phase space description can be exact and the matrix element can be refined order by order. For radiative corrections in decay PHOTOS can be used. Its phase space is described, for example, in the journal version of [492] and is exact. Approximations are in the matrix element only. Benchmarks<sup>16</sup> – comparison with other calculations

<sup>15</sup> LHCb performed MC studies for  $B_s^0 \rightarrow \mu^+ \mu^-$  and radiative decays  $B^0 \rightarrow K^* \gamma$ ,  $B_s^0 \rightarrow \phi \gamma$ , but  $\tau$  decays have not yet been taken into account. These results are not public and remain as internal documents, LHCb-ROADMAP1-002 LHCb-ROADMAP4-001.

<sup>16</sup> The purpose of that type of tests may vary. If two programs differ in physics assumptions, it may help to control physics precision. If physics assumptions are identical, but technical construction differ, then comparison checks correctness of the algorithm implementation. Finally, comparison of results from the same program, but installed at different computers may

<sup>14</sup> E.g.,  $\tau$  lepton decay products including neutrinos.

which are actually based on second-order matrix element and exponentiation was found to be excellent [493, 494].

### 5.5 Spin effects

The lifetime of the  $\tau$  lepton is orders of magnitude larger not only of its formation time in high energy experiments, but also of the time of all phenomena related to higher-order corrections such as bremsstrahlung.

Separation of  $\tau$  production and decay is excellent due to the small width of the  $\tau$  lepton. Its propagator can be well approximated by a delta function for phase space and matrix elements. The cross section for the process  $f\bar{f} \rightarrow \tau^+\tau^-Y; \tau^+ \rightarrow X^+\bar{\nu}_\tau; \tau^- \rightarrow l^-\nu_l\nu_\tau$  reads:

$$d\sigma = \sum_{spin} |\mathcal{M}|^2 d\Omega = \sum_{spin} |\mathcal{M}|^2 d\Omega_{prod} d\Omega_{\tau^+} d\Omega_{\tau^-}$$

, where  $Y$  and  $X^+$  stand for particles produced together with  $\tau^+\tau^-$  and in  $\tau^+$  decay, respectively;  $d\Omega$ ,  $d\Omega_{prod}$ ,  $d\Omega_{\tau^+}$ ,  $d\Omega_{\tau^-}$  denote the phase space in the original process, in production and decay, respectively.

This formalism looks simple, but because of over 20  $\tau$  decay channels there are more than 400 distinct processes.

Let us write the spin amplitude separated into parts for  $\tau$  pair production and decay:

$$\mathcal{M} = \sum_{\lambda_1\lambda_2=1}^2 \mathcal{M}_{\lambda_1\lambda_2}^{prod} \mathcal{M}_{\lambda_1}^{\tau^+} \mathcal{M}_{\lambda_2}^{\tau^-}.$$

After integrating out the  $\tau$  propagators, the formula for the cross section can be rewritten as

$$d\sigma = \left( \sum_{spin} |\mathcal{M}^{prod}|^2 \right) \left( \sum_{spin} |\mathcal{M}^{\tau^+}|^2 \right) \left( \sum_{spin} |\mathcal{M}^{\tau^-}|^2 \right) wt d\Omega_{prod} d\Omega_{\tau^+} d\Omega_{\tau^-},$$

where

$$wt = \left( \sum_{i,j=0,3} R_{ij} h_+^i h_-^j \right),$$

$$R_{00} = 1, \quad < wt > = 1, \quad 0 \leq wt \leq 4.$$

$R_{ij}$  can be calculated from  $\mathcal{M}_{\lambda_1\lambda_2}$  while  $h_+^i$  and  $h_-^j$  from  $\mathcal{M}^{\tau^+}$  and  $\mathcal{M}^{\tau^-}$ , respectively. Bell inequalities (related to Einstein-Rosen-Podolsky paradox [495]) tell us that in general it is impossible to rewrite  $wt$  in the following factorized form,  $wt^{factorized}$ :

$$wt \neq wt^{factorized} = \left( \sum_{i,j=0,3} R_i^A h_+^i \right) \left( \sum_{i,j=0,3} R_j^B h_-^j \right),$$

where  $R_i^A$  and  $R_j^B$  are four-component objects calculated from variables of the process of  $\tau$  pair production. In the

check correctness of the code implementation in a new software environment. Such comparisons or just the data necessary for comparisons will be referred to as respectively physical, technical and installation benchmarks. They are indispensable for reliable use of Monte Carlo programs.

Monte Carlo construction it is thus impossible to generate a  $\tau^+$  and  $\tau^-$  pair, each of the two in some quantum state, and later independently perform decays of  $\tau^+$  and  $\tau^-$ . This holds at all orders of perturbation expansion. The  $\tau$  production and decay are correlated through the spin effects only which can be represented by the well behaving factor  $wt$ , previously introduced. Above formulae do not lead to any loss of precision and hold in presence of radiative corrections as well. Different options of formalisms based on those expressions are used in Monte Carlo programs and basically well founded. That should be confronted with processes where instead of  $\tau$  leptons short-lived intermediate states are considered. Then, in general, ambiguities appear and corrections proportional to the ratio of the resonance width to its mass (or other energy scales of the process resulting, for example, from the cut-offs) must be included. Interfering background diagrams may cause additional problems. For details we refer to [496, 467, 466].

### 5.6 $\tau$ lepton production

KORALB was published [465, 497] more than *twenty years ago*. It included first-order QED corrections and complete mass and spin effects. It turned out to be very useful, and still remains in broad use. On the other hand, some of its ingredients are outdated and do not match the present day requirements, even for technical tests. For example the function PIRET(S), which describes the real part of the photon hadronic vacuum polarisation as measured by the data collected until early 80's should be replaced by one of the new precise codes (see Section 6 for details).

Unfortunately, this replacement does not solve all normalisation problems of KORALB. For example, it is well known that the one-loop corrections are not sufficient. The two major improvements which were developed during the LEP era are the introduction of higher-order QED corrections into Monte Carlo simulation and a better way to combine loop corrections with the rest of the field theory calculations. For energies up to 10 GeV (typical of the B-factories), the KKMC Monte Carlo [466] provides a realization of the above improvements. This program includes higher-order QED matrix elements with the help of exclusive exponentiation, and explicit matrix elements up to the second order. Also in this case the function calculating the vacuum polarisation must be replaced by a version appropriate for low energy (see Section 6).

Once this is completed, and if the two-loop photon vacuum polarisation can be neglected, KORALB and KKMC can form a base for tests and studies of systematic errors for cross section normalisations at low energies. Using a strategy similar to the one for Bhabha scattering [498], the results obtained in [499, 273] allow to expect a precision of 0.35-0.45% using KKMC at Belle/BaBar energies. Certainly, a precision tag similar to that for linear colliders can be also achieved for lower energies. Work beyond [499] and explained in that paper would be then necessary.



## 5.7 Separation into leptonic current and hadronic one

The matrix element used in TAUOLA for semileptonic decay  $\tau(P, s) \rightarrow \nu_\tau(N)X$ ,

$$\mathcal{M} = \frac{G}{\sqrt{2}} \bar{u}(N) \gamma^\mu (v + a\gamma_5) u(P) J_\mu \quad (204)$$

requires the hadronic current  $J_\mu$  to be known. The expression is easy to manipulate. One obtains:

$$\begin{aligned} |\mathcal{M}|^2 &= G^2 \frac{v^2 + a^2}{2} (\omega + H_\mu s^\mu), \\ \omega &= P^\mu (\Pi_\mu - \gamma_{va} \Pi_\mu^5), \quad , \\ H_\mu &= \frac{1}{M} (M^2 \delta_\mu^\nu - P_\mu P^\nu) (\Pi_\nu^5 - \gamma_{va} \Pi_\nu), \\ \Pi_\mu &= 2[(J^* \cdot N) J_\mu + (J \cdot N) J_\mu^* - (J^* \cdot J) N_\mu], \\ \Pi^{5\mu} &= 2 \operatorname{Im} \epsilon^{\mu\nu\rho\sigma} J_\nu^* J_\rho N_\sigma, \\ \gamma_{va} &= -\frac{2va}{v^2 + a^2}. \end{aligned} \quad (205)$$

If  $\tau$  coupling  $v + a\gamma_5$  and  $m_{\nu_\tau} \neq 0$  is allowed, one has to add to  $\omega$  and  $H_\mu$ :

$$\begin{aligned} \hat{\omega} &= 2 \frac{v^2 - a^2}{v^2 + a^2} m_\nu M (J^* \cdot J), \\ \hat{H}^\mu &= -2 \frac{v^2 - a^2}{v^2 + a^2} m_\nu \operatorname{Im} \epsilon^{\mu\nu\rho\sigma} J_\nu^* J_\rho P_\sigma. \end{aligned} \quad (206)$$

The expressions are useful for Monte Carlo applications and are also calculable from the first principles. The resulting expression can be used to the precision level of the order of 0.2-0.3%.

In contrast to other parts, the hadronic current  $J_\mu$  still can not be calculated reliably from the first principles. Some theoretical constraints need to be fulfilled, but in general it has to be obtained from the experimental data. We will return to this point later (see Section 5.9).

## 5.8 Bremsstrahlung in decays

The PHOTOS Monte Carlo is widely used for generation of radiative corrections in cascade decays, starting from the early papers [500, 501]. With time the precision of its prediction improved significantly, but the main principle remains the same. Its algorithm is aimed to modify the content of the event record filled in with complete cascade decays at earlier steps of the generation. PHOTOS modifies the content of the event record; it adds additional photons to the decay vertices and at the same time modifies a kinematic configuration of other decay products.

One could naively expect that this strategy is bound to substantial approximations. However, the algorithm is compatible with NLO calculations, leads to a complete coverage of the phase space for multi-photon final states and provides correct distributions in soft photon limits. For more details of the program organisation and its phase space generation we address the reader to [492].

The changes introduced over the last few years into the PHOTOS Monte Carlo program itself were rather small and the work concentrated on its theoretical foundations. This wide and complex subject goes beyond the scope of this Review and the interested reader can consult [502], where some of the topics are discussed. Previous tests of two-body decays of the  $Z$  into a pair of charged leptons [494] and a pseudoscalar  $B$  into a pair of scalars [492] were recently supplemented [503] with the study of  $W^\pm \rightarrow l^\pm \nu \gamma$ . The study of the process  $\gamma^* \rightarrow \pi^+ \pi^-$  is on-going [504]. In all of these cases a universal kernel of PHOTOS was replaced with the one matching the exact first-order matrix element. In this way terms for the NLO/NLL level are implemented. The algorithm covers the full multi-photon phase space and it is exact in the infrared region of the phase space. One should not forget that PHOTOS generates weight-one events.

The results of all tests of PHOTOS with an NLO kernel are at a sub-per mill level. No differences with benchmarks were found, even for samples of  $10^9$  events. When simpler physics assumptions were used, differences between total rates at sub-per mill level were observed or they were matching a precision of the programs used for tests.

This is very encouraging and points to the possible extension of the approach beyond the QED (scalar QED). In particular, to the QCD and/or models with phenomenological Lagrangians for interactions of photons with hadrons. For this work to be completed, spin amplitudes have to be further studied [505].

The refinements discussed above affect the practical side of simulations for  $\tau$  physics only indirectly. Changes in the kernels necessary for NLO may remain as options for tests only. They are available from the PHOTOS web page [503], but are not recommended for wider use. The corrections are small, distributions visualizing their size are available. On the other hand, their use could be perilous, it requires control of the decaying particle spin state. That is known, see, e.g., [506], to be not easy because of technical reasons.

We will show later that radiative corrections do not provide a limitation in the quest for improved precision of matching theoretical models to experimental data until issues discussed in subsection 5.12 are solved.

## 5.9 Hadronic currents

So far all discussed contributions to the predictions were found to be controlled to the precision level of 0.5% with respect to decay rate under study<sup>17</sup>

This is not the case for the hadronic current. It is the main source of our difficulties. It can not be obtained from QCD as involved energy scales are too small. On the other hand, for the low energy limits the scale is too large. That

<sup>17</sup> This 0.5% uncertainty is for the QED radiative effects. One should bear in mind other mechanisms involving production of photons, like, for example, the decay channel  $\omega \rightarrow \pi \gamma$ , which occurs with a probability of  $(8.28 \pm 0.28)\%$  and does not belong to a category of radiative corrections.

is the source of difficulties, but on the other hand, one can obtain theoretically a clear object if enough effort is devoted. That may lead to better understanding of boundaries of the perturbative domain of QCD as well.

The unquestionable property which hadronic current must fulfil is Lorentz invariance. For example, if the final state consists of three scalars, respectively of momenta  $p_1$ ,  $p_2$ ,  $p_3$ , it must take the form

$$J^\mu = N \{ T_\nu^\mu [c_1(p_2 - p_3)^\nu F_1 + c_2(p_3 - p_1)^\nu F_2 + c_3(p_1 - p_2)^\nu F_3] + c_4 q^\mu F_4 - \frac{ic_5 \epsilon^\mu_{\nu\rho\sigma}}{4\pi^2 f_\pi^2} p_1^\nu p_2^\rho p_3^\sigma F_5 \},$$

$T_{\mu\nu} = g_{\mu\nu} - q_\mu q_\nu / q^2$  is the transverse projector,  $q = p_1 + p_2 + p_3$ . Functions  $F_i$  depend on three variables that can be chosen as  $q^2 = (p_1 + p_2 + p_3)^2$  and two of the following ones  $s_1 = (p_2 + p_3)^2$ ,  $s_2 = (p_1 + p_3)^2$ ,  $s_3 = (p_2 + p_3)^2$ . This form is obtained from Lorentz invariance only.

Among the first four hadronic structure functions ( $F_1$ ,  $F_2$ ,  $F_3$ ,  $F_4$ ), only three are independent. We leave the  $F_4$  structure function in the basis because, neglecting the pseudoscalar resonance mechanism production, the contribution due to  $F_4$  is negligible ( $\sim m_\pi^2/q^2$ ) [507] and drops out (depending on the decay channel) one of  $F_1$ ,  $F_2$  or  $F_3$ , exactly as it is in TAUOLA since long.

In each case, the number of independent functions is four (rather than five) and it is not larger than the dimension of our space-time. That is why the projection operators can be defined, for two- and three-scalar final states. Work in that direction has already been done in Ref. [471] and then implemented in tests of TAUOLA too. Thanks to such a method, hadronic currents can be obtained from data without any need of phenomenological assumptions. Since long such methods were useful for data analysis, but only in part. Experimental samples were simply too small.

At present, for high statistics and precision the method may be revisited. That is why, it is of great interest to verify whether detector deficiencies will not invalidate the method or if adjustments due to incomplete phase space coverage are necessary. We will return to that question later. In the mean time let us return to other theoretical considerations which constrain the form of hadronic currents, but not always to the precision of today data.

## 5.10 The resonance chiral approximation and its result for the currents

Once the allowed Lorentz structures are determined and a proper minimal set of them is chosen, one should impose the QCD symmetries valid at low energies. The chiral symmetry of the massless QCD allows to develop an effective field theory description valid for momenta much smaller than the  $\rho$  mass,  $\chi PT$  [508, 509].

Although  $\chi PT$  cannot provide predictions valid over all  $\tau$  decay phase space, it constrains the form and the normalisation of the form factors in such limits.

The model, proposed in [441] for  $\tau$  decaying to pions, and used also for extensions to other decay channels, employs weighted products of Breit-Wigner functions to take into account resonance exchange. The form factors used

there have the right chiral limit at LO. However, as it was demonstrated in [507], they do not reproduce the NLO chiral limit.

The step towards incorporating the right low-energy limit up to NLO and the contributions from meson resonances which reflect the experimental data was done within Resonance Chiral Theory ( $R\chi T$  [433, 432]). The current state-of-the-art for the hadronic form factors ( $F_i$ ) appearing in the  $\tau$  decays is described in [510]. Apart from the correct low energy properties, it includes a right falloff [511, 512] at high energies.

The energy-dependent imaginary parts in the propagators of the vector and the axial-vector mesons  $1/(m^2 - q^2 - im\Gamma(q^2))$  were calculated in [513] at one-loop, exploiting the optical theorem that relates the appropriate hadronic matrix elements of  $\tau$  decays and the cuts with on-shell mesons in the (axial-) vector-(axial-) vector correlators.

This formalism has been shown to describe successfully the invariant mass spectra of experimental data in  $\tau$  decays for the following hadronic systems:  $\pi\pi$  [514, 515, 516],  $\pi K$  [517, 518],  $3\pi$  [507, 510, 519] and  $KK\pi$  [510, 519]. Other channels will be worked out along the same lines.

It has already been checked that the  $R\chi T$  results provide also a good description of the three-meson processes:  $\Gamma(\tau \rightarrow 3\pi\nu_\tau)$  [520] and  $\sigma(e^+e^- \rightarrow KK\pi)_{I=1}$  [396].

Both the spin-one resonance widths and the form factors of the decays  $\tau^- \rightarrow (\pi\pi, \pi K, 3\pi, KK\pi)^- \nu_\tau$  computed within  $R\chi T$  are only being implemented in TAUOLA now.

Starting from a certain precision level, the predictions, like the presented above, may turn out to be not sufficiently precise. Nonetheless, even in such a case they can provide some essential constraints on the form of the functions  $F_i$ . Further refinements will require large and combined efforts of experimental and theoretical physicists. We will elaborate on possible technical solutions later in the review. Such attempts turned out to be difficult in the past and a long time was needed for parameterisations given in [473] to become public. Even now they are semi-official and are not based on the final ALEPH and/or CLEO data.

## 5.11 Isospin symmetry of the hadronic currents

If one neglects quark masses, QCD is invariant under a transformation replacing quark flavours. As a consequence, hadronic currents describing vector  $\tau$  decays ( $2\pi, 4\pi, \eta\pi\pi, \dots$ ) and low energy  $e^+e^-$  annihilation into corresponding isovector final states are related and can be obtained one from another [521, 522]. This property, often referred to as conservation of vector current (CVC) in  $\tau$  decays, results in a possibility to predict invariant mass distributions of the hadronic system as well as the corresponding branching fractions in the  $\tau$  decays using  $e^+e^-$  data. A systematic check of these predictions showed that at the (5-10)% level they work rather well [523].

In principle, the corrections due to mass and charge difference between  $u$  and  $d$  quarks are not expected to provide significant and impossible to control effects [524, 525]. However, the high-precision data of the CLEO [526],

ALEPH [527], OPAL [528], Belle [474], CMD-2 [283, 382, 384, 386], SND [282] and KLOE [368] collaborations in the  $2\pi$  channel challenged this statement and as it was shown in [27, 11, 3, 529, 28, 374, 19], the spectral functions for  $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$  significantly differ from those obtained using  $e^+e^- \rightarrow \pi^+\pi^-$  data. Some evidence for similar discrepancy is also observed in the  $\tau^- \rightarrow 2\pi^- \pi^+ \pi^0 \nu_\tau$  decay [530, 531, 333]. This effect remains unexplained. The magnitude of the isospin-breaking corrections has been recently updated making the discrepancy in the  $2\pi$  channel smaller [29].

These CVC based relations were originally used in the TAUOLA form factors parameterisation, but they were often modified to improve fits to the data. Let us point here to an example where experimental  $e^+e^-$  data were used for the model of the  $\tau \rightarrow 4\pi \nu_\tau$  decay channels [472]. In this case, only a measurement of the distribution in the total invariant mass of the hadronic system was available. That is not enough to fix the distribution over multidimensional phase space. For other dimensions one had to rely on phenomenological models or other experiments. In the future, this may not be necessary, but will always remain as a method of benchmarks construction.

## 5.12 The challenges

As we have argued before, refined techniques for fits, involving simultaneous fits to many  $\tau$  decay channels, are necessary to improve phenomenological description of  $\tau$  decays. Complex backgrounds (where each channel contributes to signatures of other decay modes as well), different sensitivities of experiments for measurements of some angular distributions within the same hadronic system, and sometimes even an incomplete reconstruction of final states, are the main cause of this necessity. Moreover, theoretical models based on the Lagrangian approach describe simultaneously more than one  $\tau$  decay channel with the same set of parameters and only simultaneous fits allow to establish their experimental constraints in a consistent way. Significant efforts are thus necessary and close collaboration between phenomenologists and experimental physicists is indispensable. As a result, techniques of automated calculations of hadronic currents may become necessary [532].

## 5.13 Technical solutions for fits

For the final states of up to three scalars use of projection operators [471] is popular since long [530]. It enables, in principle, to obtain form factors used in hadronic currents directly from the data, at least in principle, and for one scalar function defined in 207 at a time. Only recently experimental samples became sufficiently large. However, to exploit the method one may have to improve it first: systematically include effects of limited detector acceptance. Implementation of the projection operators into packages

like MC-TESTER [533] may be useful. Efforts in that direction are being pursued now<sup>18</sup> [535].

On theoretical side one may need to choose predictions from many models before sufficiently good agreement with data will be achieved. Some automated methods of calculations may be then useful [536]. This is especially important, for hadronic multiplicities larger than three, when projector operators have never been defined.

Certain automation of the methods is thus advisable. To discriminate from the broad spectrum of choices new methods of data analysis may become useful [537]. Such methods may require simulating samples of events where several options for the matrix element calculation are used simultaneously<sup>19</sup>.

The neutrino coming from  $\tau$  decay escapes detection and as a result the  $\tau$  rest frame can not be reconstructed. Nevertheless, as it was shown in Ref. [471], angular distributions can be used for construction of projection operators, which allows extraction of the hadronic structure functions from the data. This is possible as they depend on  $s_1$ ,  $s_2$  and  $q^2$  only.

A dedicated module for the MC-TESTER [533], implementing the moments of different angular functions defined in Eq. (39)-(47) of Ref. [471], is under development. The moments are proportional to combinations of type of  $\alpha|F_i|^2 + \beta|F_j|^2 + \gamma\text{Re}(F_i F_j^*)$ , where the coefficients  $\alpha$ ,  $\beta$ , and  $\gamma$  are functions of hadron 4-momenta components in the hadronic rest frame. Preliminary results obtained with large statistics of 5 million  $\tau \rightarrow a_1 \nu_\tau \rightarrow 3\pi \nu_\tau$  decays, and assuming vanishing  $F_3$ , and  $F_5$  form factors, show that it is possible to extract the  $|F_1|^2$ ,  $|F_2|^2$ , and  $|F_1 \cdot F_2^*|^2$  as functions of  $s_1$ ,  $s_2$  and  $Q^2$ . This extraction requires solving a set of equations. Since the solution is sensitive to the precision of the estimation of the moments entering the equation, large data samples of the order of  $\mathcal{O}(10^6 - 10^7)$  are necessary. The calculation of the moments requires also knowledge of the initial  $\sqrt{s}$  of the  $\tau$  pair, which makes the analysis sensitive to initial state radiation (ISR) effects. The same studies show that the analysis is easier if one, instead of extracting the  $|F_i|^2$  form factors, compares the moments obtained from the experimental data with theoretical predictions. Such comparison does not require repetition of the Monte Carlo simulation of  $\tau$  decays with different form factors, and only the calculation of combinations of  $|F_i|^2$ , and  $\text{Re}(F_i F_j^*)$  is necessary. This is much simpler than comparing the kinematic distributions obtained from data with distributions coming from Monte Carlo simulations with various theoretical models. Further complications, for example, due to the presence of an initial state bremsstrahlung or an incomplete acceptance of decay phase space were not yet taken into account.

<sup>18</sup> This may help to embed the method in the modern software for fits, see, e.g., [534].

<sup>19</sup> Attempts to code such methods into TAUOLA combined with programs for  $\tau$  pair production and experimental detector environment were recently performed [538], but they were applied so far as prototypes only, see Fig. 1 of Ref. [539].

## 5.14 Perspectives

Definitely the improvements of  $\tau$  decay simulation packages and fit strategies are of interest for phenomenology of low energy. As a consequence, their input for such domains like phenomenology of  $g-2$  or  $\alpha_{QED}$ ,  $\alpha_{QCD}$  and their use in constraints of new physics would improve.

In this section let us argue if possible benefits for LHC phenomenology may arise from better understanding of  $\tau$  decay channels in measurements as well. In papers [540, 541] it was shown that spin effects can indeed be useful to measure such properties of the Higgs boson as parity. Moreover, such methods were verified to work well when detector effects as proposed for a future linear collider were taken into account. Good control of the decay properties is helpful. For example, in Ref. [542] it was shown for the  $\tau \rightarrow a_1 \nu_\tau \rightarrow 3\pi \nu_\tau$  decay that the sensitivity to the  $\tau$  polarisation increases about four times when all angular variables are used compared with usual  $d\Gamma/dq^2$ , see also [543].

Even though  $\tau$  decays provide some of the most prominent signatures for the LHC physics program, see, e.g., Ref. [544], for some time it was expected that methods exploiting detailed properties of  $\tau$  cascade decays are not practical for LHC studies. Thanks to efforts on reconstruction of  $\pi^0$  and  $\rho$  invariant mass peaks, this opinion evolves. Such work was done for studies of CMS ECAL detector inter-calibration [545], and in a relatively narrow  $p_T$  range (5-10 GeV), some potentially encouraging results were obtained. Some work in context of searches for new particles recently started [546]. There, improved knowledge of distinct  $\tau$  decay modes may become important at a certain point.

One can conclude that the situation is similar to that of LEP beginning, some control of all  $\tau$  decay channels is important. Nonetheless, only if detector studies of  $\pi^0$  and  $\rho$  reconstruction will provide positive results, the gate to improve a sensitivity of  $\tau$  spin measurements with most of its decay modes, as at LEP [547, 548, 549] will be open. At this moment, however, it is difficult to judge about importance of such improvements of  $\tau$  decay description for LHC perspectives. Experience of the first years of LHC must be consolidated first. In any case such activity is important for physics of the future Linear Colliders.

## 5.15 Summary

We have shown that the most urgent challenge in the quest for better understanding of  $\tau$  decays is the development of efficient techniques for fitting multidimensional distributions, which take into account realistic detector conditions. This includes cross contamination of different  $\tau$  decay modes, their respective signatures and detector acceptance effects, which have to be simultaneously taken into account when fitting experimental data. Moreover, at the current experimental precision, theoretical concepts have to be reexamined. In contrast to the past, the precision of predictions based on chiral Lagrangians and/or isospin symmetry can not be expected to match always

the precision of the data. The use of model-independent data analyses should be encouraged whenever possible in realistic conditions.


Good understanding of  $\tau$  decays is crucial for understanding the low energy regime of strong interactions and matching between the non-perturbative and the perturbative domains. Further work on better simulations of  $\tau$  decays at LHC is needed to improve its potential to study processes of new physics, especially in the Higgs sector. In addition, an accurate simulation of  $\tau$  decays is important for control of backgrounds for very rare decays. For the project to be successful, it should lead to encapsulation of our knowledge on  $\tau$  decays in a form of a Monte Carlo library to be used by low energy as well as high energy applications.

## 6 Vacuum polarization

### 6.1 Introduction

The vacuum polarisation (VP) of the photon is a quantum effect which leads, through renormalisation, to the scale dependence (‘running’) of the electromagnetic coupling,  $\alpha(q^2)$ . It therefore plays an important role in many physical processes and its knowledge is crucial for many precision analyses. A prominent example is the precision fits of the Standard Model as performed by the electroweak working group, where the QED coupling  $\alpha(q^2 = M_Z^2)$  is the least well known of the set of fundamental parameters at the  $Z$  scale,  $\{G_\mu, M_Z, \alpha(M_Z^2)\}$ . Here we are more concerned about the VP at lower scales as it enters all photon-mediated hadronic cross sections. These are used, e.g., in the determination of the strong coupling  $\alpha_s$ , the charm and bottom quark masses from  $R_{\text{had}}$  as well as in the evaluation of the hadronic contributions to the muon  $g-2$  and  $\alpha(q^2)$  itself. It also appears in Bhabha scattering in higher orders of perturbation theory needed for a precise determination of the luminosity. It is hence clear that VP also has to be included in the corresponding Monte Carlo programs.

In the following we shall first define the relevant notations, then briefly discuss the calculation of the leptonic and hadronic VP contributions, before comparing available VP parametrisations.



**Fig. 82.** Photon vacuum polarisation  $\Pi(q^2)$ .

Conventionally the vacuum polarisation function is denoted by  $\Pi(q^2)$  where  $q$  is a space- or time-like momentum. The shaded blob in Fig. 82 stands for all possible one-particle irreducible leptonic or hadronic contributions. The full photon propagator is then the sum of the bare



photon propagator and arbitrarily many iterations of VP insertions,

$$\text{full photon propagator} \sim \frac{-i}{q^2} \cdot (1 + \Pi + \Pi \cdot \Pi + \Pi \cdot \Pi \cdot \Pi + \dots) \quad (207)$$

The Dyson summation of the real part of the one-particle irreducible blobs then defines the effective QED coupling

$$\alpha(q^2) = \frac{\alpha}{1 - \Delta\alpha(q^2)} = \frac{\alpha}{1 - \text{Re}\Pi(q^2)}, \quad (208)$$

where  $\alpha \equiv \alpha(0)$  is the usual fine structure constant,  $\alpha \sim 1/137$ . It is determined most precisely through the anomalous magnetic moment of the electron,  $a_e$ , as measured by the Harvard group to an amazing 0.28 ppt [1], in agreement with less precise determinations from caesium and rubidium atom experiments. The most precise value for  $\alpha$ , which includes the updated calculations of  $\mathcal{O}(\alpha^4)$  contributions to  $a_e$  [550], is given by  $1/\alpha = 137.035999084(51)$ .

By using Eq. (208) we have defined  $\Pi$  to include the electric charge squared,  $e^2$  for leptons, but note that different conventions are used in the literature, and sometimes  $\Pi$  is also defined with a different overall sign.

Equation (208) is the usual definition of the running effective QED coupling and has the advantage that one obtains a real coupling. However, the imaginary part of the VP function  $\Pi$  is completely neglected, which is normally a good approximation as the contributions from the imaginary part are formally suppressed. This can be seen, e.g., in the case of the ‘undressing’ of the experimentally measured hadronic cross section  $\sigma_{\text{had}}(s)$ . The measured cross section  $e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons}$  contains  $|\text{full photon propagator}|^2$ , i.e. the modulus squared of the infinite sum (207). Writing  $\Pi = e^2(P + iA)$  one easily sees that

$$|1 + e^2(P + iA) + e^4(P + iA)^2 + \dots|^2 = 1 + e^2 2P + e^4(3P^2 - A^2) + e^6 4P(P^2 - A^2) + \dots$$

and that the imaginary part  $A$  enters only at order  $\mathcal{O}(e^4)$  compared to  $\mathcal{O}(e^2)$  for the leading contribution from the real part  $P$ . To account for the imaginary part of  $\Pi$  one may therefore apply the summed form of the ‘(un)dressing’ factor with the relation

$$\sigma_{\text{had}}(s) = \frac{\sigma_{\text{had}}^0(s)}{|1 - \Pi|^2} \quad (209)$$

instead of the traditionally used relation with the real effective coupling,

$$\sigma_{\text{had}}(s) = \sigma_{\text{had}}^0(s) \left( \frac{\alpha(s)}{\alpha} \right)^2. \quad (210)$$

We shall return to a comparison of the different approaches below for the case of the hadronic VP.

It should be noted that the summation breaks down and hence can not be used if  $|\Pi(s)| \sim 1$ . This is the case if  $\sqrt{s}$  is very close to or even at narrow resonance energies.

In this case one can not include the narrow resonance in the definition of the effective coupling but has to rely on another formulation, e.g. through a Breit-Wigner propagator (or a narrow width approximation with a delta-function). For a discussion of this issue see [551]. Also note that the VP summation covers only the class of one-particle irreducible diagrams of factorisable bubbles depicted in Fig. 82. This includes photon radiation within and between single bubbles, but clearly does not take into account higher-order corrections from initial state radiation or initial-final state interference effects in  $e^+e^- \rightarrow \text{hadrons}$ .

As will be discussed in the following, leptonic and hadronic contributions to  $\Delta\alpha$  are normally calculated separately and then added,  $\Delta\alpha(q^2) = \Delta\alpha_{\text{lep}}(q^2) + \Delta\alpha_{\text{had}}(q^2)$ . While the leptonic contributions can be predicted within perturbation theory, the precise determination of the hadronic contributions relies on a dispersion relation using experimental data as input.

## 6.2 Leptonic contributions

The leptonic contributions  $\Delta\alpha_{\text{lep}}$  have been calculated to sufficiently high precision. The leading order (LO) and next-to-leading order (NLO) contributions are known as analytic expressions including the full mass dependence [552], where LO and NLO refer to the expansion in terms of  $\alpha$ . The next-to-next-to-leading order (NNLO) contribution is available as an expansion in terms of  $m_\ell^2/q^2$  [553], where  $m_\ell$  is the lepton mass. To evaluate  $\Delta\alpha_{\text{lep}}(q^2)$  for  $|q^2| \lesssim m_\tau^2$ , this expansion is not appropriate, but this is exactly the region where the hadronic uncertainties are dominant. Also from the smallness of the NNLO contribution, we conclude that we do not need to further improve the leptonic contributions beyond this approximation.

The evaluation of the LO contribution is rather simple, and we briefly summarise the results below. Hereafter, it is understood that we impose the renormalisation condition  $\Pi(0) = 0$  on  $\Pi(q^2)$ . For  $q^2 < 0$ , the VP function reads

$$\Pi(q^2) = -\frac{e^2}{36\pi^2} \left( 5 - 12\eta + 3(-1 + 2\eta)\sqrt{1 + 4\eta} \ln \frac{\sqrt{1 + 4\eta} + 1}{\sqrt{1 + 4\eta} - 1} \right), \quad (211)$$

where  $\eta \equiv m_\ell^2/(-q^2)$ . For  $0 \leq q^2 \leq 4m_\ell^2$  one obtains

$$\Pi(q^2) = -\frac{e^2}{36\pi^2} \left( 5 - 12\eta + 3(-1 + 2\eta)\sqrt{-1 - 4\eta} \arctan \frac{\sqrt{-1 - 4\eta}}{-1 - 2\eta} \right), \quad (212)$$

and for  $q^2 \geq 4m_\ell^2$

$$\Pi(q^2) = -\frac{e^2}{36\pi^2} \left( 5 - 12\eta + 3(-1 + 2\eta)\sqrt{1 + 4\eta} \cdot \ln \frac{1 + \sqrt{1 + 4\eta}}{1 - \sqrt{1 + 4\eta}} \right) - \frac{i e^2}{12\pi} (1 - 2\eta)\sqrt{1 + 4\eta}. \quad (213)$$



An easily accessible reference which gives the NLO contributions is, for instance, Ref. [554, 555]. As mentioned above, the NNLO contribution is given in Ref. [553]. For all foreseeable applications the available formulae can be easily implemented and provide a sufficient accuracy. While the uncertainty from  $\alpha$  is of course completely negligible, the uncertainty stemming from the lepton masses is only tiny. Therefore the leptonic VP poses no problem.

### 6.3 Hadronic contributions

In contrast to the leptonic case, the hadronic VP  $\Pi_{\text{had}}(q^2)$  can not be reliably calculated using perturbation theory. This is clear for time-like momentum transfer  $q^2 > 0$ , where, via the optical theorem  $\text{Im } \Pi_{\text{had}}(q^2) \sim \sigma(e^+e^- \rightarrow \text{hadrons})$  goes through all the resonances in the low energy region. However, it is possible to use a dispersion relation to obtain the real part of  $\Pi$  from the imaginary part. The dispersion integral is given by

$$\Delta\alpha_{\text{had}}^{(5)}(q^2) = -\frac{q^2}{4\pi^2\alpha} \text{P} \int_{m_\pi^2}^{\infty} \frac{\sigma_{\text{had}}^0(s) ds}{s - q^2}, \quad (214)$$

where  $\sigma_{\text{had}}^0(s)$  is the (undressed) hadronic cross section which is determined from experimental data. Only away from hadronic resonances and (heavy) quark thresholds one can apply perturbative QCD to calculate  $\sigma_{\text{had}}^0(s)$ . In this region the parametric uncertainties due to the values of the quark masses and  $\alpha_s$ , and due to the choice of the renormalisation scale, are small. Therefore the uncertainty of the hadronic VP is dominated by the statistical and systematic uncertainties of the experimental data for  $\sigma_{\text{had}}^0(s)$  used as input in (214).

Note that the dispersion integral (214) leads to a smooth function for space-like momenta  $q^2 < 0$ , whereas in the time-like region it has to be evaluated using the principal value description and shows strong variations at resonance energies, as demonstrated e.g. in Fig. 83. In Eq. (214)  $\Delta\alpha_{\text{had}}^{(5)}$  denotes the five-flavour hadronic contribution. At energies we are interested in, i.e. far below the  $t\bar{t}$  threshold, the contribution from the top quark is small and usually added separately. The analytic expressions for  $\Delta\alpha^{\text{top}}(q^2)$  obtained in perturbative QCD are the same as for the leptonic contributions given above, up to multiplicative factors taking into account the top quark charge and the corresponding SU(3) colour factors, which read  $Q_t^2 N_c$  at LO and  $Q_t^2 \frac{N_c^2 - 1}{2N_c}$  at NLO.

Contributions from narrow resonances can easily be treated using the narrow width approximation or a Breit-Wigner form. For the latter one obtains

$$\Delta\alpha^{\text{Breit-Wigner}}(s) = \frac{3\Gamma_{ee}}{\alpha M} \frac{s(s - M^2 - \Gamma^2)}{(s - M^2)^2 + M^2\Gamma^2}, \quad (215)$$

with  $M$ ,  $\Gamma$  and  $\Gamma_{ee}$  the mass, total and electronic width of the resonance. For a discussion of the undressing of  $\Gamma_{ee}$  see [551].

Although the determination of  $\Delta\alpha_{\text{had}}^{(5)}(q^2)$  via the dispersion integral (214) may appear straightforward, in practise the data combination for  $\sigma_{\text{had}}^0(s)$  is far from trivial. In the low energy region up to about 1.4 – 2 GeV many data sets from the different hadronic exclusive final states (channels) from various experiments have to be combined, before the different channels which contribute incoherently to  $\sigma_{\text{had}}^0(s)$  can be summed. For higher energies the data for the fast growing number of possible multi-hadronic final states are far from complete, and instead inclusive (hadronic) measurements are used. For the details of the data input, the treatment of the data w.r.t. radiative corrections, the estimate of missing threshold contributions and unknown subleading channels (often via isospin correlations) and the combination procedures we refer to the publications of the different groups cited below.

In the following we shall briefly describe and then compare the evaluations of the (hadronic) VP available as parametrisations or tabulations from different groups.

### 6.4 Currently available VP parametrisations


For many years Helmut Burkhardt and Bolek Pietrzyk have been providing the Fortran function named REPI for the leptonic and hadronic VP [167, 556, 252, 557, 9]. While the leptonic VP is coded in analytical form with one-loop accuracy, the hadronic VP is given as a very compact parametrisation in the space-like region, but does not cover the time-like region. For their latest update see [10]. The code can be obtained from Burkhardt's web-pages which contain also a short introduction and a list of older references, see

<http://hbu.web.cern.ch/hbu/aqed/aqed.html>.

Similarly, Fred Jegerlehner has been providing a package of Fortran routines for the running of the effective QED coupling [251, 7, 12, 11, 4, 5]. It provides leptonic and hadronic VP both in the space- and time-like region. For the leptonic VP the complete one- and two-loop results and the known high energy approximation for the three-loop corrections are included. The hadronic contributions are given in tabulated form in the subroutine HADR5N. The full set of routines can be downloaded from Jegerlehner's web-page

<http://www-com.physik.hu-berlin.de/~fjeger/>. The version available from there is the one we use in the comparisons below and was last modified in November 2003. It will be referred to as J03 in the following. An update is in progress and other versions may be available from the author upon request. Note that for quite some time his routine has been the only available code for the time-like hadronic VP. Fig. 83 shows the leptonic and hadronic contributions together with their sum as given by Jegerlehner's routine.

The experiments CMD-2 and SND at Novosibirsk are using their own VP compilation to undress hadronic cross sections, and the values used are given in tables in some of their publications. Recently CMD-2 has made their compilation publicly available, see Fedor Ignatov's web-page




**Fig. 83.** Different contributions to  $\Delta\alpha(s)$  in the time-like region as given by the routine from Fred Jegerlehner. Figure provided with the package `alphaQED.uu` from his homepage.

<http://cmd.inp.nsk.su/~ignatov/vpl/>. There links are given to a corresponding talk at the ‘4th meeting of the Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies’ (Beijing 2008), to the thesis of Ignatov (in Russian) and to a file containing the tabulation, which can be used together with a downloadable package. The tabulation is given for the real and imaginary parts of the sum of leptonic and hadronic VP, for both space- and time-like momenta, and for the corresponding errors. Fig. 84, also displayed on their web-page, shows the results from CMD-2 for  $|1 + \Pi|^2$  both for the space- and time-like momenta in the range  $-(15 \text{ GeV})^2 < q^2 < (15 \text{ GeV})^2$  (upper panel) and for the important low energy region  $-(2 \text{ GeV})^2 < q^2 < (2 \text{ GeV})^2$ . The solid (black) lines are the sum of leptonic and hadronic contributions, while the dotted (red) lines are for the leptonic contributions only.

Another independent compilation of the hadronic VP is available from the group of Hagiwara et al. [551] (HMNT), at present upon request from the authors. They provide tabulations (with a simple interpolation routine in Fortran) of  $\Delta\alpha_{\text{had}}^{(5)}(q^2)$  both in the space- and time-like region, and also a compilation of  $R_{\text{had}}(s)$ . Currently available routines are based on the analysis [13, 14]. Two different versions are provided, one including the narrow resonances  $J/\psi, \psi'$  and the Upsilon family,  $\Upsilon(1S) - \Upsilon(3S)$ , in Breit-Wigner form, one excluding them. However, for applications of  $\Delta\alpha$  it should be remembered that close to narrow resonances the resummation of such large contributions in the effective coupling breaks down. In this context, note that the compilation from Novosibirsk contains these narrow resonances, whereas the routine from Jegerlehner does contain  $J/\psi$  and  $\psi'$ , but seems to exclude (or smear over) the Upsilon resonances. When called in the charm or bottom resonance region Jegerlehner’s routine gives a warning that the “results may not be reliable close to J/Psi and Upsilon resonances”.


In the following we shall compare the parametrisations from the different groups.



**Fig. 84.**  $|1 + \Pi|^2$  from CMD-2’s compilation for space- and time-like momenta (labelled  $\sqrt{s}$ ); solid (black) lines: leptonic plus hadronic contributions, dotted (red) lines: only leptonic contributions. Upper panel:  $-(15 \text{ GeV})^2 < q^2 < (15 \text{ GeV})^2$ . Lower panel:  $-(2 \text{ GeV})^2 < q^2 < (2 \text{ GeV})^2$ . Figures provided by Fedor Ignatov.

## 6.5 Comparison of the results from different groups

In Fig. 85, we compare the parametrisations from Burkhart and Pietrzyk (BP05), Jegerlehner (J03) and Hagiwara et al. (HMNT) in the space-like (upper) and time-like region (lower panel). For the space-like region the differences among the three parametrisations are roughly within one standard deviation in the whole energy range shown. However, for the time-like region, there is disagreement between HMNT and J03 at several energy regions, most notably at  $1 \text{ GeV} \lesssim \sqrt{s} \lesssim 1.6 \text{ GeV}$ , and at  $0.8 \text{ GeV} \lesssim \sqrt{s} \lesssim 0.95 \text{ GeV}$ . As for the discrepancy at  $1 \text{ GeV} \lesssim \sqrt{s} \lesssim 1.6 \text{ GeV}$ , checking the routine from Jegerlehner, one finds that a too sparsely spaced energy grid in this region seems to be the reason. The discrepancy at  $0.8 \text{ GeV} \lesssim \sqrt{s} \lesssim 0.95 \text{ GeV}$  is further scrutinised in Fig. 86, where in addition to the two parametrisations HMNT (solid (red) line) and J03 (dotted (blue) line), the result for  $\Delta\alpha_{\text{had}}^{(5)}(s)/\alpha$  obtained by integrating over the




**Fig. 85.** Comparison of the results from Hagiwara et al. (HMNT [551]) for  $\Delta\alpha_{\text{had}}^{(5)}(q^2)$  in units of  $\alpha$  with parametrisations from Burkhardt and Pietrzyk (BP05 [10]) and Jegerlehner (J03). Upper panel:  $\Delta\alpha_{\text{had}}^{(5)}(Q^2)/\alpha$  for space-like momentum transfer ( $Q^2 < 0$ ), where the three parametrisations are indistinguishable. The differences (normalised and multiplied by 100) are highlighted by the dashed and dotted curves; the wide light (blue) band is obtained by using the error band of HMNT in the normalised difference to J03, labelled ‘(J03-HMNT)/HMNT ( $\times 100$ )’. Lower panel:  $\Delta\alpha_{\text{had}}^{(5)}(s)/\alpha$  from J03 and HMNT (as labelled) for time-like momenta ( $q^2 = s$ ). For readability, only the error band of HMNT is displayed.

$R$ -data as compiled by the PDG [259]<sup>20</sup> is shown as the dashed (green) line. While the results from HMNT and the one based on the PDG  $R$ -data agree rather well, their disagreement with the J03 compilation in the region  $0.8 \text{ GeV} \lesssim \sqrt{s} \lesssim 0.95 \text{ GeV}$  is uncomfortably large compared to the error but may be due to a different data input of the J03 parametrisation.

In the following we shall compare the parametrisation from HMNT with the one from the CMD-2 collaboration which has become available very recently. Note that for undressing their experimentally measured hadronic cross sections, CMD-2 includes the imaginary part of the VP function  $\Pi(q^2)$  in addition to the real part. Before com-

<sup>20</sup> The actual compilation of the data is available in electronic form from [http://pdg.lbl.gov/2008/hadronic-xsections/hadronicrpp\\_page1001.dat](http://pdg.lbl.gov/2008/hadronic-xsections/hadronicrpp_page1001.dat).



**Fig. 86.** Comparison of the results from Hagiwara et al. (HMNT, solid (red) line) for  $\Delta\alpha_{\text{had}}^{(5)}(s)/\alpha$  with the parametrisation from Jegerlehner (J03, dotted (blue) line) in the time-like region in the range  $\sqrt{s} = 0.7 - 1 \text{ GeV}$ . The dashed (green) line shows the result if the data compilation from the PDG [259] is used.

ing to the comparison with CMD-2, let us discuss some generalities about  $\text{Im } \Pi(q^2)$ . If we are to include the imaginary part, then the VP correction factor  $\alpha(q^2)^2$  should be replaced as


$$\left( \frac{\alpha}{1 - \Delta\alpha(q^2)} \right)^2 = \left( \frac{\alpha}{1 - \text{Re } \Pi(q^2)} \right)^2 \rightarrow \quad (216)$$

$$\left| \frac{\alpha}{1 - \Pi(q^2)} \right|^2 = \frac{\alpha^2}{(1 - \text{Re } \Pi(q^2))^2 + (\text{Im } \Pi(q^2))^2}.$$

Note that, as mentioned already in the introduction, the contribution from the real part appears at  $\mathcal{O}(e^2)$  in the denominator, while that from the imaginary part starts only at  $\mathcal{O}(e^4)$ . Because of this suppression we expect the effects from the imaginary part to be small. Nevertheless we would like to stress two points. First, field-theoretically, it is more accurate to include the imaginary part which exists above threshold. Including only  $\text{Re } \Pi(q^2)$  in the VP correction is an approximation which may be sufficient in most cases. Second, it is expected that the contribution from the imaginary part is of the order of a few per mille of the total VP corrections. While this seems small, it can be non-negligible at the  $\rho$  meson region where the accuracy of the cross section measurements reaches the order of (or even less than) 1%. Similarly, in the region of the narrow  $\phi$  resonance, the contributions from the imaginary part become non-negligible and should be taken into account.

In Fig. 87 the VP correction factor, based on the compilation from HMNT, with and without  $\text{Im } \Pi(q^2)$  is compared to  $|1 - \Pi(s)|^2$  as used by the CMD-2 collaboration in their recent analysis of the hadronic cross section in the  $2\pi$  channel in the  $\rho$  central region [386].<sup>21</sup> In the upper


<sup>21</sup> We thank Gennadiy Fedotovitch for providing us with a table including the VP correction factors not included in [386].



**Fig. 87.** Upper panel: Correction factor  $|1 - \Pi(s)|^2$  as used for ‘undressing’ by the CMD-2 collaboration in [386] (dashed line) compared to the same quantity using the HMNT compilation for the  $e^+e^- \rightarrow \text{hadrons}$  data (solid line). Also shown is the correction factor  $(1 - \text{Re } \Pi)^2 = (\alpha/\alpha(s))^2$ , based on  $\alpha(s)$  in the time-like region from HMNT (dotted line). Lower panel: Differences of the quantities as indicated on the plot.

panel the VP correction factors are given, whereas in the lower panel the differences are shown. As expected, the differences between the three are visible, and are about a few per mille at most. The difference between the CMD-2 results and the one from HMNT including  $\text{Im } \Pi(q^2)$  (solid (red) curve in the lower panel of Fig. 87 shows a marked dip followed by a peak in the  $\rho - \omega$  interference region where the  $\pi^+\pi^-$  cross section falls sharply. This is most probably a direct consequence of the different data input used. However, in most applications such a difference will be partially cancelled when integrated over an energy region including the  $\rho$  peak.

In Figs. 88 and 89 we compare  $\Delta\alpha(s)$  in the time-like region as given by the parametrisation from CMD-2 with the one from HMNT, where for HMNT we have calculated the leptonic contributions (up to including the NNLO corrections) as described above. The two panels in Fig. 88 (upper panel:  $0 < \sqrt{s} < 2$  GeV, lower panel:




**Fig. 88.**  $\Delta\alpha(s)$  in the time-like region as given by the parametrisation from CMD-2 (solid (blue) band) compared to the same quantity from HMNT (dotted (red) line). Upper panel:  $0 < \sqrt{s} < 2$  GeV, lower panel:  $2 \text{ GeV} < \sqrt{s} < 10$  GeV.

$2 \text{ GeV} < \sqrt{s} < 10$  GeV) show  $\Delta\alpha(s)$  with the  $1\sigma$  error band from CMD-2 as a solid (blue) band, whereas for HMNT the mean value for  $\Delta\alpha(s)$  is given by the dotted (red) line, which can hardly be distinguished. To highlight the differences between the two parametrisations, Fig. 89 displays the normalised difference  $(\Delta\alpha^{\text{CMD-2}}(s) - \Delta\alpha^{\text{HMNT}}(s))/\Delta\alpha^{\text{HMNT}}(s)$  as a solid (black) line, and also shows the relative errors of CMD-2 and HMNT as dashed (blue) and red (dotted) lines, respectively. As visible in Fig. 89, the error as given by the CMD-2 parametrisation is somewhat smaller than the one from HMNT. Both parametrisations agree fairly well, and for most energies the differences between the parametrisations are about as large or smaller than the error bands. Close to narrow resonances the estimated uncertainties are large, but as discussed above, there the approximation of the effective coupling  $\alpha(s)$  breaks down and resonance contributions should be treated differently.

## 6.6 Summary

Vacuum polarisation of the photon plays an important role in many physical processes. It has to be taken into account, e.g., in Monte Carlo generators for hadronic cross





**Fig. 89.** Solid (black) lines: Normalised difference  $(\Delta\alpha^{\text{CMD-2}}(s) - \Delta\alpha^{\text{HMNT}}(s))/\Delta\alpha^{\text{HMNT}}(s)$  in the time-like region. The dashed (blue) and dotted (red) lines indicate the relative error for the CMD-2 and HMNT parametrisations. Upper panel:  $0 < \sqrt{s} < 2$  GeV, lower panel:  $2 \text{ GeV} < \sqrt{s} < 10$  GeV.

sections or Bhabha scattering. When low energy data are used in dispersion integrals to predict the hadronic contributions to  $g-2$  or  $\Delta\alpha(q^2)$ , undressed data have to be used, so VP has to be subtracted from measured cross sections. The different VP contributions have been discussed, and available VP compilations have been briefly described and compared. Until recently only one parametrisation has been available in the time-like region, now three routines in the space- and time-like regions exist, from Jegerlehner, CMD-2 and HMNT, and a fourth from Burkhardt and Pietrzyk in the space-like region. While the accuracy of the hadronic cross section data themselves is the limiting factor in the precise determination of  $g-2$  and  $\Delta\alpha(M_Z^2)$ , the error of the VP (or  $\Delta\alpha(q^2)$ ) is not the limiting factor in its current applications. With the ongoing efforts to measure  $\sigma_{\text{had}}(s)$  with even better accuracy in the whole low energy region, further improvements of the various VP parametrisations are foreseen.

## 7 Summary

In this Report we have summarized the achievements of the last years of the experimental and theoretical groups

working on hadronic cross section measurements and tau physics. In addition we have sketched the prospects in this field for the years to come. We have emphasised the importance of continuous and close collaboration between the experimental and theoretical groups crucial in the quest for precision in hadronic physics. The platform set to simplify this collaboration is a *Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies (Radio Montecarlo)*, for better understanding of the needs and limitations of both experimental and theoretical communities and to facilitate the information flow between them. This Review is a result of the Working Group.

The Report was divided into five Sections covering the luminosity measurements at low energies (up to the energy of  $B$ - factories) (Section 2),  $R$  measurement by energy scan (Section 3),  $R$  measurement using radiative return (Section 4), tau physics (Section 6), and the calculation of the vacuum polarisation with emphasis on the hadronic contributions (Section 5). In all the Sections, with the exception of Section 5, we gave an overview of the experimental results and the status of the Monte Carlo event generators used in the experimental analyses with emphasis on their accuracy and tests.

Concerning the work done on the topic of precision luminosity measurement (Section 2), a particular effort was paid to arrive at an up-to-date estimate of the accuracy of the most precise MC tools used by the experimentalists. Several tuned comparisons between the predictions of independent generators were presented, considering the large-angle Bhabha process with realistic event selection criteria and at different c.m. energies. It turned out that the three most precise luminosity tools, i.e. BabaYaga@NLO, BHWIDE and MCGPJ, agree within 0.1% for the integrated cross sections and within less than 1% for the differential distributions. Therefore the main conclusion of the work on tuned comparisons is that the technical precision of MC programs is well under control, the (minor) discrepancies still observed being due to slightly different details in the treatment of radiative corrections and their implementation. The theoretical accuracy of the generators, as due to radiative corrections partially accounted, was assessed by performing detailed comparisons between the results of the generators and those of exact perturbative calculations. In particular, explicit cross-checks with the predictions of available NNLO QED calculations and with new exact results for lepton and hadron pair corrections led to the conclusion that the total theoretical uncertainty is at the one per mill level for the large-angle Bhabha process at different c.m. energies. Albeit this error estimate could be put on firmer grounds thanks to further work in progress, it appears to be already quite robust and sufficient for precise determination of the luminosity.

In Section 3 we presented the current status of the studies of  $e^+e^-$  annihilation into hadrons and muons at the energies of up to a few GeV. Accurate measurements of the ratio  $R$ , i.e. the ratio of the cross sections of hadron and muon channels, are crucial for the evaluation of the hadronic contribution to vacuum polarisation and subsequently to various precision tests of the Standard Model.

Results of several experimental collaborations have been reviewed for the most important processes with final states of  $\mu^+\mu^-$ ,  $\pi^+\pi^-$ ,  $\pi^+\pi^-\pi^0$ ,  $\pi^+\pi^-2\pi^0$ ,  $\pi^+2\pi^-$ , two kaons and heavier mesons. In particular,  $R$  scans at CMD-2, SND, CLEO and BES experiments have been discussed. Analytic expressions for the Born level cross sections of the main processes have been presented. First-order QED radiative corrections have been given explicitly for the case of muon, pion and kaon pair production. The two latter cases are computed using scalar QED to describe interactions of pseudoscalar mesons with photons in the final state. Matching with higher-order QED corrections evaluated in the leading logarithm approximation have been discussed. Good agreement between different Monte Carlo codes for the muon channel has been shown. The theoretical uncertainty in the description of these processes has been evaluated. For the two main channels  $e^+e^- \rightarrow \mu^+\mu^-$  and  $e^+e^- \rightarrow \pi^+\pi^-$  this uncertainty has been estimated to be of the order of 0.2%.

In Section 4 we have given an overview of experimental measurements via radiative return and described the Monte Carlo generators used in the analyses. Special emphasis has been put on the modelling of the meson-photon interaction, crucial for reaching the accuracy below 1%. Experimentally the radiative return has been applied successfully to the KLOE experiment in Frascati, the BaBar experiment in Stanford and the Belle experiment in Tsukuba, obtaining important results for the measurement of precise hadronic cross sections as well as in the field of hadron spectroscopy. In all three experiments, the ISR physics programme is still going on. New experiments like the BES-III detector at the BEPC-II in Beijing and the experiments at the VEPP-2000 machine in Novosibirsk will use the radiative return method to complement their standard physics programme of energy scanning in the regions of 2 - 4.6 GeV (BEPC-II) and 1 - 2 GeV (VEPP-2000). The success of this program was possible only by close collaboration between experimental and theoretical groups. Dedicated Monte Carlo generators (PHOKHARA, EKHARA, FEVA, FASTERD) were developed to make possible the experimental analysis. The physics program allowed for better modelling of the photon-meson interaction which is crucial for a precise determination of the pion form factor. The measurements of the hadronic cross sections by means of radiative return allowed to reduce the error of the hadronic contribution to the anomalous magnetic moment of the muon and to the running of the fine structure constant. Ongoing and forthcoming measurements will aim at a even better modelling of the hadron-photon interaction and the inclusion of those QED radiative corrections not yet accounted for in the Monte Carlo generators. This ongoing physics program will lead to further improvements in the precision of the calculation of the hadronic contribution to the anomalous magnetic moment of the muon and to the running of the fine structure constant, which in turn is crucial for the tests of the Standard Model and searches for New Physics.

In Section 5 we described the present status of the simulation programs for the production and decay of  $\tau$  lep-

tons. The available programs have been discussed in the context of the required accuracy to match current high-statistics experimental data. After a review of the existing programs used in the data analysis we have emphasised the topics which will require particular attention in the future. We have elaborated on the efforts which are going on at present and focused on the necessary improvements. The techniques for fitting  $\tau$  decay currents require particular attention. The observed spectra and angular distributions are a convolution of theoretical predictions with experimental effects which should be taken into account in the fitting procedures. Background contributions also play an important role if high precision is requested. We have also commented on the impact of these efforts for forthcoming high energy experiments (like at LHC), where  $\tau$  decays are used to constrain hard processes rather than to measure properties of  $\tau$  decays.

In Section 6 the different VP contributions have been discussed, and available parametrisations have been compared. Vacuum polarisation (VP) forms an universal part of radiative corrections and as such is an important ingredient in the Monte Carlo programs. In addition, to evaluate the hadronic contributions to  $g - 2$  and  $\Delta\alpha(q^2)$  via dispersion relations, one has to use ‘undressed’ hadronic cross section data *i.e.* with the VP effects removed. Therefore precise knowledge of VP is required. While in the space-like region the VP is a smooth function and the parametrisations are in excellent agreement, in the time-like region the VP is a fast varying function and differences exist between different parametrisations, especially around resonances. However, the accuracy, which is typically of the order of or below a few per mille and the agreement of the more recent compilations indicate that the current precision of VP is sufficient for the envisaged applications. In the future better hadronic cross section data will lead to further improved accuracy.

## Acknowledgements

This work was supported in part by:

European Union Marie-Curie Research Training Networks MRTN-CT-2006-035482 “FLAVIANet” and MRTN-CT-2006-035505 “HEPTOOLS”;

European Union Research Programmes at LNF, FP7, Transnational Access to Research Infrastructure (TARI), Hadron Physics2-Integrating Activity, Contract No. 227431;

Generalitat Valenciana

under Grant No. PROMETEO/2008/069;

German Federal Ministry of Education and Research (BMBF) grants 05HT4VKA/3, 06-KA-202 and 06-MZ-9171I;

German Research Foundation (DFG): ‘Emmy Noether Programme’, contracts DE839/1-4, ‘Heisenberg Programme’ and Sonderforschungsbereich/Transregio SFB/TRR 9;

Initiative and Networking Fund of the Helmholtz Association, contract HA-101 (“Physics at the Terascale”);

INTAS project Nr 05-1000008-8328 “Higher-order effects in  $e^+e^-$  annihilation and muon anomalous magnetic moment”;

Ministerio de Ciencia e Innovación under Grant No. FPA2007-60323, and CPAN (Grant No. CSD2007-00042);

National Natural Science Foundation of China under Contracts Nos. 10775142, 10825524 and 10935008;  
Polish Government grant N202 06434 (2008-2010);  
PST.CLG.980342

RFBR grants 03-02-16477, 04-02-16217, 04-02-1623, 04-02-16443, 04-02-16181-a, 04-02-16184-a, 05-02-16250-a, 06-02-16192-a, 07-02-00816-a, 08-02-13516 08-02-91969 and 09-02-01143;

US DOE contract DE-FG02-09ER41600.

We thank J. Libby for useful correspondence about the luminosity measurement at CLEO-c. S.E. and V.C. are grateful to the Cracow Institute of Nuclear Physics where part of this work has been performed.

## References

1. D. Hanneke, S. Fogwell, G. Gabrielse, Phys. Rev. Lett. **100**, 120801 (2008), 0801.1134
2. F. Jegerlehner, J. Phys. **G29**, 101 (2003), hep-ph/0104304
3. F. Jegerlehner, Nucl. Phys. Proc. Suppl. **131**, 213 (2004), hep-ph/0312372
4. F. Jegerlehner, Nucl. Phys. Proc. Suppl. **162**, 22 (2006), hep-ph/0608329
5. F. Jegerlehner, Nucl. Phys. Proc. Suppl. **181-182**, 135 (2008), 0807.4206
6. N. Cabibbo, R. Gatto, Phys. Rev. **124**, 1577 (1961)
7. S. Eidelman, F. Jegerlehner, Z. Phys. **C67**, 585 (1995), hep-ph/9502298
8. M. Davier, A. Hocker, Phys. Lett. **B419**, 419 (1998), hep-ph/9711308
9. H. Burkhardt, B. Pietrzyk, Phys. Lett. **B513**, 46 (2001)
10. H. Burkhardt, B. Pietrzyk, Phys. Rev. **D72**, 057501 (2005), hep-ph/0506323
11. F. Jegerlehner, Nucl. Phys. Proc. Suppl. **126**, 325 (2004), hep-ph/0310234
12. F. Jegerlehner (2003), hep-ph/0308117
13. K. Hagiwara, A.D. Martin, D. Nomura, T. Teubner, Phys. Rev. **D69**, 093003 (2004), hep-ph/0312250
14. K. Hagiwara, A.D. Martin, D. Nomura, T. Teubner, Phys. Lett. **B649**, 173 (2007), hep-ph/0611102
15. (2008), 0811.4682
16. M. Gourdin, E. De Rafael, Nucl. Phys. **B10**, 667 (1969)
17. M. Davier, A. Hoecker, B. Malaescu, C.Z. Yuan, Z. Zhang (2009), 0908.4300
18. F. Jegerlehner, Nucl. Phys. Proc. Suppl. **181-182**, 26 (2008)
19. F. Jegerlehner, A. Nyffeler, Phys. Rept. **477**, 1 (2009), 0902.3360
20. S.I. Eidelman (2009), 0904.3275
21. J. Prades, E. de Rafael, A. Vainshtein (2009), 0901.0306
22. J. Prades (2009), 0909.2546
23. G.W. Bennett et al. (Muon G-2), Phys. Rev. **D73**, 072003 (2006), hep-ex/0602035
24. R.M. Carey et al., FERMILAB-PROPOSAL-0989 (2009)
25. J. Imazato, Nucl. Phys. Proc. Suppl. **129**, 81 (2004)
26. R. Alemany, M. Davier, A. Hocker, Eur. Phys. J. **C2**, 123 (1998), hep-ph/9703220
27. M. Davier, S. Eidelman, A. Hocker, Z. Zhang, Eur. Phys. J. **C27**, 497 (2003), hep-ph/0208177
28. M. Davier, S. Eidelman, A. Hocker, Z. Zhang, Eur. Phys. J. **C31**, 503 (2003), hep-ph/0308213
29. M. Davier et al. (2009), 0906.5443
30. F. Ambrosino et al. (KLOE), Eur. Phys. J. **C47**, 589 (2006), hep-ex/0604048
31. H. Bhabha, Proc. Roy. Soc. **A154**, 195 (1936)
32. G. Barbiellini et al., Nucl. Instrum. Meth. **123**, 125 (1975)
33. H.C. Dehne, M.A. Preger, S. Tazzari, G. Vignola, Nucl. Instrum. Meth. **116**, 345 (1974)
34. S. Jadach et al. (1996), hep-ph/9602393
35. A.B. Arbuzov et al., Nucl. Phys. **B485**, 457 (1997), hep-ph/9512344
36. G. Montagna, O. Nicrosini, F. Piccinini, Phys. Lett. **B385**, 348 (1996), hep-ph/9605252
37. A. Arbuzov et al., Phys. Lett. **B383**, 238 (1996), hep-ph/9605239
38. G. Montagna, O. Nicrosini, F. Piccinini, Riv. Nuovo Cim. **21N9**, 1 (1998), hep-ph/9802302
39. B.F.L. Ward, S. Jadach, M. Melles, S.A. Yost, Phys. Lett. **B450**, 262 (1999), hep-ph/9811245
40. S. Jadach (2003), hep-ph/0306083
41. S. Dobbs et al. (CLEO), Phys. Rev. **D76**, 112001 (2007), 0709.3783
42. A. Hafner, Diploma thesis, University of Karlsruhe (2007)
43. D.M. Asner et al. (2008), 0809.1869
44. L.M. Brown, R.P. Feynman, Phys. Rev. **85**, 231 (1952)
45. F. Redhead, Proc. Roy. Soc. **220**, 219 (1953)
46. R. Polovin, JETP **31**, 449 (1956)
47. M. Consoli, Nucl. Phys. **B160**, 208 (1979)
48. M. Böhm, A. Denner, W. Hollik, R. Sommer, Phys. Lett. **B144**, 414 (1984)
49. D. Bardin, W. Hollik, T. Riemann, Z. Phys. **C49**, 485 (1991)
50. D. Bardin, P. Christova, M. Jack, L. Kalinovskaya, A. Olchevski, S. Riemann, T. Riemann, Comput. Phys. Commun. **133**, 229 (2001), hep-ph/9908433
51. A. Arbuzov, M. Awramik, M. Czakon, A. Freitas, M. Grünewald, K. Mönig, S. Riemann, T. Riemann, Comput. Phys. Commun. **174**, 728 (2006), hep-ph/0507146
52. G. Montagna, F. Piccinini, O. Nicrosini, G. Passarino, R. Pittau, Nucl. Phys. **B401**, 3 (1993)
53. G. Montagna, F. Piccinini, O. Nicrosini, G. Passarino, R. Pittau, Comput. Phys. Commun. **76**, 328 (1993)
54. G. Montagna, O. Nicrosini, F. Piccinini, G. Passarino, Comput. Phys. Commun. **117**, 278 (1999), hep-ph/9804211
55. A. Djouadi, C. Verzegnassi, Phys. Lett. **B195**, 265 (1987)
56. A. Djouadi, Nuovo Cim. **A100**, 357 (1988)
57. B.A. Kniehl, J.H. Kuhn, R.G. Stuart, Phys. Lett. **B214**, 621 (1988)
58. J.J. van der Bij, F. Hoogeveen, Nucl. Phys. **B283**, 477 (1987)
59. R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, A. Vicere, Nucl. Phys. **B409**, 105 (1993)
60. J. Fleischer, O.V. Tarasov, F. Jegerlehner, Phys. Lett. **B319**, 249 (1993)
61. J. Fleischer, O.V. Tarasov, F. Jegerlehner, Phys. Rev. **D51**, 3820 (1995)
62. R. Boughezal, M. Czakon, Nucl. Phys. **B755**, 221 (2006), hep-ph/0606232
63. K.G. Chetyrkin, M. Faisst, J.H. Kuhn, P. Maierhofer, C. Sturm, Phys. Rev. Lett. **97**, 102003 (2006), hep-ph/0605201



64. Y. Schroder, M. Steinhauser, Phys. Lett. **B622**, 124 (2005), [hep-ph/0504055](#)
65. K.G. Chetyrkin, J.H. Kuhn, M. Steinhauser, Phys. Lett. **B351**, 331 (1995), [hep-ph/9502291](#)
66. K.G. Chetyrkin, J.H. Kuhn, M. Steinhauser, Phys. Rev. Lett. **75**, 3394 (1995), [hep-ph/9504413](#)
67. R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, A. Vicere, Phys. Lett. **B288**, 95 (1992), [hep-ph/9205238](#)
68. J.J. van der Bij, K.G. Chetyrkin, M. Faisst, G. Jikia, T. Seidensticker, Phys. Lett. **B498**, 156 (2001), [hep-ph/0011373](#)
69. M. Faisst, J.H. Kuhn, T. Seidensticker, O. Veretin, Nucl. Phys. **B665**, 649 (2003), [hep-ph/0302275](#)
70. R. Boughezal, J.B. Tausk, J.J. van der Bij, Nucl. Phys. **B713**, 278 (2005), [hep-ph/0410216](#)
71. R. Boughezal, J.B. Tausk, J.J. van der Bij, Nucl. Phys. **B725**, 3 (2005), [hep-ph/0504092](#)
72. M. Awramik, M. Czakon, A. Freitas, G. Weiglein, Phys. Rev. **D69**, 053006 (2004), [hep-ph/0311148](#)
73. M. Awramik, M. Czakon, A. Freitas, G. Weiglein, Phys. Rev. Lett. **93**, 201805 (2004), [hep-ph/0407317](#)
74. W. Hollik, U. Meier, S. Uccirati, Nucl. Phys. **B731**, 213 (2005), [hep-ph/0507158](#)
75. W. Hollik, U. Meier, S. Uccirati, Nucl. Phys. **B765**, 154 (2007), [hep-ph/0610312](#)
76. M. Awramik, M. Czakon, A. Freitas, JHEP **11**, 048 (2006), [hep-ph/0608099](#)
77. M. Awramik, M. Czakon, A. Freitas, Phys. Lett. **B642**, 563 (2006), [hep-ph/0605339](#)
78. J.H. Kuhn, A.A. Penin, V.A. Smirnov, Eur. Phys. J. **C17**, 97 (2000), [hep-ph/9912503](#)
79. J.H. Kuhn, A.A. Penin, V.A. Smirnov, Nucl. Phys. Proc. Suppl. **89**, 94 (2000), [hep-ph/0005301](#)
80. J.H. Kuhn, S. Moch, A.A. Penin, V.A. Smirnov, Nucl. Phys. **B616**, 286 (2001), [hep-ph/0106298](#)
81. B. Feucht, J.H. Kuhn, A.A. Penin, V.A. Smirnov, Phys. Rev. Lett. **93**, 101802 (2004), [hep-ph/0404082](#)
82. B. Jantzen, J.H. Kuhn, A.A. Penin, V.A. Smirnov, Phys. Rev. **D72**, 051301 (2005), [hep-ph/0504111](#)
83. B. Jantzen, J.H. Kuhn, A.A. Penin, V.A. Smirnov, Nucl. Phys. **B731**, 188 (2005), [hep-ph/0509157](#)
84. A. Denner, B. Jantzen, S. Pozzorini, Nucl. Phys. **B761**, 1 (2007), [hep-ph/0608326](#)
85. A. Denner, B. Jantzen, S. Pozzorini (2008), 0801.2647
86. F. Berends, K. Gaemers, R. Gastmans, Nucl. Phys. **B68**, 541 (1974)
87. W. Beenakker, F. Berends, S. van der Marck, Nucl. Phys. **B349**, 323 (1991)
88. M. Terentyev, Yad. Fiz. **9**, 1212 (1969)
89. F. Berends, R. Gastmans, Nucl. Phys. **B61**, 414 (1973)
90. S. Eidelman, E. Kuraev, Nucl. Phys. **B143**, 353 (1978)
91. F. Berends, R. Kleiss, P. De Causmaecker, R. Gastmans, W. Troost, T. Wu, Nucl. Phys. **B206**, 61 (1982)
92. M. Bohm, Z. Sack, Z. Phys. **C33**, 157 (1986)
93. S. Actis, P. Mastrolia, G. Ossola (2009), 0909.1750
94. A.B. Arbuzov, E.A. Kuraev, N.P. Merenkov, L. Trentadue, Phys. Atom. Nucl. **60**, 591 (1997), [http://ccdb4fs.kek.jp/cgi-bin/img\\_index?9604099](http://ccdb4fs.kek.jp/cgi-bin/img_index?9604099)
95. A.B. Arbuzov, E.A. Kuraev, N.P. Merenkov, L. Trentadue, J. Exp. Theor. Phys. **81**, 638 (1995), [hep-ph/9509405](#)
96. A.B. Arbuzov, E.A. Kuraev, N.P. Merenkov, L. Trentadue, Nucl. Phys. **B474**, 271 (1996)
97. S. Laporta, E. Remiddi, Phys. Lett. **B379**, 283 (1996), [hep-ph/9602417](#)
98. S. Laporta, Int. J. Mod. Phys. **A15**, 5087 (2000), [hep-ph/0102033](#)
99. F.V. Tkachov, Phys. Lett. **B100**, 65 (1981)
100. K. Chetyrkin, F. Tkachov, Nucl. Phys. **B192**, 159 (1981)
101. A.V. Kotikov, Phys. Lett. **B254**, 158 (1991)
102. A. Kotikov, Phys. Lett. **B259**, 314 (1991)
103. A.V. Kotikov, Phys. Lett. **B267**, 123 (1991)
104. E. Remiddi, Nuovo Cim. **A110**, 1435 (1997), [hep-th/9711188](#)
105. M. Caffo, H. Czyz, S. Laporta, E. Remiddi, Acta Phys. Polon. **B29**, 2627 (1998), [hep-th/9807119](#)
106. M. Caffo, H. Czyz, S. Laporta, E. Remiddi, Nuovo Cim. **A111**, 365 (1998), [hep-th/9805118](#)
107. M. Argeri, P. Mastrolia, Int. J. Mod. Phys. **A22**, 4375 (2007), 0707.4037
108. V. Smirnov, "Evaluating Feynman Integrals" (Springer Verlag, Berlin, 2004)
109. S. Friot, D. Greynat, E. De Rafael, Phys. Lett. **B628**, 73 (2005), [hep-ph/0505038](#)
110. N. Usyukina, Teor. Mat. Fiz. **22**, 300 (1975)
111. V. Smirnov, Phys. Lett. **B460**, 397 (1999), [hep-ph/9905323](#)
112. B. Tausk, Phys. Lett. **B469**, 225 (1999), [hep-ph/9909506](#)
113. V. Smirnov, O. Veretin, Nucl. Phys. **B566**, 469 (2000), [hep-ph/9907385](#)
114. V.A. Smirnov, Phys. Lett. **B524**, 129 (2002), [hep-ph/0111160](#)
115. G. Heinrich, V.A. Smirnov, Phys. Lett. **B598**, 55 (2004), [hep-ph/0406053](#)
116. M. Czakon, Comput. Phys. Commun. **175**, 559 (2006), [hep-ph/0511200](#)
117. J. Gluza, K. Kajda, T. Riemann, Comput. Phys. Commun. **177**, 879 (2007), [arXiv:0704.2423](#)
118. A. Goncharov, Math. Res. Letters **5**, 497 (1998), <http://www.math.uiuc.edu/K-theory/0297>
119. D.J. Broadhurst, Eur. Phys. J. **C8**, 311 (1999), [hep-th/9803091](#)
120. E. Remiddi, J. Vermaseren, Int. J. Mod. Phys. **A15**, 725 (2000), [hep-ph/9905237](#)
121. T. Gehrmann, E. Remiddi, Comput. Phys. Commun. **141**, 296 (2001), [<http://arXiv.org/abs/hep-ph/0107173>]
122. T. Gehrmann, E. Remiddi, Comput. Phys. Commun. **144**, 200 (2002), [hep-ph/0111255](#)
123. D. Maitre, Comput. Phys. Commun. **174**, 222 (2006), [hep-ph/0507152](#)
124. D. Maitre, *Extension of HPL to complex arguments* (2007), [hep-ph/0703052](#)
125. J. Vollinga, S. Weinzierl, Comput. Phys. Commun. **167**, 177 (2005), [hep-ph/0410259](#)
126. S. Weinzierl (2007), 0705.0900
127. A.A. Penin, Phys. Rev. Lett. **95**, 010408 (2005), [hep-ph/0501120](#)
128. A.A. Penin, Nucl. Phys. **B734**, 185 (2006), [hep-ph/0508127](#)
129. A. Mitov, S. Moch, JHEP **05**, 001 (2007), [hep-ph/0612149](#)
130. T. Becher, K. Melnikov, JHEP **06**, 084 (2007), [arXiv:0704.3582 \[hep-ph\]](#)
131. Z. Bern, L. Dixon, A. Ghinculov, Phys. Rev. **D63**, 053007 (2001), [hep-ph/0010075](#)



132. R. Bonciani, A. Ferroglia, Phys. Rev. **D72**, 056004 (2005), [hep-ph/0507047](#)
133. R. Bonciani, P. Mastrolia, E. Remiddi, Nucl. Phys. **B661**, 289 (2003), [hep-ph/0301170](#)
134. R. Bonciani, P. Mastrolia, E. Remiddi, Nucl. Phys. **B690**, 138 (2004), [hep-ph/0311145](#)
135. R. Bonciani, P. Mastrolia, E. Remiddi, Nucl. Phys. **B676**, 399 (2004), [hep-ph/0307295](#)
136. M. Czakon, J. Gluza, T. Riemann, Phys. Rev. **D71**, 073009 (2005), [hep-ph/0412164](#)
137. M. Czakon, J. Gluza, T. Riemann, Nucl. Phys. **B751**, 1 (2006), [hep-ph/0604101](#)
138. D.J. Broadhurst, J. Fleischer, O.V. Tarasov, Z. Phys. **C60**, 287 (1993), [hep-ph/9304303](#)
139. A.I. Davydychev, M.Y. Kalmykov, Nucl. Phys. **B699**, 3 (2004), [hep-th/0303162](#)
140. A.B. Arbuzov, E.A. Kuraev, B.G. Shaikhatdenov, Mod. Phys. Lett. **A13**, 2305 (1998), [hep-ph/9806215](#)
141. E.W.N. Glover, J.B. Tausk, J.J. Van der Bij, Phys. Lett. **B516**, 33 (2001), [hep-ph/0106052](#)
142. G.J.H. Burgers, Phys. Lett. **B164**, 167 (1985)
143. R. Bonciani, A. Ferroglia, P. Mastrolia, E. Remiddi, J. van der Bij, Nucl. Phys. **B681**, 261 (2004), [hep-ph/0310333](#)
144. R. Bonciani, A. Ferroglia, P. Mastrolia, E. Remiddi, J. van der Bij, Nucl. Phys. **B701**, 121 (2004), [hep-ph/0405275](#)
145. R. Bonciani, A. Ferroglia, P. Mastrolia, E. Remiddi, J.J. van der Bij, Nucl. Phys. **B716**, 280 (2005), [hep-ph/0411321](#)
146. S. Actis, M. Czakon, J. Gluza, T. Riemann, Nucl. Phys. **B786**, 26 (2007), [arXiv:0704.2400v.2 \[hep-ph\]](#)
147. R. Bonciani, A. Ferroglia, A.A. Penin, Phys. Rev. Lett. **100**, 131601 (2008), [0710.4775](#)
148. R. Bonciani, A. Ferroglia, A.A. Penin, JHEP **02**, 080 (2008), [arXiv:0802.2215](#)
149. S. Actis, M. Czakon, J. Gluza, T. Riemann, Phys. Rev. Lett. **100**, 131602 (2008), [arXiv:0711.3847](#)
150. S. Actis, M. Czakon, J. Gluza, T. Riemann, Phys. Rev. **D78**, 085019 (2008), [arXiv:0807.4691](#)
151. J. Frenkel, J.C. Taylor, Nucl. Phys. **B116**, 185 (1976)
152. J. Fleischer, A.V. Kotikov, O.L. Veretin, Nucl. Phys. **B547**, 343 (1999), [hep-ph/9808242](#)
153. U. Aglietti, R. Bonciani, Nucl. Phys. **B668**, 3 (2003), [hep-ph/0304028](#)
154. U. Aglietti, R. Bonciani, Nucl. Phys. **B698**, 277 (2004), [hep-ph/0401193](#)
155. R.E. Cutkosky, J. Math. Phys. **1**, 429 (1960)
156. Fortran program *rintpl.f*, the Fortran program *rintpl.F* is based on the data compilation performed for [13, 14]. The publication is in preparation. The routine is available upon request from the authors, E-mails: dno-mura@post.kek.jp, thomas.teubner@liverpool.ac.uk.
157. R.V. Harlander, M. Steinhauser, Comput. Phys. Commun. **153**, 244 (2003), [hep-ph/0212294](#)
158. F. Berends, G. Komen, Phys. Lett. **B63**, 432 (1976)
159. B. Kniehl, M. Krawczyk, J. Kühn, R. Stuart, Phys. Lett. **B209**, 337 (1988)
160. T. van Ritbergen, R.G. Stuart, Phys. Lett. **B437**, 201 (1998), [hep-ph/9802341](#)
161. R. Barbieri, J.A. Mignaco, E. Remiddi, Nuovo Cim. **A11**, 824 (1972)
162. R. Barbieri, J.A. Mignaco, E. Remiddi, Nuovo Cim. **A11**, 865 (1972)
163. P. Mastrolia, E. Remiddi, Nucl. Phys. **B664**, 341 (2003), [hep-ph/0302162](#)
164. S. Actis, M. Czakon, J. Gluza, T. Riemann, Acta Phys. Polon. **B38**, 3517 (2007), [arXiv:0710.5111](#)
165. J.H. Kühn, S. Uccirati, Nucl. Phys. **B806**, 300 (2009), [arXiv:0807.1284](#)
166. DESY Zeuthen webpage: <http://www-zeuthen.desy.de/theory/research/bhabha/bhabha.html>
167. H. Burkhardt, *New numerical analysis of the hadronic vacuum polarization*, TASSO-NOTE-192 (1981), and Fortran program *repi.f* (1986)
168. G. Passarino, M. Veltman, Nucl. Phys. **B160**, 151 (1979)
169. Z. Bern, L.J. Dixon, D.A. Kosower, Ann. Rev. Nucl. Part. Sci. **46**, 109 (1996), [hep-ph/9602280](#)
170. L.J. Dixon (1996), [hep-ph/9601359](#)
171. T. Binoth, J.P. Guillet, G. Heinrich, Nucl. Phys. **B572**, 361 (2000), [hep-ph/9911342](#)
172. A. Denner, S. Dittmaier, Nucl. Phys. **B658**, 175 (2003), [hep-ph/0212259](#)
173. T. Binoth, J.P. Guillet, G. Heinrich, E. Pilon, C. Schubert, JHEP **10**, 015 (2005), [hep-ph/0504267](#)
174. A. Denner, S. Dittmaier, Nucl. Phys. **B734**, 62 (2006), [hep-ph/0509141](#)
175. A. Denner, S. Dittmaier, Nucl. Phys. Proc. Suppl. **157**, 53 (2006), [hep-ph/0601085](#)
176. A. Davydychev, Phys. Lett. **B263**, 107 (1991)
177. O.V. Tarasov, Phys. Rev. **D54**, 6479 (1996), [hep-th/9606018](#)
178. J. Fleischer, F. Jegerlehner, O.V. Tarasov, Nucl. Phys. **B566**, 423 (2000), [hep-ph/9907327](#)
179. D.B. Melrose, Nuovo Cim. **40**, 181 (1965)
180. T. Diakonidis, J. Fleischer, J. Gluza, K. Kajda, T. Riemann, J. Tausk (2008), [arXiv:0807.2984](#)
181. T. Diakonidis, J. Fleischer, J. Gluza, K. Kajda, T. Riemann, J. Tausk (2008), [0812.2134](#)
182. Z. Bern, L.J. Dixon, D.A. Kosower, Annals Phys. **322**, 1587 (2007), [0704.2798](#)
183. L.D. Landau, Nucl. Phys. **13**, 181 (1959)
184. S. Mandelstam, Phys. Rev. **112**, 1344 (1958)
185. S. Mandelstam, Phys. Rev. **115**, 1741 (1959)
186. R.J. Eden, P.V. Landshoff, D.I. Olive, J.C. Polkinghorne, Cambridge University Press (1966)
187. G. 't Hooft, M. Veltman, Nucl. Phys. **B153**, 365 (1979)
188. Z. Bern, L.J. Dixon, D.A. Kosower, Phys. Lett. **B302**, 299 (1993), [hep-ph/9212308](#)
189. Z. Bern, L.J. Dixon, D.A. Kosower, Nucl. Phys. **B412**, 751 (1994), [hep-ph/9306240](#)
190. G. Duplancic, B. Nizic, Eur. Phys. J. **C35**, 105 (2004), [hep-ph/0303184](#)
191. R.K. Ellis, G. Zanderighi, JHEP **02**, 002 (2008), [0712.1851](#)
192. R. Britto, F. Cachazo, B. Feng, Nucl. Phys. **B725**, 275 (2005), [hep-th/0412103](#)
193. P. Mastrolia, Phys. Lett. **B644**, 272 (2007), [hep-th/0611091](#)
194. D. Forde, Phys. Rev. **D75**, 125019 (2007), [0704.1835](#)
195. N.E.J. Bjerrum-Bohr, D.C. Dunbar, W.B. Perkins, JHEP **04**, 038 (2008), [0709.2086](#)
196. R. Britto, E. Buchbinder, F. Cachazo, B. Feng, Phys. Rev. **D72**, 065012 (2005), [hep-ph/0503132](#)

197. R. Britto, B. Feng, P. Mastrolia, Phys. Rev. **D73**, 105004 (2006), [hep-ph/0602178](#)
198. C. Anastasiou, R. Britto, B. Feng, Z. Kunszt, P. Mastrolia, Phys. Lett. **B645**, 213 (2007), [hep-ph/0609191](#)
199. C. Anastasiou, R. Britto, B. Feng, Z. Kunszt, P. Mastrolia, JHEP **03**, 111 (2007), [hep-ph/0612277](#)
200. R. Britto, B. Feng, Phys. Rev. **D75**, 105006 (2007), [hep-ph/0612089](#)
201. E.W. Nigel Glover, C. Williams, JHEP **12**, 067 (2008), 0810.2964
202. R. Britto, B. Feng, P. Mastrolia, Phys. Rev. **D78**, 025031 (2008), 0803.1989
203. G. Ossola, C.G. Papadopoulos, R. Pittau, Nucl. Phys. **B763**, 147 (2007), [hep-ph/0609007](#)
204. G. Ossola, C.G. Papadopoulos, R. Pittau, JHEP **07**, 085 (2007), 0704.1271
205. F. del Aguila, R. Pittau, JHEP **07**, 017 (2004), [hep-ph/0404120](#)
206. R. Pittau (2004), [hep-ph/0406105](#)
207. R. Pittau, Comput. Phys. Commun. **104**, 23 (1997), [hep-ph/9607309](#)
208. R. Pittau, Comput. Phys. Commun. **111**, 48 (1998), [hep-ph/9712418](#)
209. P. Mastrolia, G. Ossola, C.G. Papadopoulos, R. Pittau, JHEP **06**, 030 (2008), 0803.3964
210. G. Ossola, C.G. Papadopoulos, R. Pittau, JHEP **05**, 004 (2008), 0802.1876
211. A. Kanaki, C.G. Papadopoulos, Comput. Phys. Commun. **132**, 306 (2000), [hep-ph/0002082](#)
212. C.G. Papadopoulos, Comput. Phys. Commun. **137**, 247 (2001), [hep-ph/0007335](#)
213. C.G. Papadopoulos, M. Worek, Eur. Phys. J. **C50**, 843 (2007), [hep-ph/0512150](#)
214. A. Cafarella, C.G. Papadopoulos, M. Worek (2007), 0710.2427
215. H. Czyz, E. Nowak-Kubat, Acta Phys. Polon. **B36**, 3425 (2005), [hep-ph/0510287](#)
216. H. Czyz, E. Nowak-Kubat, Phys. Lett. **B634**, 493 (2006), [hep-ph/0601169](#)
217. H. Czyz, M. Gunia, unpublished (2009)
218. M. Caffo, H. Czyz, Comput. Phys. Commun. **100**, 99 (1997), [hep-ph/9607357](#)
219. G. Montagna, M. Moretti, O. Nicrosini, A. Pallavicini, F. Piccinini, Nucl. Phys. **B547**, 39 (1999), [hep-ph/9811436](#)
220. E. Kuraev, V. Fadin, Sov. J. Nucl. Phys. **41**, 466 (1985)
221. G. Altarelli, G. Martinelli (1986), in Ellis, J. (Ed.), Peccei, R.d. (Ed.): Physics At Lep, Vol. 1, 47-57
222. O. Nicrosini, L. Trentadue, Phys. Lett. **B196**, 551 (1987)
223. O. Nicrosini, L. Trentadue, Z. Phys. **C39**, 479 (1988)
224. D. Yennie, S. Frautschi, H. Suura, Annals Phys. **13**, 379 (1961)
225. C.M. Carloni Calame, C. Lunardini, G. Montagna, O. Nicrosini, F. Piccinini, Nucl. Phys. **B584**, 459 (2000), [hep-ph/0003268](#)
226. C.M. Carloni Calame, G. Montagna, O. Nicrosini, F. Piccinini, Nucl. Phys. Proc. Suppl. **131**, 48 (2004), [hep-ph/0312014](#)
227. G. Balossini, C.M. Carloni Calame, G. Montagna, O. Nicrosini, F. Piccinini, Nucl. Phys. **B758**, 227 (2006), [hep-ph/0607181](#)
228. A.B. Arbuzov, G.V. Fedotov, F.V. Ignatov, E.A. Kuraev, A.L. Sibidanov, Eur. Phys. J. **C46**, 689 (2006), [hep-ph/0504233](#)
229. S. Jadach, W. Placzek, B.F.L. Ward, Phys. Lett. **B390**, 298 (1997), [hep-ph/9608412](#)
230. V. Gribov, L. Lipatov, Sov. J. Nucl. Phys. **15**, 675 (1972)
231. G. Altarelli, G. Parisi, Nucl. Phys. **B126**, 298 (1977)
232. Y. Dokshitzer, Sov. Phys. JETP **46**, 641 (1977)
233. M. Cacciari, A. Deandrea, G. Montagna, O. Nicrosini, Europhys. Lett. **17**, 123 (1992)
234. M. Skrzypek, Acta Phys. Polon. **B24**, 123 (1992)
235. M. Przybycien, Acta Phys. Polon. **B24**, 1105 (1993)
236. A.B. Arbuzov, Phys. Lett. **B470**, 252 (1999)
237. A. Arbuzov, G. Fedotov, E. Kuraev, N. Merenkov, V. Rushai, L. Trentadue, JHEP **9710**, 001 (1997), 9702262
238. A.B. Arbuzov, E. Scherbakova, JETP Lett. **83**, 427 (2006)
239. M. Greco, O. Nicrosini, Phys. Lett. **B240**, 219 (1990)
240. C.M. Carloni Calame, Phys. Lett. **B520**, 16 (2001), [hep-ph/0103117](#)
241. A.B. Arbuzov, E. Scherbakova, Phys. Lett. **B660**, 37 (2008)
242. C.M. Carloni Calame, G. Montagna, O. Nicrosini, M. Treccani, Phys. Rev. **D69**, 037301 (2004), [hep-ph/0303102](#)
243. C.M. Carloni Calame, G. Montagna, O. Nicrosini, M. Treccani, JHEP **05**, 019 (2005), [hep-ph/0502218](#)
244. C.M. Carloni Calame, G. Montagna, O. Nicrosini, A. Vicini, JHEP **12**, 016 (2006), [hep-ph/0609170](#)
245. C.M. Carloni Calame, G. Montagna, O. Nicrosini, A. Vicini, JHEP **10**, 109 (2007), 0710.1722
246. K.T. Mahanthappa, Phys. Rev. **126**, 329 (1962)
247. M. Bohm, A. Denner, W. Hollik, Nucl. Phys. **B304**, 687 (1988)
248. E. Drago, G. Venanzoni, *A Bhabha generator for DAPHNE including radiative corrections and Phi resonance*, INFN-AE-97-48 (1997)
249. F. Berends, R. Kleiss, Nucl. Phys. **B228**, 537 (1983)
250. F. Berends, R. Kleiss, Nucl. Phys. **B186**, 22 (1981)
251. F. Jegerlehner, Z. Phys. **C32**, 195 (1986)
252. H. Burkhardt, F. Jegerlehner, G. Penso, C. Verzegnassi, Z. Phys. **C43**, 497 (1989)
253. G. Balossini et al., Phys. Lett. **B663**, 209 (2008), 0801.3360
254. P. Rong-Gang (2009), private communication
255. S. Jadach, M. Melles, B.F.L. Ward, S.A. Yost, Phys. Lett. **B377**, 168 (1996), [hep-ph/9603248](#)
256. C. Glosser, S. Jadach, B.F.L. Ward, S.A. Yost, Phys. Lett. **B605**, 123 (2005), [hep-ph/0406298](#)
257. S. Jadach, B.F.L. Ward, S.A. Yost, Phys. Rev. **D73**, 073001 (2006), [hep-ph/0602197](#)
258. C. Carloni Calame et al., in preparation (????)
259. C. Amsler et al. (Particle Data Group), Phys. Lett. **B667**, 1 (2008)
260. R.R. Akhmetshin et al. (CMD-2), Phys. Lett. **B527**, 161 (2002), [hep-ex/0112031](#)
261. E. Shintani, S. Aoki, T. Chiu et al., Phys. Rev. **D79**, 074510 (2009)
262. T. Kinoshita, B. Nizic, Y. Okamoto, Phys. Rev. **D31**, 2108 (1985)
263. B.E. Lautrup, A. Peterman, E. de Rafael, Phys. Rept. **3**, 193 (1972)
264. F.A. Berends, R. Kleiss, Nucl. Phys. **B177**, 237 (1981)
265. A.B. Arbuzov, D.Y. Bardin, A. Leike, Mod. Phys. Lett. **A7**, 2029 (1992)

266. A. Bramon, M. Greco (1997), in \*Maiani, L. (ed.) et al.: The second DAPHNE physics handbook, vol. 2\* 451-466
267. A.I. Ahmedov, G.V. Fedotov, E.A. Kuraev, Z.K. Silagadze, JHEP **09**, 008 (2002), [hep-ph/0201157](#)
268. E.A. Kuraev, Z.K. Silagadze, Phys. Atom. Nucl. **58**, 1589 (1995), [hep-ph/9502406](#)
269. A.B. Arbuzov, Nuovo Cim. **107A**, 1263 (1994)
270. S.I. Eidelman, E.A. Kuraev, V.S. Panin, Nucl. Phys. **B148**, 245 (1979)
271. F.A. Berends, K.J.F. Gaemers, R. Gastmans, Nucl. Phys. **B57**, 381 (1973)
272. S. Jadach, B.F.L. Ward, Z. Was, Comput. Phys. Commun. **130**, 260 (2000), [hep-ph/9912214](#)
273. S. Jadach, B.F.L. Ward, Z. Was, Phys. Rev. **D63**, 113009 (2001), [hep-ph/0006359](#)
274. A. Arbuzov, V. Astakhov, A. Fedorov, G. Fedotov, E. Kuraev, N. Merenkov, A. Sibidanov, JHEP **9710**, 006 (1997)
275. J.S. Schwinger, *Particles, Sources, and Fields.*, Vol. Vol. 3 (Redwood City, USA: Addison-Wesley, 1989)
276. K. Melnikov, Int. J. Mod. Phys. **A16**, 4591 (2001), [hep-ph/0105267](#)
277. A. Hoefer, J. Gluza, F. Jegerlehner, Eur. Phys. J. **C24**, 51 (2002), [hep-ph/0107154](#)
278. E. Anashkin, V. Aulchenko, S. Baru et al. (1988), iCFA Instr. Bulletin 5 (1988) 18
279. M.N. Achasov et al., Nucl. Instrum. Meth. **A449**, 125 (2000), [hep-ex/9909015](#)
280. R.R. Akhmetshin et al. (CMD-2) (1999), [hep-ex/9904027](#)
281. M.N. Achasov et al., J. Exp. Theor. Phys. **101**, 1053 (2005), [hep-ex/0506076](#)
282. M.N. Achasov et al., J. Exp. Theor. Phys. **103**, 380 (2006), [hep-ex/0605013](#)
283. R.R. Akhmetshin et al. (CMD-2), Phys. Lett. **B578**, 285 (2004), [hep-ex/0308008](#)
284. M.N. Achasov et al., Phys. Rev. **D66**, 032001 (2002), [hep-ex/0201040](#)
285. M.N. Achasov et al., Phys. Rev. **D68**, 052006 (2003), [hep-ex/0305049](#)
286. R.R. Akhmetshin et al., Phys. Lett. **B642**, 203 (2006)
287. A. Aloisio et al. (KLOE), Phys. Lett. **B561**, 55 (2003), [hep-ex/0303016](#)
288. R.R. Akhmetshin et al. (CMD2), Phys. Lett. **B466**, 392 (1999), [hep-ex/9904024](#)
289. M.N. Achasov et al., J. Exp. Theor. Phys. **96**, 789 (2003)
290. R.R. Akhmetshin et al. (CMD-2), Phys. Lett. **B595**, 101 (2004), [hep-ex/0404019](#)
291. R.R. Akhmetshin et al. (CMD-2), Phys. Lett. **B551**, 27 (2003), [hep-ex/0211004](#)
292. M.N. Achasov et al., J. Exp. Theor. Phys. **103**, 720 (2006), [hep-ex/0606057](#)
293. M.N. Achasov et al., Phys. Rev. **D76**, 072012 (2007), 0707.2279
294. D. Besson et al. (CLEO), Phys. Rev. **D76**, 072008 (2007), 0706.2813
295. D. Cronin-Hennessy et al. (CLEO) (2008), 0801.3418
296. A. Osterheld et al. (1986), sLAC-PUB-4160
297. J.Z. Bai et al. (BES), Phys. Rev. Lett. **88**, 101802 (2002), [hep-ex/0102003](#)
298. A.E. Blinov et al., Z. Phys. **C70**, 31 (1996)
299. R. Ammar et al. (CLEO), Phys. Rev. **D57**, 1350 (1998), [hep-ex/9707018](#)
300. J. Siegrist et al., Phys. Rev. **D26**, 969 (1982)
301. J.Z. Bai et al. (BES), Phys. Rev. Lett. **84**, 594 (2000), [hep-ex/9908046](#)
302. M. Ablikim et al. (BES), Phys. Lett. **B660**, 315 (2008), 0705.4500
303. M. Ablikim et al. (BES Collaboration), Phys. Lett. **B677**, 239 (2009), 0903.0900
304. Nucl. Phys. Proc. Suppl. **131**, 1 (2004), Workshop on Hadronic Cross Section at Low Energy SIGHAD03; Pisa; Oct. 8-10 2003
305. Nucl. Phys. Proc. Suppl. **144**, 1 (2005), 8th Int. Workshop on Tau Lepton Physics (Tau 04); Nara; Japan; 14-17 Sept. 2004
306. Nucl. Phys. Proc. Suppl. **162**, 1 (2006), Proc. of the Int. Workshop on e+ e- Collisions from phi to psi; Novosibirsk; Feb. 27 - March 2; 2006
307. Nucl. Phys. Proc. Suppl. **169**, 1 (2007), 9th Int. Workshop on Tau Lepton Physics; 19-22 September 2006; Pisa (Italy)
308. Nucl. Phys. Proc. Suppl. **181+182**, 1 (2008), Proc. of the Int. Workshop on e+ e- Collisions from Phi to Psi; Frascati 7-10 April 2008
309. Nucl. Phys. Proc. Suppl. **???**, 1 (2008), 10th Int. Workshop on Tau Lepton Physics; Novosibirsk; Russia Sept. 22-25, 2008
310. V.N. Baier, V.A. Khoze, Sov. Phys. JETP **21**, 629 (1965)
311. V.N. Baier, V.A. Khoze, Sov. Phys. JETP **21**, 1145 (1965)
312. G. Pancheri, Nuovo Cim. **A60**, 321 (1969)
313. M. Greco, G. Pancheri-Srivastava, Y. Srivastava, Nucl. Phys. **B101**, 234 (1975)
314. M.S. Chen, P.M. Zerwas, Phys. Rev. **D11**, 58 (1975)
315. S. Spagnolo, Eur. Phys. J. **C6**, 637 (1999)
316. V.A. Khoze et al., Eur. Phys. J. **C18**, 481 (2001), [hep-ph/0003313](#)
317. M. Benayoun, S.I. Eidelman, V.N. Ivanchenko, Z.K. Silagadze, Mod. Phys. Lett. **A14**, 2605 (1999), [hep-ph/9910523](#)
318. A.B. Arbuzov, E.A. Kuraev, N.P. Merenkov, L. Trentadue, JHEP **12**, 009 (1998), [hep-ph/9804430](#)
319. G. Abbiendi et al. (OPAL), Eur. Phys. J. **C33**, 173 (2004), [hep-ex/0309053](#)
320. J. Abdallah et al. (DELPHI), Eur. Phys. J. **C45**, 589 (2006), [hep-ex/0512012](#)
321. P. Achard et al. (L3), Eur. Phys. J. **C47**, 1 (2006), [hep-ex/0603022](#)
322. S. Schael et al. (ALEPH), Eur. Phys. J. **C49**, 411 (2007), [hep-ex/0609051](#)
323. S. Binner, J.H. Kühn, K. Melnikov, Phys. Lett. **B459**, 279 (1999), [hep-ph/9902399](#)
324. H. Czyż, J.H. Kühn, Eur. Phys. J. **C18**, 497 (2001), [hep-ph/0008262](#)
325. G. Rodrigo, A. Gehrmann-De Ridder, M. Guilleaume, J.H. Kühn, Eur. Phys. J. **C22**, 81 (2001), [hep-ph/0106132](#)
326. J.H. Kühn, G. Rodrigo, Eur. Phys. J. **C25**, 215 (2002), [hep-ph/0204283](#)
327. G. Rodrigo, H. Czyż, J.H. Kühn, M. Szopa, Eur. Phys. J. **C24**, 71 (2002), [hep-ph/0112184](#)
328. H. Czyż, A. Grzelińska, J.H. Kühn, G. Rodrigo, Eur. Phys. J. **C27**, 563 (2003), [hep-ph/0212225](#)
329. H. Czyż, A. Grzelińska, J.H. Kühn, G. Rodrigo, Eur. Phys. J. **C33**, 333 (2004), [hep-ph/0308312](#)



330. H. Czyż, A. Grzelińska, J.H. Kühn, G. Rodrigo, Eur. Phys. J. **C39**, 411 (2005), [hep-ph/0404078](#)
331. H. Czyż, A. Grzelińska, J.H. Kühn, Phys. Lett. **B611**, 116 (2005), [hep-ph/0412239](#)
332. H. Czyż, A. Grzelińska, J.H. Kühn, G. Rodrigo, Eur. Phys. J. **C47**, 617 (2006), [hep-ph/0512180](#)
333. H. Czyż, J.H. Kühn, A. Wapient, Phys. Rev. **D77**, 114005 (2008), [0804.0359](#)
334. H. Czyż, J.H. Kühn, E. Nowak, G. Rodrigo, Eur. Phys. J. **C35**, 527 (2004), [hep-ph/0403062](#)
335. H. Czyż, A. Grzelińska, J.H. Kühn, Phys. Rev. **D75**, 074026 (2007), [hep-ph/0702122](#)
336. H. Czyż, A. Grzelińska, A. Wapient, Acta Phys. Polon. **B38**, 3491 (2007), [0710.4227](#)
337. H. Czyż, Nucl. Phys. Proc. Suppl. **162**, 76 (2006), [hep-ph/0606227](#)
338. H. Czyż, Nucl. Phys. Proc. Suppl. **181-182**, 264 (2008)
339. A. Grzelińska, H. Czyż, A. Wapient (2008), [0812.1939](#)
340. M. Caffo, H. Czyż, E. Remiddi, Phys. Lett. **B327**, 369 (1994)
341. K. Melnikov, F. Nguyen, B. Valeriani, G. Venanzoni, Phys. Lett. **B477**, 114 (2000), [hep-ph/0001064](#)
342. S. Dubinsky, A. Korchin, N. Merenkov, G. Pancheri, O. Shekhovtsova, Eur. Phys. J. **C40**, 41 (2005), [hep-ph/0411113](#)
343. G. Pancheri, O. Shekhovtsova, G. Venanzoni, Phys. Lett. **B642**, 342 (2006), [hep-ph/0605244](#)
344. G. Pancheri, O. Shekhovtsova, G. Venanzoni, J. Exp. Theor. Phys. **106**, 470 (2008), [0706.3027](#)
345. O. Shekhovtsova, G. Venanzoni, G. Pancheri, Comput. Phys. Commun. **180**, 1206 (2009), [0901.4440](#)
346. G. Cataldi, A. Denig, S. Müller, W. Kluge, G. Venanzoni (1999), KLOE memo 195 (Aug. 1999), Frascati Physics Series (2000) 569
347. A. Denig et al. (KLOE) (2001), [hep-ex/0106100](#)
348. A. Aloisio et al. (KLOE) (2001), [hep-ex/0107023](#)
349. A.G. Denig et al. (the KLOE), Nucl. Phys. Proc. Suppl. **116**, 243 (2003), [hep-ex/0211024](#)
350. B. Valeriani et al. (KLOE) (2002), [hep-ex/0205046](#)
351. G. Venanzoni et al. (KLOE), Nucl. Phys. Proc. Suppl. **123**, 177 (2003), [hep-ex/0210013](#)
352. S.E. Müller (KLOE), Nucl. Phys. Proc. Suppl. **126**, 335 (2004)
353. A. Aloisio et al. (KLOE), Eur. Phys. J. **C33**, s656 (2004), [hep-ex/0307051](#)
354. A.G. Denig (KLOE), AIP Conf. Proc. **717**, 83 (2004), [hep-ex/0311012](#)
355. B. Valeriani (KLOE), Nucl. Phys. Proc. Suppl. **131**, 75 (2004)
356. W. Kluge, Nucl. Phys. Proc. Suppl. **135**, 357 (2004)
357. A.G. Denig (KLOE), Int. J. Mod. Phys. **A20**, 1935 (2005)
358. W. Kluge (KLOE), Acta Physica Slovaca **55**, 49 (2005)
359. A. Aloisio et al. (KLOE), Nucl. Phys. Proc. Suppl. **144**, 231 (2005)
360. A. Denig, Nucl. Phys. Proc. Suppl. **162**, 81 (2006), [hep-ex/0611024](#)
361. S.E. Müller, F. Nguyen (KLOE), Nucl. Phys. Proc. Suppl. **162**, 90 (2006)
362. D. Leone (KLOE), Nucl. Phys. Proc. Suppl. **162**, 95 (2006)
363. G. Venanzoni (KLOE), Nucl. Phys. Proc. Suppl. **169**, 277 (2007)
364. F. Nguyen (KLOE), Nucl. Phys. Proc. Suppl. **181-182**, 106 (2008), [0807.1612](#)
365. S.E. Müller (KLOE), Acta Phys. Polon. **B38**, 3007 (2007)
366. F. Ambrosino et al. (KLOE) (2007), [0707.4078](#)
367. A. Aloisio et al. (KLOE), Phys. Lett. **B606**, 12 (2005), [hep-ex/0407048](#)
368. F. Ambrosino et al. (KLOE), Phys. Lett. **B670**, 285 (2009), [0809.3950](#)
369. W. Kluge, Nucl. Phys. Proc. Suppl. **181-182**, 280 (2008), [0805.4708](#)
370. G. Venanzoni (for the KLOE Collaboration) (2009), [0906.4331](#)
371. G.W. Bennett et al. (Muon g-2), Phys. Rev. Lett. **89**, 101804 (2002), [hep-ex/0208001](#)
372. G.W. Bennett et al. (Muon g-2), Phys. Rev. Lett. **92**, 161802 (2004), [hep-ex/0401008](#)
373. D.W. Hertzog, Nucl. Phys. Proc. Suppl. **181-182**, 5 (2008)
374. F. Jegerlehner, *The anomalous magnetic moment of the muon*, Berlin, Germany: Springer 426 p (2008)
375. M. Davier, W.J. Marciano, Ann. Rev. Nucl. Part. Sci. **54**, 115 (2004)
376. T. Teubner, Nucl. Phys. Proc. Suppl. **181-182**, 20 (2008)
377. D. Stockinger, Nucl. Phys. Proc. Suppl. **181-182**, 32 (2008)
378. J.H. Kühn, M. Steinhauser, Phys. Lett. **B437**, 425 (1998), [hep-ph/9802241](#)
379. S. Eidelman, F. Jegerlehner, A.L. Kataev, O. Veretin, Phys. Lett. **B454**, 369 (1999), [hep-ph/9812521](#)
380. K.G. Chetyrkin, J.H. Kühn, A. Kwiatkowski, Phys. Rept. **277**, 189 (1996)
381. R.R. Akhmetshin et al. (CMD-2), Phys. Lett. **B562**, 173 (2003), [hep-ex/0304009](#)
382. V.M. Aulchenko et al. (CMD-2), JETP Lett. **82**, 743 (2005), [hep-ex/0603021](#)
383. R.R. Akhmetshin et al. (CMD2), Phys. Lett. **B605**, 26 (2005), [hep-ex/0409030](#)
384. R.R. Akhmetshin et al., JETP Lett. **84**, 413 (2006), [hep-ex/0610016](#)
385. F. Ignatov (CMD-2 and SND), Nucl. Phys. Proc. Suppl. **181-182**, 101 (2008)
386. R.R. Akhmetshin et al. (CMD-2), Phys. Lett. **B648**, 28 (2007), [hep-ex/0610021](#)
387. M.N. Achasov et al., Nucl. Phys. Proc. Suppl. **162**, 11 (2006), [hep-ex/0604052](#)
388. E.P. Solodov (BABAR) (2002), [hep-ex/0107027](#)
389. B. Aubert et al. (BABAR), Phys. Rev. **D69**, 011103 (2004), [hep-ex/0310027](#)
390. B. Aubert et al. (BABAR), Phys. Rev. **D70**, 072004 (2004), [hep-ex/0408078](#)
391. B. Aubert et al. (BABAR), Phys. Rev. **D71**, 052001 (2005), [hep-ex/0502025](#)
392. B. Aubert et al. (BABAR), Phys. Rev. **D73**, 012005 (2006), [hep-ex/0512023](#)
393. B. Aubert et al. (BABAR), Phys. Rev. **D73**, 052003 (2006), [hep-ex/0602006](#)
394. B. Aubert et al. (BABAR), Phys. Rev. **D74**, 091103 (2006), [hep-ex/0610018](#)
395. B. Aubert et al. (BABAR), Phys. Rev. **D76**, 012008 (2007), [0704.0630](#)
396. B. Aubert et al. (BaBar), Phys. Rev. **D77**, 092002 (2008), [0710.4451](#)



397. B. Aubert et al. (BABAR), Phys. Rev. **D76**, 092005 (2007), 0708.2461
398. B. Aubert et al. (BABAR), Phys. Rev. **D76**, 092006 (2007), 0709.1988
399. B. Aubert et al. (BABAR), Phys. Rev. **D76**, 111105 (2007), hep-ex/0607083
400. B. Aubert et al. (BaBar) (2008), 0808.1543
401. B. Aubert et al. (BABAR) (2009), 0903.1597
402. A.G. Denig, P.A. Lukin (BABAR), Nucl. Phys. Proc. Suppl. **181-182**, 111 (2008)
403. M. Davier (2009), to be published in Nucl. Phys. **B**, Proc. Suppl.(2009), <http://tau08.inp.nsk.su/prog.php>
404. K. Abe et al. (Belle), Phys. Rev. Lett. **98**, 092001 (2007), hep-ex/0608018
405. C.Z. Yuan et al. (Belle), Phys. Rev. Lett. **99**, 182004 (2007), 0707.2541
406. X.L. Wang et al. (Belle), Phys. Rev. Lett. **99**, 142002 (2007), 0707.3699
407. G. Pakhlova et al. (Belle), Phys. Rev. **D77**, 011103 (2008), 0708.0082
408. C.Z. Yuan et al. (Belle), Phys. Rev. **D77**, 011105 (2008), 0709.2565
409. G. Pakhlova, Nucl. Phys. Proc. Suppl. **181-182**, 117 (2008)
410. G. Pakhlova et al. (Belle), Phys. Rev. Lett. **100**, 062001 (2008), 0708.3313
411. G. Pakhlova et al. (Belle), Phys. Rev. Lett. **101**, 172001 (2008), 0807.4458
412. B. Aubert et al. (BABAR), Phys. Rev. Lett. **95**, 142001 (2005), hep-ex/0506081
413. E. Prencipe, Nucl. Phys. Proc. Suppl. **181-182**, 333 (2008)
414. K. Abe et al. (Belle) (2006), hep-ex/0612006
415. V. Balagura (Belle), Nucl. Phys. Proc. Suppl. **181-182**, 338 (2008)
416. T.E. Coan et al. (CLEO), Phys. Rev. Lett. **96**, 162003 (2006), hep-ex/0602034
417. Q. He et al. (CLEO), Phys. Rev. **D74**, 091104 (2006), hep-ex/0611021
418. Y.S. Kalashnikova, Nucl. Phys. Proc. Suppl. **181-182**, 363 (2008)
419. F.E. Maas (PANDA), Nucl. Phys. Proc. Suppl. **181-182**, 45 (2008)
420. G. Salme, Nucl. Phys. Proc. Suppl. **181-182**, 51 (2008), 0807.0344
421. V.F. Dmitriev, A.I. Milstein, Nucl. Phys. Proc. Suppl. **181-182**, 66 (2008)
422. R. Baldini, S. Pacetti, A. Zallo, A. Zichichi, Eur. Phys. J. **A39**, 315 (2009), 0711.1725
423. J. Arrington, Phys. Rev. **C68**, 034325 (2003), nucl-ex/0305009
424. G. Rodrigo, Acta Phys. Polon. **B32**, 3833 (2001), hep-ph/0111151
425. M. Caffo, H. Czyż, E. Remiddi, Nuovo Cim. **A110**, 515 (1997), hep-ph/9704443
426. F.A. Berends, W.L. van Neerven, G.J.H. Burgers, Nucl. Phys. **B297**, 429 (1988)
427. J.S. Schwinger (1989), Particles, Sources, and Fields. Vol. 3, Redwood City, USA: Addison-Wesley (1989), p.99
428. F. Jegerlehner, K. Kolodziej, Eur. Phys. J. **C12**, 77 (2000), hep-ph/9907229
429. K. Kolodziej, M. Zralek, Phys. Rev. **D43**, 3619 (1991)
430. R. Tarrach, Nuovo Cim. **A28**, 409 (1975)
431. D. Drechsel, G. Knochlein, A. Metz, S. Scherer, Phys. Rev. **C55**, 424 (1997), nucl-th/9608061
432. G. Ecker, J. Gasser, H. Leutwyler, A. Pich, E. de Rafael, Phys. Lett. **B223**, 425 (1989)
433. G. Ecker, J. Gasser, A. Pich, E. de Rafael, Nucl. Phys. **B321**, 311 (1989)
434. N.N. Achasov, V.V. Gubin, E.P. Solodov, Phys. Rev. **D55**, 2672 (1997), hep-ph/9610282
435. N.N. Achasov, A.V. Kiselev, Phys. Rev. **D73**, 054029 (2006), hep-ph/0512047
436. O. Shekhovtsova, PHOKHARA6.1, unpublished (2008), <http://ific.uv.es/rodrigo/phokhara/>
437. F. Ambrosino et al. (KLOE), Eur. Phys. J. **C49**, 473 (2007), hep-ex/0609009
438. P. Beltrame, *Measurement of the pion form factor via Radiative Return for  $\sqrt{s} = 1$  GeV with the KLOE detector*, PhD thesis, KA-IEKP-2009-8 (2009)
439. D. Leone, *Measurement of the hadronic cross section  $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$  with the KLOE detector using radiative return with tagged photons*, PhD thesis, KA-IEKP-2007-7 (2007)
440. C. Bruch, A. Khodjamirian, J.H. Kuhn, Eur. Phys. J. **C39**, 41 (2005), hep-ph/0409080
441. J.H. Kuhn, A. Santamaria, Z. Phys. **C48**, 445 (1990)
442. F. Ambrosino et al. (KLOE), Phys. Lett. **B670**, 285 (2009), 0809.3950
443. F. Jegerlehner, Fortran code for the effective fine structure constant. <http://www-com.physik.hu-berlin.de/fjeger/alphaQED.uu>
444. H. Czyż et al., unpublished
445. S. Mueller et al., <http://www.lnf.infn.it/kloe/pub/knote/kn221.pdf>
446. F. Ambrosino et al., Nucl. Instrum. Meth. **A534**, 403 (2004), physics/0404100
447. A. Denig et al., <http://www.lnf.infn.it/kloe/pub/knote/kn192.ps>
448. A. Bramon, G. Colangelo, P.J. Franzini, M. Greco, Phys. Lett. **B287**, 263 (1992)
449. J.L. Lucio Martinez, M. Napsuciale, Phys. Lett. **B331**, 418 (1994)
450. B. Aubert (The BABAR) (2009), 0908.3589
451. A. Antonelli et al. (DM2), Z. Phys. **C56**, 15 (1992)
452. M.N. Achasov et al. (SND), Preprint Budker INP 2001-34 (2001)
453. P.L. Frabetti et al. (E687), Phys. Lett. **B514**, 240 (2001), hep-ex/0106029
454. G. Bardin et al., Nucl. Phys. **B411**, 3 (1994)
455. D. Bisello et al. (DM2), Z. Phys. **C48**, 23 (1990)
456. E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane, T.M. Yan, Phys. Rev. **D21**, 203 (1980)
457. G. Pakhlova (Belle) (2009), 0908.0231
458. K. Abe et al. (Belle), Phys. Rev. Lett. **88**, 181803 (2002), hep-ex/0202017
459. A.M. Badalian, B.L.G. Bakker, I.V. Danilkin, Phys. Atom. Nucl. **72**, 638 (2009), 0805.2291
460. B.Q. Li, K.T. Chao, Phys. Rev. D **79**, 094004 (2009), 0903.5506
461. E. van Beveren, X. Liu, R. Coimbra, G. Rupp (2008), 0809.1151
462. I. Adachi et al. (Belle) (2008), 0808.0006
463. L. Wang, Ph.D. thesis, Institute of High Energy Physics, Chinese Academy of Sciences, and Laboratoire de l'Accelérateur Lineaire, Université Paris-Sud 11 (2009)

464. S. Dobbs et al. (CLEO), Phys. Rev. **D76**, 112001 (2007), 0709.3783
465. S. Jadach, Z. Was, Comput. Phys. Commun. **36**, 191 (1985)
466. S. Jadach, Z. Was, B.F.L. Ward, Comput. Phys. Commun. **130**, 260 (2000), up to date source available from <http://home.cern.ch/jadach/>
467. S. Jadach, J.H. Kuhn, Z. Was, Comput. Phys. Commun. **64**, 275 (1990)
468. M. Jezabek, Z. Was, S. Jadach, J.H. Kuhn, Comput. Phys. Commun. **70**, 69 (1992)
469. S. Jadach, Z. Was, R. Decker, J.H. Kuhn, Comput. Phys. Commun. **76**, 361 (1993)
470. E. Barberio, Z. Was, Comput. Phys. Commun. **79**, 291 (1994)
471. J.H. Kuhn, E. Mirkes, Z. Phys. **C56**, 661 (1992)
472. A.E. Bondar et al., Comput. Phys. Commun. **146**, 139 (2002), [hep-ph/0201149](#)
473. P. Golonka et al., Comput. Phys. Commun. **174**, 818 (2006), [hep-ph/0312240](#)
474. M. Fujikawa et al. (Belle), Phys. Rev. **D78**, 072006 (2008), 0805.3773
475. B. Aubert et al. (BaBar), Phys. Rev. **D76**, 051104 (2007), 0707.2922
476. D. Epifanov et al. (Belle), Phys. Lett. **B654**, 65 (2007), 0706.2231
477. B. Aubert et al. (BaBar), Phys. Rev. Lett. **100**, 011801 (2008), 0707.2981
478. I. Adachi et al. (Belle) (2008), 0812.0480
479. K. Inami et al. (Belle), Phys. Lett. **B643**, 5 (2006), [hep-ex/0609018](#)
480. D.M. Asner et al. (CLEO), Phys. Rev. **D61**, 012002 (2000), [hep-ex/9902022](#)
481. K. Inami et al. (Belle), Phys. Lett. **B672**, 209 (2009), 0811.0088
482. D. Buskulic et al. (ALEPH), Z. Phys. **C74**, 263 (1997)
483. K.W. Edwards et al. (CLEO), Phys. Rev. **D61**, 072003 (2000), [hep-ex/9908024](#)
484. R. Decker, M. Finkemeier, P. Heiliger, H.H. Jonsson, Z. Phys. **C70**, 247 (1996), [hep-ph/9410260](#)
485. B. Aubert et al. (BaBar), Phys. Rev. **D72**, 072001 (2005), [hep-ex/0505004](#)
486. D. Gibaut et al. (CLEO), Phys. Rev. Lett. **73**, 934 (1994)
487. J.H. Kuhn, Z. Was, Acta Phys. Polon. **B39**, 147 (2008), [hep-ph/0602162](#)
488. A. Anastassov et al. (CLEO), Phys. Rev. Lett. **86**, 4467 (2001), [hep-ex/0010025](#)
489. R. Kitano, Y. Okada, Phys. Rev. **D63**, 113003 (2001), [hep-ph/0012040](#)
490. B.M. Dassinger, T. Feldmann, T. Mannel, S. Turczyk, JHEP **10**, 039 (2007), 0707.0988
491. D.R. Yennie, S. Frautschi, H. Suura, Ann. Phys. (NY) **13**, 379 (1961)
492. G. Nanava, Z. Was, Eur. Phys. J. **C51**, 569 (2007), [hep-ph/0607019](#)
493. P. Golonka, Z. Was, Eur. Phys. J. **C45**, 97 (2006), [hep-ph/0506026](#)
494. P. Golonka, Z. Was, Eur. Phys. J. **C50**, 53 (2007), [hep-ph/0604232](#)
495. A. Einstein, B. Podolsky, N. Rosen, Phys. Rev. **47**, 777 (1935)
496. S. Jadach, Z. Was, Comput. Phys. Commun. **36**, 191 (1985)
497. S. Jadach, Z. Was, Acta Phys. Polon. **B15**, 1151 (1984)
498. S. Jadach, E. Richter-Was, B.F.L. Ward, Z. Was, Phys. Lett. **B260**, 438 (1991)
499. S. Banerjee, B. Pietrzyk, J.M. Roney, Z. Was, Phys. Rev. **D77**, 054012 (2008), 0706.3235
500. E. Barberio, B. van Eijk, Z. Was, Comput. Phys. Commun. **66**, 115 (1991)
501. E. Barberio, Z. Was, Comput. Phys. Commun. **79**, 291 (1994)
502. Z. Was, Acta Phys. Polon. **B39**, 1761 (2008), 0807.2775
503. P. Golonka, G. Nanava, Z. Was, *Tests of photos hard bremsstrahlung* (2002-now), tests of PHOTOS Hard Bremsstrahlung, <http://mc-tester.web.cern.ch/MC-TESTER/PHOTOS-MCTESTER/>
504. Z. Was, Q. Xu (2008), in preparation
505. A. van Hameren, Z. Was (2008), 0802.2182
506. Z. Was, Nucl. Instrum. Meth. **A534**, 260 (2004), [hep-ph/0402129](#)
507. D. Gomez Dumm, A. Pich, J. Portoles, Phys. Rev. **D69**, 073002 (2004), [hep-ph/0312183](#)
508. J. Gasser, H. Leutwyler, Ann. Phys. **158**, 142 (1984)
509. J. Gasser, H. Leutwyler, Nucl. Phys. **B250**, 465 (1985)
510. D. Gómez-Dumm, P. Roig, A. Pich, J. Portolés, to appear
511. S.J. Brodsky, G.R. Farrar, Phys. Rev. Lett. **31**, 1153 (1973)
512. G.P. Lepage, S.J. Brodsky, Phys. Rev. **D22**, 2157 (1980)
513. D. Gomez Dumm, A. Pich, J. Portoles, Phys. Rev. **D62**, 054014 (2000), [hep-ph/0003320](#)
514. F. Guerrero, A. Pich, Phys. Lett. **B412**, 382 (1997), [hep-ph/9707347](#)
515. A. Pich, J. Portoles, Phys. Rev. **D63**, 093005 (2001), [hep-ph/0101194](#)
516. A. Pich, J. Portoles, Nucl. Phys. Proc. Suppl. **121**, 179 (2003), [hep-ph/0209224](#)
517. M. Jamin, A. Pich, J. Portoles, Phys. Lett. **B640**, 176 (2006), [hep-ph/0605096](#)
518. M. Jamin, A. Pich, J. Portoles, Phys. Lett. **B664**, 78 (2008), 0803.1786
519. P. Roig, Nucl. Phys. B (Proc. Suppl.) **189**, 78 (2009), 0810.5764
520. R. Barate et al. (ALEPH), Eur. Phys. J. **C4**, 409 (1998)
521. Y.S. Tsai, Phys. Rev. **D4**, 2821 (1971)
522. H.B. Thacker, J.J. Sakurai, Phys. Lett. **B36**, 103 (1971)
523. S.I. Eidelman, V.N. Ivanchenko, Phys. Lett. **B257**, 437 (1991)
524. V. Cirigliano, G. Ecker, H. Neufeld, Phys. Lett. **B513**, 361 (2001), [hep-ph/0104267](#)
525. V. Cirigliano, G. Ecker, H. Neufeld, JHEP **08**, 002 (2002), [hep-ph/0207310](#)
526. S. Anderson et al. (CLEO), Phys. Rev. **D61**, 112002 (2000), [hep-ex/9910046](#)
527. S. Schael et al. (ALEPH), Phys. Rept. **421**, 191 (2005), [hep-ex/0506072](#)
528. K. Ackerstaff et al. (OPAL), Eur. Phys. J. **C7**, 571 (1999), [hep-ex/9808019](#)
529. S. Ghozzi, F. Jegerlehner, Phys. Lett. **B583**, 222 (2004), [hep-ph/0310181](#)
530. M. Davier, A. Hocker, Z. Zhang, Rev. Mod. Phys. **78**, 1043 (2006), [hep-ph/0507078](#)
531. V.P. Druzhinin (2007), 0710.3455
532. A. Korchin et al., preliminary
533. N. Davidson, P. Golonka, T. Przedzinski, Z. Was (2008), 0812.3215

534. R. Brun et al., J. Phys. Conf. Ser. **119**, 042006 (2008), 0901.0886
535. O. Shekhovtsova, A. Kalinowski, T. Przedzinski, *preliminary* (2009)
536. A. Korchin, J. Fujimoto, H. Czyz, F. Jegerlehner, *preliminary* (2009)
537. A. Hocker et al., PoS **ACAT**, 040 (2007), physics/0703039
538. V. Cherepanov, T. Przedzinski, Z. Was, *preliminary* (2009), resulting software is available upon request.
539. Z. Was, Nucl. Phys. B (Proc. Suppl.) **189**, 43 (2009), 0901.1865
540. G.R. Bower, T. Pierzchala, Z. Was, M. Worek, Phys. Lett. **B543**, 227 (2002), hep-ph/0204292
541. K. Desch, A. Imhof, Z. Was, M. Worek, Phys. Lett. **B579**, 157 (2004), hep-ph/0307331
542. P. Privitera, Phys. Lett. **B308**, 163 (1993)
543. J.H. Kuhn, Phys. Rev. **D52**, 3128 (1995), hep-ph/9505303
544. E. Richter-Was (Atlas) (2009), 0903.4198
545. G.L. Bayatian et al. (CMS), *CMS physics: Technical design report*, CERN-LHCC-2006-001 (2006)
546. T. Nattermann, K. Desch, P. Wienemann, C. Zendler, JHEP **04**, 057 (2009), 0903.0714
547. G. Aad et al. (ATLAS), JINST **3**, S08003 (2008)
548. G. Aad et al. (ATLAS) (2009), 0901.0512
549. A. Heister et al. (ALEPH), Eur. Phys. J. **C20**, 401 (2001), hep-ex/0104038
550. T. Aoyama, M. Hayakawa, T. Kinoshita, M. Nio, Phys. Rev. **D77**, 053012 (2008), 0712.2607
551. K. Hagiwara, R. Liao, A.D. Martin, D. Nomura, T. Teubner (2009), in preparation
552. A.O.G. Kallen, A. Sabry, Kong. Dan. Vid. Sel. Mat. Fys. Med. **29N17**, 1 (1955)
553. M. Steinhauser, Phys. Lett. **B429**, 158 (1998), hep-ph/9803313
554. A. Djouadi, P. Gambino, Phys. Rev. **D49**, 3499 (1994), hep-ph/9309298
555. A. Djouadi, P. Gambino, Erratum Phys. Rev. **D53**, 4111 (1996)
556. H. Burkhardt (1982), thesis: DESY Hamburg - Internal Rep. F35-82-03, 118p
557. H. Burkhardt, B. Pietrzyk, Phys. Lett. **B356**, 398 (1995)